



Scott and White Health Plan

SeniorCare Advantage (PPO)
SeniorCare Advantage (HMO)

2020 Prior Authorization Criteria

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

ACTEMRA IV (S)

Products Affected

- Actemra INJ 200MG/10ML, 400MG/20ML, 80MG/4ML

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Rheumatoid Arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. One of the following: Trial and failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR for continuation of prior Actemra therapy. Systemic Juvenile Idiopathic arthritis (SJIA) (Initial): Diagnosis of active SJIA. Trial and failure, contraindication, or intolerance to one NSAID or systemic glucocorticoid. Polyarticular Juvenile Idiopathic Arthritis (PJIA) (Initial): Diagnosis of active PJIA. One of the following: Trial and failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR for continuation of prior Actemra therapy. Cytokine Release Syndrome (CRS) Risk due to CAR T-cell Therapy: Patient will receive or is receiving chimeric antigen receptor (CAR) T-cell immunotherapy [i.e., Kymriah (tisagenlecleuce), Yescarta (axicabtagene ciloleuce)]. |
| Age Restrictions | N/A |
| Prescriber Restrictions | RA, SJIA, PJIA (initial): Prescribed by or in consultation with a rheumatologist. CRS Risk due to CAR T-cell Therapy: Prescribed by or in consultation with an oncologist or hematologist. |
| Coverage Duration | RA, SJIA, PJIA (Initial, reauth): 12 months. CRS risk due to CAR T-cell therapy: 2 months |
| Other Criteria | RA, SJIA, PJIA (Reauth): Documentation of positive clinical response to Actemra therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

ACTEMRA SC (S)

Products Affected

- Actemra INJ 162MG/0.9ML
- Actemra Actpen

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Rheumatoid Arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. One of the following: Trial and failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR for continuation of prior Actemra therapy. Giant Cell Arteritis (GCA) (Initial): Diagnosis of GCA. Trial and failure, contraindication, or intolerance to a glucocorticoid (i.e., prednisone). Systemic Juvenile Idiopathic Arthritis (SJIA) (Initial): Diagnosis of active SJIA. Trial and failure, contraindication, or intolerance to one of the following: NSAID or systemic glucocorticoid. Polyarticular Juvenile Idiopathic Arthritis (PJIA) (Initial): Diagnosis of active PJIA. One of the following: Trial and failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR for continuation of prior Actemra therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | RA, GC, SJIA, PJIA (initial): Prescribed by or in consultation with a rheumatologist. |
| Coverage Duration | RA, GC, SJIA, PJIA (initial, reauth): 12 months |
| Other Criteria | RA, GC, SJIA, PJIA (Reauth): Documentation of positive clinical response to Actemra therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ACTIMMUNE (S)

Products Affected

- Actimmune

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Diagnosis of one of the following: 1) Chronic granulomatous disease (CGD), or 2) severe malignant osteopetrosis (SMO). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ADCETRIS (S)

Products Affected

- Adcetris

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Hodgkin Lymphoma (HL): Diagnosis of HL. One of the following: failure of autologous hematopoietic stem cell transplant (auto-HSCT) or failure of at least two prior multi-agent chemotherapy regimens or used as post-auto-HSCT consolidation therapy for patients at high risk of relapse or progression. Systemic Anaplastic Large Cell Lymphoma (sALCL): Diagnosis of sALCL. Failure of at least one prior multi-agent chemotherapy regimen. |
| Age Restrictions | N/A |
| Prescriber Restrictions | HL, sALCL: Prescribed by or in consultation with an oncologist/hematologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ADCIRCA (S)

Products Affected

- Alyq
- Tadalafil TABS 20MG

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH. |
| Age Restrictions | N/A |
| Prescriber Restrictions | PAH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist. |
| Coverage Duration | PAH: Initial: 6 months. Reauth: 12 months. |
| Other Criteria | PAH (Reauth): Documentation of positive clinical response to therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ADDERALL XR (S)

Products Affected

- Amphetamine/dextroamphetamine
CP24

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | One of the following: a) diagnosis of attention deficit hyperactivity disorder (ADHD), OR b) diagnosis of attention deficit disorder (ADD) |
| Age Restrictions | PA applies to members 19 years of age or older |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

ADEMPAS (S)

Products Affected

- Adempas

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH AND PAH is symptomatic AND One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH. Chronic thromboembolic pulmonary hypertension (CTEPH) (Initial): One of the following: A) Both of the following: 1) Diagnosis of inoperable or persistent/recurrent CTEPH and 2) CTEPH is symptomatic OR B) Patient is currently on any therapy for the diagnosis of CTEPH. |
| Age Restrictions | N/A |
| Prescriber Restrictions | PAH, CTEPH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist. |
| Coverage Duration | PAH, CTEPH: Initial: 6 months. Reauth: 12 months. |
| Other Criteria | PAH, CTEPH (Reauth): Documentation of positive clinical response to therapy. |

AFINITOR (S)

Products Affected

- Afinitor
- Everolimus TABS 2.5MG, 5MG, 7.5MG

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Subependymal Giant Cell Astrocytoma (SEGA) associated with tuberous sclerosis complex (TSC): Diagnosis of SEGA associated with TSC that requires therapeutic intervention but patient is not a candidate for curative surgical resection. Renal cell carcinoma: Diagnosis of advanced or metastatic renal cell carcinoma AND trial and failure, contraindication, or intolerance to SUTENT (sunitinib) or NEXAVAR (sorafenib). Neuroendocrine tumors of pancreatic origin (pNET): Diagnosis of progressive pNET that are unresectable, locally advanced, or metastatic. Renal angiomyolipoma: Diagnosis of renal angiomyolipoma and TSC AND Patient does not require immediate surgery. Breast Cancer: Patient is a postmenopausal woman AND Diagnosis of advanced hormone receptor-positive, HER2-negative breast cancer AND trial and failure, contraindication, or intolerance to FEMARA (letrozole) or ARIMIDEX (anastrozole) AND used in combination with AROMASIN (exemestane). Neuroendocrine tumors of gastrointestinal (GI) or lung origin: Diagnosis of progressive, well-differentiated, non-functional NET of GI or lung origin AND patient has unresectable, locally advanced or metastatic disease. |
| Age Restrictions | N/A |
| Prescriber Restrictions | All uses: Prescribed by or in consultation with an oncologist |
| Coverage Duration | All uses: 12 months |
| Other Criteria | All Indications: Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

AFINITOR DISPERZ (S)

Products Affected

- Afinitor Disperz

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Subependymal Giant Cell Astrocytoma (SEGA): Diagnosis of SEGA associated with tuberous sclerosis complex (TSC) that requires therapeutic intervention but patient is not a candidate for curative surgical resection. TSC-associated partial-onset seizures: Diagnosis of TSC-associated partial-onset seizures. Used as adjunctive therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | SEGA: Prescribed by or in consultation with an oncologist. TSC-associated partial-onset seizures: Prescribed by or in consultation with a neurologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

AIMOVIG (S)

Products Affected

- Aimovig

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Episodic Migraines (EM) (initial): Diagnosis of EM. Patient has 4 to 14 migraine days per month, but no more than 14 headache days per month. Chronic Migraines (CM) (initial): Diagnosis of CM. Medication overuse headache has been considered and potentially offending medication(s) have been discontinued. Patient has greater than or equal to 15 headache days per month, of which at least 8 must be migraine days for at least 3 months. |
| Age Restrictions | EM, CM (initial): 18 years of age or older. |
| Prescriber Restrictions | EM, CM (initial, reauth): Prescribed by or in consultation with a neurologist or pain specialist. |
| Coverage Duration | EM, CM (initial): 6 months. EM, CM (reauth): 12 months. |
| Other Criteria | EM, CM (reauth): Patient has experienced a positive response to therapy, demonstrated by a reduction in headache frequency and/or intensity. Use of acute migraine medications (e.g., non-steroidal anti-inflammatory drugs [NSAIDs], triptans) has decreased since the start of CGRP therapy. CM (reauth): Patient continues to be monitored for medication overuse headache. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ALDURAZYME (S)

Products Affected

- Aldurazyme

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Mucopolysaccharidosis I: confirmed diagnosis of Hurler and Hurler-Scheie forms of Mucopolysaccharidosis I (MPS I), OR confirmed diagnosis of Scheie form of Mucopolysaccharidosis I (MPS I) who have moderate to severe symptoms. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ALECENSA (S)

Products Affected

- Alecensa

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Non-small cell lung cancer (NSCLC): A) Diagnosis of metastatic NSCLC AND B) Patient has anaplastic lymphoma kinase (ALK)-positive disease as detected with a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ALIQOPA (S)

Products Affected

- Aliqopa

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Relapsed Follicular Lymphoma: Diagnosis of relapsed follicular lymphoma AND patient has received at least two prior systemic therapies. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist/oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

ALPHA-1 PROTEINASE INHIBITOR, NON-PREFERRED (S)

Products Affected

- Aralast Np
- Glassia
- Zemaira

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Alpha-1 antitrypsin (AAT) deficiency: Diagnosis of congenital AAT deficiency. Diagnosis of emphysema. Continued optimal conventional treatment for emphysema (e.g., bronchodilators). One of the following: 1) PiZZ, PiZ(null), or Pi(null)(null) protein phenotypes (homozygous) OR 2) other rare AAT disease-causing alleles associated with serum AAT level less than 11 µM/L [e.g., Pi(Malton, Malton)]. Circulating serum concentration of AAT level less than 11 µM/L (which corresponds to less than 80 mg/dL if measured by radial immunodiffusion or less than 57 mg/dL if measured by nephelometry). One of the following: FEV1 level is between 30% and 65% of predicted OR the patient has experienced a rapid decline in lung function (i.e., reduction of FEV1 more than 120 mL/year) that warrants treatment. Trial and failure, or intolerance to Prolastin. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

ALPHA-1 PROTEINASE INHIBITOR, PROLASTIN (S)

Products Affected

- Prolastin-c INJ 1000MG

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Alpha-1 antitrypsin (AAT) deficiency: Diagnosis of congenital AAT deficiency. Diagnosis of emphysema. Continued optimal conventional treatment for emphysema (e.g., bronchodilators). One of the following: 1) PiZZ, PiZ(null), or Pi(null)(null) protein phenotypes (homozygous) OR 2) other rare AAT disease-causing alleles associated with serum AAT level less than 11 µM/L [e.g., Pi(Malton, Malton)]. Circulating serum concentration of AAT level less than 11 µM/L (which corresponds to less than 80 mg/dL if measured by radial immunodiffusion or less than 57 mg/dL if measured by nephelometry). One of the following: FEV1 level is between 30% and 65% of predicted OR the patient has experienced a rapid decline in lung function (i.e., reduction of FEV1 more than 120 mL/year) that warrants treatment. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ALUNBRIG (S)

Products Affected

- Alunbrig

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Non-small cell lung cancer (NSCLC): Diagnosis of metastatic NSCLC. Patient has an anaplastic lymphoma kinase (ALK)-positive tumor as detected with a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

AMPHETAMINE (S)

Products Affected

- Amphetamine/dextroamphetamine
TABS

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | One of the following: a) diagnosis of ADHD, OR b) diagnosis of attention deficit disorder (ADD), OR c) diagnosis of narcolepsy as confirmed by a sleep study (unless the prescriber provides justification confirming that a sleep study would not be feasible). |
| Age Restrictions | PA applies to members 19 years of age or older |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

AMPYRA (S)

Products Affected

- Ampyra

- Dalfampridine Er

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Multiple Sclerosis (MS) (initial): Diagnosis of MS. Physician confirmation that patient has difficulty walking (eg, timed 25 foot walk test). One of the following: expanded disability status scale (EDSS) score less than or equal to 7, or not restricted to using a wheelchair (if EDSS is not measured). |
| Age Restrictions | N/A |
| Prescriber Restrictions | MS (initial): Prescribed by or in consultation with a neurologist. |
| Coverage Duration | MS (Initial): 6 months. (Reauth): 12 months. |
| Other Criteria | MS (Reauth): Physician confirmation that the patient's walking improved with therapy. One of the following: EDSS score less than or equal to 7, or not restricted to using a wheelchair (if EDSS is not measured). |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ANADROL-50 (S)

Products Affected

- Anadrol-50

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Anemia (initial): Diagnosis of anemia caused by deficient red cell production AND trial and failure or intolerance to two standard therapies for anemia (i.e., erythropoiesis-stimulating agents, immunosuppressants) AND Treatment will not replace other supportive measures (e.g., transfusion, correction of iron, folic acid, vitamin B12 or pyridoxine deficiency, antibacterial therapy, corticosteroids). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | Initial, reauth: 12 months |
| Other Criteria | Anemia (reauth): Documentation of a positive clinical response to Anadrol-50 therapy as evidenced by an improvement in anemia (e.g., increased hemoglobin, increased reticulocyte count, reduction/elimination for need of blood transfusions). |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

ANDROXY (S)

Products Affected

- Androxy

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Hypogonadism (HG) (Initial): Diagnosis (dx) of HG in a male patient at birth. 1) Two pre-treatment serum total testosterone (T) levels less than 300 ng/dL (10.4 nmol/L) or less than the reference range for the lab OR 2) Both of the following: Has a condition that may cause altered sex-hormone binding globulin (SHBG) (eg, thyroid disorder, HIV disease, liver disorder, diabetes, obesity), and one pre-treatment calculated free or bioavailable T level less than 5 ng/dL (0.17 nmol/L) or less than reference range for the lab, OR 3) History of bilateral orchiectomy, panhypopituitarism, or a genetic disorder known to cause HG (eg, congenital anorchia, Klinefelter's syndrome). Delayed puberty (DP): Dx of DP in males at birth. Breast cancer (BC): Dx for the palliative treatment of inoperable BC in females at birth. Gender Dysphoria (GD) (off-label): Dx of GD. Patient is a female-to-male transsexual. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | HG(init): (New to T tx:6 mo. Cont T tx:12 mo), (reauth): 12 mo. BC, GD: 12 mo. DP: 6 mo. |
| Other Criteria | HG (Reauth): 1) Follow-up total serum T level within or below the normal limits of the reporting lab, or 2) Follow-up total serum T level outside of upper limits of normal for the reporting lab and the dose is adjusted, OR 3) Has a condition that may cause altered SHBG (eg, thyroid disorder, HIV disease, liver disorder, diabetes, obesity), and one of the following: Follow-up calculated free or bioavailable T level within or below the normal limits of the reporting lab, or follow-up calculated free or bioavailable T level outside of upper limits of normal for the reporting lab and the dose is adjusted. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

APOKYN (S)

Products Affected

- Apokyn INJ 30MG/3ML

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | PD (Initial): Patient is using Apokyn with any 5-HT3 antagonist (eg, ondansetron, granisetron, dolasetron, palonosetron, alosetron) |
| Required Medical Information | Parkinson's disease (PD) (Initial): Diagnosis of advanced PD. Patient is experiencing acute intermittent hypomobility (defined as “off” episodes characterized by muscle stiffness, slow movements, or difficulty starting movements). Patient is receiving Apokyn in combination with other medications for the treatment of PD (e.g., carbidopa/levodopa, pramipexole, ropinirole, etc.). |
| Age Restrictions | N/A |
| Prescriber Restrictions | PD (Initial): Prescribed by or in consultation with a neurologist. |
| Coverage Duration | PD (Initial, reauth): 12 months |
| Other Criteria | PD (Reauth): Documentation of positive clinical response to Apokyn therapy. |

ARANESP (S)

Products Affected

- Aranesp Albumin Free INJ
 100MCG/0.5ML, 100MCG/ML,
 10MCG/0.4ML, 150MCG/0.3ML,
 200MCG/0.4ML, 200MCG/ML,
 25MCG/0.42ML, 25MCG/ML,
 300MCG/0.6ML, 300MCG/ML,
 40MCG/0.4ML, 40MCG/ML,
 500MCG/ML, 60MCG/0.3ML,
 60MCG/ML

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Anemia with Chronic Kidney Disease (CKD) (Initial): Diagnosis (Dx) of CKD. Anemia by lab values (Hct less than 30% or Hgb less than 10 g/dL) collected within 30 days of request. One of the following: a) both of the following: Patient is on dialysis, patient is without ESRD OR b) all of the following: patient is not on dialysis, the rate of hemoglobin decline indicates the likelihood of requiring a red blood cell (RBC) transfusion, and reducing the risk of alloimmunization and/or other RBC transfusion-related risks is a goal. Anemia with chemo (Initial): Other causes of anemia have been ruled out. Anemia by lab values (Hct less than 30%, Hgb less than 10 g/dL) collected within the prior 2 weeks of request. Cancer is a non-myeloid malignancy. Patient is concurrently on chemo, or will receive concomitant chemo for a minimum of 2 months, or anemia is caused in part by cancer chemo. CKD (init, reauth), Chemo (init), MDS (init): Verify iron evaluation for adequate iron stores. Anemia in Myelodysplastic Syndrome (MDS) (Initial): Dx of MDS. Serum erythropoietin level is 500 mU/mL or less, or dx of transfusion-dependent MDS. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|--------------------------|--|
| Coverage Duration | CKD (Init): 6 mo. CKD (reauth): 12 mo. Chemo(init, reauth): 3 mo. MDS: (init) 3 mo,(reauth) 12 mo. |
| Other Criteria | Subject to ESRD review. CKD (Reauth): Dx of CKD. One of the following: 1) Most recent or average (avg) Hct over 3 mo is 33% or less (Hgb 11 g/dL or less) for patients on dialysis, without ESRD, 2) Most recent or avg Hct over 3 mo is 30% or less (Hgb 10 g/dL or less) for patients not on dialysis, OR 3) Most recent or avg Hct over 3 mo is 36% or less (Hgb 12 g/dL or less) for pediatric patients. Decrease in the need for blood transfusion or Hgb increased by 1 g/dL or more from pre-treatment level. Chemo (Reauth): Anemia by lab values (Hgb less than 10 g/dl or Hct less than 30%) collected within the prior 2 weeks of request. Decrease in the need for blood transfusion or Hgb increased by 1 g/dL or more from pre-treatment level. Patient is concurrently on chemo, or will receive concomitant chemo for a minimum of 2 months, or anemia is caused in part by cancer chemo. MDS (Reauth): Most recent or avg Hct over 3 months is 36% or less, OR most recent or avg Hgb over 3 months is 12 g/dl or less. Decrease in the need for blood transfusion or Hgb increased by 1 g/dL or more from pre-treatment level. Off-label uses (except MDS, HCV): Will not be approved if patient has Hgb greater than 10 g/dL or Hct greater than 30%. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

ARCALYST (S)

Products Affected

- Arcalyst

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Cryopyrin-Associated Periodic Syndromes (CAPS) (Initial): Diagnosis of CAPS, including Familial Cold Auto-inflammatory Syndrome (FCAS) and/or Muckle-Wells Syndrome (MWS). The medication will not be used in combination with another biologic. |
| Age Restrictions | CAPS (Initial): 12 years of age or older |
| Prescriber Restrictions | CAPS (initial): Prescribed by or in consultation with an immunologist, allergist, dermatologist, rheumatologist, neurologist or other medical specialist. |
| Coverage Duration | CAPS (initial, reauth): 12 months |
| Other Criteria | CAPS (Reauth): Patient has experienced disease stability or improvement in clinical symptoms while on therapy as evidence by one of the following: A) improvement in rash, fever, joint pain, headache, conjunctivitis, B) decreased number of disease flare days, C) normalization of inflammatory markers (CRP, ESR, SAA), D) corticosteroid dose reduction, OR E) improvement in MD global score or active joint count. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ARZERRA (S)

Products Affected

- Arzerra

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Refractory chronic lymphocytic leukemia (CLL): Diagnosis of CLL. Disease is refractory to both fludarabine and alemtuzumab. Previously untreated CLL: Diagnosis of CLL. Patient is previously untreated for CLL. Patient is not an appropriate candidate for fludarabine-based therapy. Used in combination with chlorambucil. Recurrent or progressive Chronic Lymphocytic Leukemia (CLL): Diagnosis of chronic lymphocytic leukemia (CLL). Disease is recurrent or progressive. Arzerra is used for an extended treatment for patients who are in complete or partial response after at least two lines of therapy. Relapsed Chronic Lymphocytic Leukemia (CLL): Diagnosis of CLL. Disease has relapsed. Used in combination with fludarabine and cyclophosphamide. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist/hematologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

AUBAGIO (S)

Products Affected

- Aubagio

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Multiple Sclerosis (MS): Diagnosis of a relapsing form of MS (eg, relapsing-remitting MS, secondary-progressive MS with relapses). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

AURYXIA (S)

Products Affected

- Auryxia

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | Exclude if used for iron deficiency anemia in chronic kidney disease (CKD) not on dialysis. |
| Required Medical Information | Hyperphosphatemia in chronic kidney disease: Diagnosis of hyperphosphatemia. Patient has chronic kidney disease (CKD). Patient is on dialysis. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

AUSTEDO (S)

Products Affected

- Austedo

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Chorea associated with Huntington's disease (initial): Diagnosis of Chorea associated with Huntington's disease. Tardive dyskinesia (initial): Diagnosis of moderate to severe tardive dyskinesia. One of the following: 1) Patient has persistent symptoms of tardive dyskinesia despite a trial of dose reduction, tapering, or discontinuation of the offending medication or 2) Patient is not a candidate for a trial of dose reduction, tapering, or discontinuation of the offending medication. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Huntington's disease chorea (initial): Prescribed by a neurologist. Tardive dyskinesia (initial): Prescribed by or in consultation with a neurologist or psychiatrist. |
| Coverage Duration | Initial: 3 months. Reauth: 12 months |
| Other Criteria | All indications (Reauth): Documentation of positive clinical response to Austedo therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

AYVAKIT (S)

Products Affected

- Ayvakit

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Gastrointestinal stromal tumor (GIST): Diagnosis of GIST. Disease is one of the following: unresectable or metastatic. Presence of platelet-derived growth factor receptor alpha (PDGFRA) exon 18 mutation, including PDGFRA D842V mutations. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

BALVERSA (S)

Products Affected

- Balversa

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Urothelial Carcinoma: Diagnosis of urothelial carcinoma (UC). One of the following: Locally advanced or Metastatic AND Patient has fibroblast growth factor receptor (FGFR) 3 or FGFR2 genetic alterations as detected by an U.S. Food and Drug Administration (FDA)-approved test (therascreen FGFR RGQ RT-PCR Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). One of the following: 1) Patient has progressed during or following at least one line of prior chemotherapy or immunotherapy OR 2) Patient has progressed within 12 months of neoadjuvant or adjuvant platinum-containing chemotherapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

BAVENCIO (S)

Products Affected

- Bavencio

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Merkel Cell Carcinoma (MCC): Diagnosis of metastatic Merkel cell carcinoma. Urothelial Carcinoma (UC): Diagnosis of locally advanced or metastatic urothelial carcinoma. One of the following: 1) Patient has disease progression during or following platinum-containing chemotherapy, OR 2) Patient has disease progression within 12 months of neoadjuvant or adjuvant treatment with platinum-containing chemotherapy |
| Age Restrictions | MCC: Patient is 12 years of age or older. |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

BELEODAQ (S)

Products Affected

- Beleodaq

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Peripheral T-cell lymphoma (PTCL): Diagnosis of PTCL. Trial and failure, contraindication, or intolerance to at least one prior therapy (e.g., conventional chemotherapy). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist/oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

BENLYSTA (S)

Products Affected

- Benlysta

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Systemic lupus erythematosus (SLE) (init): Diagnosis of active SLE. Autoantibody positive (ie, anti-nuclear antibody [ANA] titer greater than or equal to 1:80 or anti-dsDNA level greater than or equal to 30 IU/mL). Currently receiving at least one standard of care treatment for active SLE (eg, antimalarials [eg, Plaquenil (hydroxychloroquine)], corticosteroids [eg, prednisone], or immunosuppressants [eg, methotrexate, Imuran (azathioprine)]). |
| Age Restrictions | N/A |
| Prescriber Restrictions | SLE (init): Prescribed by or in consultation with a rheumatologist |
| Coverage Duration | SLE (init, reauth): 6 months |
| Other Criteria | SLE (reauth): Documentation of positive clinical response to Benlysta therapy. |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

BENZODIAZEPINES (S)

Products Affected

- Alprazolam TABS
- Alprazolam Er
- Alprazolam Intensol
- Alprazolam Odt
- Alprazolam Xr
- Chlordiazepoxide Hcl CAPS 10MG, 5MG
- Chlordiazepoxide Hydrochloride
- Estazolam
- Lorazepam CONC
- Lorazepam INJ 2MG/ML, 4MG/ML
- Lorazepam TABS
- Lorazepam Intensol
- Oxazepam
- Temazepam

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Verify the medication is being used for an FDA-approved diagnosis. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

BERINERT (S)

Products Affected

- Berinert

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Treatment of hereditary angioedema (HAE) attacks: Diagnosis of HAE. For the treatment of acute HAE attacks. Not used in combination with other approved treatments for acute HAE attacks. |
| Age Restrictions | N/A |
| Prescriber Restrictions | HAE: Prescribed by or in consultation with an immunologist, allergist, or rheumatologist |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

BESPONSA (S)

Products Affected

- Besponsa

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | B-cell precursor acute lymphoblastic leukemia (ALL): Diagnosis of B-cell precursor acute lymphoblastic leukemia (ALL). Disease is relapsed or refractory. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist/oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

BLINCYTO (S)

Products Affected

- Blincyto

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Relapsed or refractory B-cell precursor acute lymphoblastic leukemia (ALL): Diagnosis of relapsed or refractory B-cell precursor acute lymphoblastic leukemia/acute lymphoblastic lymphoma. Minimal residual disease (MRD)-positive B-cell precursor ALL (MRD+ ALL): Diagnosis of B-cell precursor ALL. Patient is in their first or second complete remission. Documentation of MRD greater than or equal to 0.1%. |
| Age Restrictions | N/A |
| Prescriber Restrictions | All Indications: Prescribed by or in consultation with a hematologist/oncologist |
| Coverage Duration | ALL: 12 months. MRD+ ALL: 6 months. |
| Other Criteria | Subject to Part B vs. Part D review. All Indications: Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

BORTEZOMIB (S)

Products Affected

- Bortezomib

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Multiple myeloma (MM): Diagnosis of MM. Mantle cell lymphoma (MCL): Diagnosis of MCL AND patient has received at least one prior therapy for MCL. |
| Age Restrictions | N/A |
| Prescriber Restrictions | MM, MCL: Prescribed by or in consultation with an oncologist/hematologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

BOSULIF (S)

Products Affected

- Bosulif

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Chronic myelogenous/myeloid leukemia (CML): Diagnosis of Philadelphia chromosome-positive (Ph+) CML. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

BOTOX (S)

Products Affected

- Botox

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | <p>Neuromuscular Disorders (init): Strabismus, blepharospasm associated with dystonia (eg, benign essential blepharospasm), treatment of upper or lower limb spasticity, VII cranial nerve disorders (hemifacial spasms), cervical dystonia</p> <p>Hyperhidrosis(HH): (Init) Dx of primary axillary HH. Score of 3 or 4 on the Hyperhidrosis Disease Severity Scale (HDSS) or skin maceration with secondary infection.</p> <p>Migraine:(Init) Dx of chronic migraines (greater than or equal to 15 migraine headache days per month with headache lasting 4 hours a day or longer). TF/C/I to prophylactic therapy with at least two of the following agents, each given for a trial of at least two months: antidepressants [ie, Effexor (venlafaxine)], antiepileptics [ie, Depakote/Depakote ER (divalproex sodium), Topamax (topiramate)], beta-blockers [eg, atenolol, Inderal (propranolol), nadolol, timolol, Toprol XL (metoprolol)]</p> <p>Achalasia:(Init) High risk of complication from or failure to pneumatic dilation or myotomy, or prior dilation caused esophageal perforation, or patient has an epiphrenic diverticulum or hiatal hernia.</p> <p>Anal Fissure (AF)(Init): Dx of chronic AF. At least 2 months of either nocturnal pain and bleeding or postdefecation pain.</p> <p>Chronic Back Pain (CBP):(Init) Dx of low back pain lasting greater than or equal to six months.</p> <p>Urinary incont (UI):(init) Neurogenic detrusor overactivity associated with a neurologic condition (eg, spinal cord injury [SCI], multiple sclerosis) or detrusor sphincter dyssynergia with SCI.</p> <p>Overactive bladder (OAB): (init) Dx of OAB. One of the following symptoms: urge urinary incontinence, urgency, frequency.</p> |
| Age Restrictions | N/A |
| Prescriber Restrictions | <p>Migraine (initial): Prescribed by a neurologist or pain specialist.</p> <p>CBP (initial): Prescribed by a neurologist, neurosurgeon, orthopedist, or pain specialist.</p> <p>UI, OAB (initial): Prescribed by a neurologist, neurosurgeon, or urologist.</p> |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|--------------------------|--|
| Coverage Duration | Achalasia: 6moCBP:1 tx(series of injxs)UI:3mo(1 dose,200units)Other:3mo |
| Other Criteria | UI, OAB, CBP, Neuromuscular Disorders:(Reauth) Confirmed improvement in symptoms with initial Botox treatment. At least 3 months have or will have elapsed since the last treatment with Botox HH:(Reauth) At least a 2-point improvement in HDSS. Migraine:(Reauth) Reduction in headache frequency or intensity. Confirmation of decreased utilization of pain medications (eg, narcotic analgesics, NSAIDs) or triptans, or a reduction in the number of ER visits. Achalasia:(Reauth) Documentation of improvement or reduction in symptoms of achalasia (ie, dysphagia, regurgitation, chest pain). At least 6 months have or will have elapsed since last series of injections AF: (Reauth) Incomplete healing of fissure or recurrence of fissure. Improved symptoms with prior treatment with Botox. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

BRAFTOVI (S)

Products Affected

- Braftovi

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Melanoma: Diagnosis of unresectable melanoma or metastatic melanoma. Cancer is BRAF V600E or V600K mutant type (MT) as detected by a U.S. Food and Drug Administration (FDA)-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Used in combination with Mektovi (binimetinib). Colorectal Cancer: One of the following diagnoses: Colon Cancer or Rectal Cancer. One of the following: 1) Unresectable or advanced disease or 2) Metastatic disease. Patient has received prior therapy. Cancer is BRAF V600E mutant type as detected by a U.S. Food and Drug Administration (FDA)-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Used in combination with Erbitux (cetuximab). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

BRIVIACT (S)

Products Affected

- Briviact

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Partial-onset seizures: Diagnosis of partial-onset seizures. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

BRUKINSA (S)

Products Affected

- Brukinsa

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Diagnosis of relapsed or refractory mantle cell lymphoma (MCL). Trial and failure, contraindication, or intolerance to at least ONE combination treatment of rituximab and chemotherapy (e.g., BR, R-CHOP, R-CVP, R-FCM). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist/hematologist. |
| Coverage Duration | 12 months. |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

BYNFEZIA (S)

Products Affected

- Bynfezia Pen

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Acromegaly (initial): Diagnosis of acromegaly. One of the following: A) Inadequate response to surgical resection and/or pituitary irradiation OR B) Patient is not a candidate for surgical resection or pituitary irradiation. Trial and failure, contraindication or intolerance to a dopamine agonist (e.g., bromocriptine or cabergoline) at maximally tolerated doses. Carcinoid tumor (initial): Diagnosis of metastatic carcinoid tumor requiring symptomatic treatment of severe diarrhea or flushing episodes. Vasoactive intestinal peptide tumor (initial): Diagnosis of vasoactive intestinal peptide tumor requiring treatment of profuse watery diarrhea. All indications (initial): Trial and failure, or intolerance to generic octreotide. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | All uses (initial, reauth): 12 months |
| Other Criteria | Acromegaly (reauth): Documentation of positive clinical response to therapy (e.g., reduction or normalization of IGF-1/GH level for same age and sex, reduction in tumor size). Carcinoid tumor (reauth): Patient has improvement in number of diarrhea or flushing episodes. Vasoactive intestinal peptide tumor (reauth): Patient has improvement in number of diarrhea episodes. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

CABLIVI (S)

Products Affected

- Cablivi

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Acquired thrombocytic thrombocytopenic purpura (aTTP): Diagnosis of aTTP. First dose was/will be administered by a healthcare provider as a bolus intravenous injection. Used in combination with immunosuppressive therapy (e.g. rituximab, glucocorticoids). One of the following: 1) Used in combination with plasma exchange or 2) both of the following: patient has completed plasma exchange and less than 59 days have or will have elapsed beyond the last plasma exchange. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist or oncologist. |
| Coverage Duration | 3 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

CABOMETYX (S)

Products Affected

- Cabometyx

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Renal cell carcinoma (RCC): Diagnosis of RCC. RCC is advanced. Hepatocellular Carcinoma (HCC): Diagnosis of HCC. One of the following: a) Trial and failure or intolerance to Nexavar (sorafenib tosylate), or b) Patient has metastatic disease, or c) Patient has extensive liver tumor burden, or d) Patient is inoperable by performance status or comorbidity (local disease or local disease with minimal extrahepatic disease only), or e) Disease is unresectable. |
| Age Restrictions | N/A |
| Prescriber Restrictions | RCC: Prescribed by or in consultation with one of the following: an oncologist or nephrologist. HCC: Prescribed by or in consultation with one of the following: oncologist, hepatologist, or gastroenterologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

CALQUENCE (S)

Products Affected

- Calquence

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Mantle Cell Lymphoma: Diagnosis of mantle cell lymphoma (MCL) AND patient has received at least one prior therapy for MCL. Chronic Lymphocytic Leukemia (CLL) or Small Lymphocytic Lymphoma (SLL): Diagnosis of CLL or SLL. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist/oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

CAPRELSA (S)

Products Affected

- Caprelsa

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Thyroid Cancer: Diagnosis of medullary thyroid cancer (MTC). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with oncologist or endocrinologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

CARISOPRODOL (S)

Products Affected

- Carisoprodol TABS
- Vanadom

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | The drug is being prescribed for an FDA-approved indication. If the patient is 65 years of age or older, the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

CAYSTON (S)

Products Affected

- Cayston

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Cystic fibrosis (CF) (Initial, Reauth): Diagnosis of CF AND Patient has evidence of Pseudomonas aeruginosa in the lungs. |
| Age Restrictions | CF (Initial): 7 years of age or older |
| Prescriber Restrictions | N/A |
| Coverage Duration | CF (Initial, reauth): 12 months |
| Other Criteria | CF (Reauth): Patient is benefiting from treatment (i.e. improvement in lung function [forced expiratory volume in one second (FEV1)], decreased number of pulmonary exacerbations). |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

CERDELGA (S)

Products Affected

- Cerdelga

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Gaucher disease (Initial): Diagnosis of Gaucher disease type 1. Patient is an extensive metabolizer (EM), intermediate metabolizer (IM), or poor metabolizer (PM) of cytochrome P450 enzyme (CYP) 2D6 as detected by an FDA-cleared test. |
| Age Restrictions | Gaucher disease (initial): 18 years of age or older |
| Prescriber Restrictions | N/A |
| Coverage Duration | Gaucher disease (initial, reauth): 12 months |
| Other Criteria | Gaucher disease (Reauth): Patient's condition has not progressed, as defined by ALL of the following: A) Hemoglobin level decreased greater than 1.5 g/dL from baseline, AND B) Platelet count decreased greater than 25% from baseline, AND C) Spleen volume increased greater than 25% from baseline, AND D) Liver volume increased greater than 20% from baseline. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

CEREZYME (S)

Products Affected

- Cerezyme

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Gaucher disease: Diagnosis of type 1 Gaucher disease. Patient has evidence of symptomatic disease (e.g., moderate to severe anemia, thrombocytopenia, bone disease, hepatomegaly, or splenomegaly). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | Gaucher disease: 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

CHOLBAM (S)

Products Affected

- Cholbam

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Bile acid synthesis disorders due to single enzyme defects (BAS) (initial): diagnosis of a bile acid synthesis disorder due to a single enzyme defect. Peroxisomal disorders (PD) (initial): All of the following: 1) diagnosis of peroxisomal disorder, 2) patient exhibits at least one of the following: a) liver disease (eg, jaundice, elevated serum transaminases), OR b) steatorrhea, OR c) complications from decreased fat-soluble vitamin absorption (eg, poor growth), AND 3) Cholbam will be used as an adjunctive treatment. |
| Age Restrictions | N/A |
| Prescriber Restrictions | All uses (initial): Prescribed by a hepatologist, medical geneticist, pediatric gastroenterologist, OR other specialist that treats inborn errors of metabolism. |
| Coverage Duration | All uses (initial, reauth): 12 months |
| Other Criteria | All uses (reauth): documentation of positive clinical response to Cholbam therapy. |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

CHORIONIC GONADOTROPIN (S)

Products Affected

- Chorionic Gonadotropin INJ
- Novarel
- Pregnyl W/diluent Benzyl Alcohol/nacl

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Prepubertal Cryptorchidism: Diagnosis of prepubertal cryptorchidism not due to anatomical obstruction. Male Hypogonadotropic Hypogonadism (MHH) (initial): Diagnosis of male hypogonadism secondary to pituitary deficiency, and low testosterone (below normal reference value provided by the physician's laboratory) and one of the following: a) low LH (below normal reference value provided by the physician's laboratory) or b) low FSH (below normal reference value provided by the physician's laboratory). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | Prepubertal Cryptorchidism: 6 wks. MHH (initial, reauth): 12 months. |
| Other Criteria | Excluded if used to promote fertility. MHH (Reauth): Documentation of positive clinical response to therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

CIALIS (S)

Products Affected

- Tadalafil TABS 2.5MG, 5MG

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | Concurrent use of nitrates. |
| Required Medical Information | Diagnosis of benign prostatic hyperplasia (BPH). Trial and failure, contraindication, or intolerance to an alpha-blocker (e.g., doxazosin, prazosin, tamsulosin) or a 5-alpha reductase inhibitor (e.g., dutasteride, finasteride). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

CICLOPIROX (S)

Products Affected

- Ciclodan SOLN
- Ciclopirox Nail Lacquer

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | All of the following: 1) Patient does not have lunula (matrix) involvement, 2) one of the following: a) Diagnosis of onychomycosis of the toenails, OR b) Diagnosis of onychomycosis of the fingernails, 3) Diagnosis of onychomycosis has been confirmed by one of the following: a) positive potassium hydroxide (KOH) preparation, OR b) culture, OR c) histology, 4) If toenail onychomycosis, patient has mild to moderate disease involving at least 1 great toenail, AND 5) Trial and failure, contraindication, or intolerance to oral terbinafine. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 48 weeks. |
| Other Criteria | N/A |

CIMZIA (S)

Products Affected

- Cimzia

- Cimzia Starter Kit

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Rheumatoid Arthritis (RA, initial): Diagnosis (dx) of moderately to severely active RA. TF/C/I to both Enbrel and Humira, or attestation demonstrating a trial may be inappropriate, OR for continuation of prior Cimzia therapy. Crohn's Disease (CD, initial): Dx of moderately to severely active CD. Trial and failure, contraindication, or intolerance (TF/C/I) to one of the following conventional therapies: 6-mercaptopurine (Purinethol), Azathioprine (Imuran), Corticosteroid (eg, prednisone, methylprednisolone), Methotrexate (Rheumatrex, Trexall). TF/C/I to Humira OR for continuation of prior Cimzia therapy. Psoriatic Arthritis (PsA, initial): Dx of active PsA. TF/C/I to both Enbrel and Humira, or attestation demonstrating a trial may be inappropriate, OR for continuation of prior Cimzia therapy. Ankylosing Spondylitis (AS, initial): Dx of active AS. TF/C/I to both Enbrel and Humira, or attestation demonstrating a trial may be inappropriate, OR for continuation of prior Cimzia therapy. Plaque Psoriasis (initial): Dx of moderate to severe plaque psoriasis. TF/C/I to Cosentyx AND either Humira or Enbrel OR for continuation of prior Cimzia therapy. Non-radiographic axial spondyloarthritis (nr-axSpA, initial): Dx of nr-axSpA with objective signs of inflammation (eg, C-reactive protein [CRP] levels above the upper limit of normal and/or sacroiliitis on magnetic resonance imaging [MRI], indicative of inflammatory disease, but without definitive radiographic evidence of structural damage on sacroiliac joints.) TF/C/I to two non-steroidal anti-inflammatory drugs (NSAIDs). |
| Age Restrictions | N/A |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|--------------------------------|--|
| Prescriber Restrictions | CD (init): Prescribed by or in consultation with a gastroenterologist. RA, AS, nr-axSpA (init): Prescribed by or in consultation with a rheumatologist. PsA (init): Prescribed by or in consultation with a dermatologist or rheumatologist. Plaque psoriasis (init): Prescribed by or in consultation with a dermatologist. |
| Coverage Duration | RA, PsA, AS, Plaque psoriasis, nr-axSpA (init, reauth): 12 mos. CD (init): 16 wks. (reauth): 12 mos. |
| Other Criteria | Reauth (all indications): Documentation of positive clinical response to Cimzia therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

CINRYZE (S)

Products Affected

- Cinryze

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Prophylaxis of hereditary angioedema (HAE) attacks: Diagnosis of HAE. For prophylaxis against HAE attacks. |
| Age Restrictions | N/A |
| Prescriber Restrictions | HAE (prophylaxis): Prescribed by or in consultation with an immunologist, allergist, or rheumatologist |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

COMETRIQ (S)

Products Affected

- Cometriq

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Medullary thyroid cancer (MTC): Diagnosis of Metastatic MTC. |
| Age Restrictions | N/A |
| Prescriber Restrictions | MTC: Prescribed by or in consultation with an oncologist/hematologist or endocrinologist. |
| Coverage Duration | All uses: 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

COPIKTRA (S)

Products Affected

- Copiktra

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Chronic Lymphocytic Leukemia (CLL)/Small Lymphocytic Lymphoma (SLL): Diagnosis of CLL or SLL. Disease is relapsed or refractory. Trial and failure, contraindication, or intolerance to at least two prior therapies for CLL/SLL (e.g., Leukeran [chlorambucil], Gazyva [obinutuzumab], Arzerra [ofatumumab], Bendeka [bendamustine], Imbruvica [ibrutinib], Rituxan [rituximab], etc.). Follicular Lymphoma: Diagnosis of follicular lymphoma. Disease is relapsed or refractory. Trial and failure, contraindication, or intolerance to at least two prior systemic therapies for follicular lymphoma (e.g., Leukeran [chlorambucil], Gazyva [obinutuzumab], Arzerra [ofatumumab], Bendeka [bendamustine], Imbruvica [ibrutinib], Rituxan [rituximab], etc.). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist/oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

CORLANOR (S)

Products Affected

- Corlanor

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Chronic heart failure (CHF) (initial): Diagnosis of CHF. Patient has NYHA Class II, III, or IV symptoms. Patient has a left ventricular ejection fraction less than or equal to 35%. Patient is in sinus rhythm. Patient has a resting heart rate of greater than or equal to 70 beats per minute. One of the following: patient is on a beta-blocker at a maximally tolerated dose, or patient has a contraindication or intolerance to beta-blocker therapy. Patient has been hospitalized for worsening HF in the previous 12 months. Trial and failure, contraindication, or intolerance to maximally tolerated doses of an ACE inhibitor or ARB. Dilated Cardiomyopathy (DCM) (initial): Diagnosis of heart failure due to DCM. Patient has NYHA Class II, III, or IV symptoms. Patient is in sinus rhythm. Patient has an elevated heart rate. Trial and failure, contraindication or intolerance to one of the following: 1) Beta blocker (e.g., bisoprolol, metoprolol succinate extended release), 2) Angiotensin-converting enzyme (ACE) inhibitor (e.g., captopril, enalapril), or 3) Diuretic Agent (e.g., spironolactone, furosemide). |
| Age Restrictions | N/A |
| Prescriber Restrictions | CHF, DCM (initial): Prescribed by or in consultation with a cardiologist |
| Coverage Duration | CHF, DCM (initial, reauth): 12 months |
| Other Criteria | CHF, DCM (reauth): Documentation of positive clinical response to therapy. |

COSENTYX (S)

Products Affected

- Cosentyx
- Cosentyx Sensoready Pen

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Plaque psoriasis (Initial): Diagnosis of moderate to severe plaque psoriasis. One of the following: Trial and failure, contraindication, or intolerance to Enbrel (etanercept) OR Humira (adalimumab), OR for continuation of prior Cosentyx therapy. Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. One of the following: Trial and failure, contraindication, or intolerance to Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR for continuation of prior Cosentyx therapy. Ankylosing Spondylitis (AS) (Initial): Diagnosis of active AS. One of the following: Trial and failure, contraindication, or intolerance to Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR for continuation of prior Cosentyx therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Plaque psoriasis (initial): Prescribed by or in consultation with a dermatologist. PsA (initial): Prescribed by or in consultation with a rheumatologist or dermatologist. AS (initial): Prescribed by or in consultation with a rheumatologist. |
| Coverage Duration | All uses (initial, reauth): 12 months |
| Other Criteria | All indications (Reauth): Documentation of positive clinical response to Cosentyx therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

COTELLIC (S)

Products Affected

- Cotellic

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Melanoma: Diagnosis of unresectable or metastatic melanoma. Patient has a BRAF V600E or V600K mutation as detected by a U.S. Food and Drug Administration (FDA)-approved test (e.g., cobas 4800 BRAF V600 Mutation Test) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Used in combination with vemurafenib. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

CRINONE (S)

Products Affected

- Crinone

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | All indications: Excluded if for fertility uses. |
| Required Medical Information | Secondary amenorrhea: Diagnosis of secondary amenorrhea (the absence of menses in women who have already started menstruation who are not pregnant, breastfeeding, or in menopause). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

CYRAMZA (S)

Products Affected

- Cyramza

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Gastric cancer: All of the following: 1) diagnosis of one of the following: a) gastric adenocarcinoma, OR b) gastro-esophageal junction (GEJ) adenocarcinoma, AND 2) disease is one of the following: a) locally advanced, OR b) metastatic, AND 3) disease has progressed on or after one of the following first-line therapies: a) fluoropyrimidine-containing chemotherapy (eg, fluorouracil, capecitabine), OR b) platinum-containing chemotherapy (eg, cisplatin, carboplatin, oxaliplatin). Non-small cell lung cancer: All of the following: 1) diagnosis of metastatic non-small cell lung cancer, AND 2) used in combination with docetaxel, AND 3) disease has progressed on or after platinum-based chemotherapy (eg, cisplatin, carboplatin, oxaliplatin). Metastatic colorectal cancer (mCRC): 1) Diagnosis of metastatic CRC AND 2) Used in combination with irinotecan or FOLFIRI (irinotecan, folinic acid, and 5-fluorouracil), AND 3) Patient had disease progression on or after prior therapy with bevacizumab, oxaliplatin, and a fluoropyrimidine. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

CYSTARAN (S)

Products Affected

- Cystaran

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Cystinosis: Diagnosis of cystinosis, confirmed by elevated leukocyte cystine levels (LCL), genetic analysis of the CTNS gene or corneal cystine crystal accumulation AND Patient is concomitantly receiving treatment with oral cysteamine. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

DACOGEN (S)

Products Affected

- Decitabine

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Myelodysplastic syndrome (MDS): Diagnosis of myelodysplastic syndrome. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist/oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

DALIRESP (S)

Products Affected

- Daliresp

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Chronic Obstructive Pulmonary Disease (COPD): (Initial) Diagnosis of COPD. History of COPD exacerbations which required the use of systemic corticosteroids, antibiotics, or hospital admission. Trial and failure, intolerance, or contraindication to two prior therapies for COPD. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | COPD (init, reauth): 12 months |
| Other Criteria | COPD (reauth): Documentation of positive clinical response to Daliresp therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

DARAPRIM (S)

Products Affected

- Daraprim

- Pyrimethamine TABS

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Toxoplasmosis: 1) Patient is using pyrimethamine for the treatment of toxoplasmic encephalitis, secondary prophylaxis of toxoplasmic encephalitis, or treatment of congenital toxoplasmosis OR 2) Patient is using pyrimethamine for the primary prophylaxis of toxoplasmic encephalitis, patient has experienced intolerance to prior prophylaxis with trimethoprim-sulfamethoxazole (TMP-SMX), and one of the following: patient has been re-challenged with TMP-SMX using a desensitization protocol and is still unable to tolerate, or evidence of life-threatening reaction to TMP-SMX in the past (eg, toxic epidermal necrolysis, Stevens-Johnson syndrome). Malaria: Patient is using pyrimethamine for the treatment of acute malaria or chemoprophylaxis of malaria. Patient does not have megaloblastic anemia due to folate deficiency. The provider acknowledges that pyrimethamine is not recommended by the Centers for Disease Control and Prevention (CDC) for the treatment and/or prophylaxis of malaria. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an infectious disease specialist |
| Coverage Duration | 12 months |
| Other Criteria | Toxoplasmosis only: Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

DARZALEX (S)

Products Affected

- Darzalex

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Relapsed/Refractory Multiple Myeloma (MM): Diagnosis of MM. One of the following: A) Both of the following: Used as monotherapy and Patient has received at least three prior treatment regimens which included both a proteasome inhibitor (eg, bortezomib [Velcade], carfilzomib [Kyprolis]) and an immunomodulatory agent (eg, lenalidomide [Revlimid], thalidomide [Thalomid]) or patient is double-refractory to a proteasome inhibitor and an immunomodulatory agent. OR B) Patient has received at least one prior therapy. Darzalex will be used in combination with either 1) lenalidomide and dexamethasone or 2) bortezomib and dexamethasone. OR C) Both of the following: used in combination with both pomalidomide and dexamethasone, AND patient has received at least two prior therapies including lenalidomide and a proteasome inhibitor (e.g., bortezomib [Velcade], carfilzomib [Kyprolis]). Newly Diagnosed Multiple Myeloma: Newly diagnosed multiple myeloma, patient is ineligible for autologous stem cell transplant and used in combination with all of the following: bortezomib, melphalan, and prednisone. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist/hematologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

DAURISMO (S)

Products Affected

- Daurismo

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Acute myeloid leukemia (AML): Diagnosis of newly-diagnosed acute myeloid leukemia (AML) AND Used in combination with low-dose cytarabine AND One of the following: 1) Patient is greater than or equal to 75 years old, or 2) Patient has comorbidities that preclude the use of intensive induction chemotherapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist/oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

DEFERASIROX (S)

Products Affected

- Deferasirox
- Exjade
- Jadenu
- Jadenu Sprinkle

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Chronic Iron Overload Due to Blood Transfusions (Initial): Diagnosis of chronic iron overload due to blood transfusions (transfusional hemosiderosis). Patient has a baseline ferritin level more than 1,000 mcg/L. Patient has required the transfusion of at least 100 mL/kg packed red blood cells. Myelodysplastic Syndrome (MDS) (Initial): Diagnosis of MDS. Patient has Low or Intermediate-1 disease or is a potential transplant patient. Patient has received more than 20 red blood cell transfusions. Chronic iron overload due to non-transfusion-dependent thalassemia (NTDT) (Initial): Diagnosis of chronic iron overload due to NTDT. Liver iron concentration (LIC) 5 milligrams of iron per gram of liver dry weight (mg Fe/g dw) or higher. Serum ferritin level greater than 300 mcg/L. |
| Age Restrictions | Iron Overload Due to Blood Transfusions (initial): 2 years of age or older. NTDT (initial): 10 years of age or older |
| Prescriber Restrictions | N/A |
| Coverage Duration | Iron Overload Due to Blood Transfusions, MDS (initial, reauth):12 mo. NTDT (initial, reauth): 6mo. |
| Other Criteria | Iron Overload Due to Blood Transfusions, MDS (Reauth): Patient experienced a reduction from baseline in serum ferritin level or LIC. NTDT (Reauth): Patient has LIC 3 mg Fe/g dw or higher. Patient experienced a reduction from baseline in serum ferritin level or LIC. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

DEXMETHYLPHENIDATE (S)

Products Affected

- Dexmethylphenidate Hcl TABS 10MG, 5MG
- Dexmethylphenidate Hcl Er
- Dexmethylphenidate Hydrochloride TABS 2.5MG

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | One of the following: a) diagnosis of attention deficit hyperactivity disorder (ADHD), OR b) diagnosis of attention deficit disorder (ADD). |
| Age Restrictions | PA applies to members 19 years of age or older |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

DEXTROAMPHETAMINE (S)

Products Affected

- Dexedrine TABS
- Dextroamphetamine Sulfate SOLN
- Dextroamphetamine Sulfate TABS
- Dextroamphetamine Sulfate Er
- Zenedi

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | One of the following: a) diagnosis of attention deficit hyperactivity disorder (ADHD), OR b) diagnosis of attention deficit disorder (ADD), OR c) diagnosis of narcolepsy as confirmed by a sleep study (unless the prescriber provides justification confirming that a sleep study would not be feasible). |
| Age Restrictions | PA applies to members 19 years of age or older |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

DIACOMIT (S)

Products Affected

- Diacomit

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Diagnosis of seizures associated with Dravet syndrome (DS). Used in combination with clobazam. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a neurologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

DOXEPIN TOPICAL (S)

Products Affected

- Doxepin Hydrochloride CREA

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Diagnosis of moderate pruritus. Patient has atopic dermatitis or lichen simplex chronicus. Trial and failure, contraindication, or intolerance to at least one medium potency topical corticosteroid, or is not a candidate for topical corticosteroids (e.g., treatment is on face, axilla, or groin). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 14 days |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

DUOBRII (S)

Products Affected

- Duobrii

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Plaque Psoriasis: Diagnosis of plaque psoriasis. Both of the following: 1) Trial and failure, intolerance or contraindication to one high potency corticosteroid topical treatment (e.g., halobetasol propionate, clobetasol propionate, fluocinonide) AND 2) Trial and failure or intolerance to tazarotene. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Plaque Psoriasis: Prescribed by or in consultation with a dermatologist. |
| Coverage Duration | Plaque Psoriasis: 12 months |
| Other Criteria | N/A |

DUPIXENT (S)

Products Affected

- Dupixent

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Atopic dermatitis (initial): Diagnosis of moderate to severe atopic dermatitis. Trial and failure, contraindication, or intolerance to one medium to high potency topical corticosteroid. One of the following: A) Trial and failure or intolerance to pimecrolimus topical cream, unless the patient is not a candidate for pimecrolimus therapy (e.g., immunocompromised, severe atopic dermatitis), B) Trial and failure or intolerance to tacrolimus topical ointment, unless the patient is not a candidate for tacrolimus ointment therapy (e.g., immunocompromised), or C) Eucrisa (crisaborole). Eosinophilic Asthma (initial): Diagnosis of moderate to severe asthma. Asthma is an eosinophilic phenotype as defined by a baseline (pre-treatment) peripheral blood eosinophil level greater than or equal to 150 cells per microliter. One of the following: 1) Patient has had at least one or more asthma exacerbations requiring systemic corticosteroids within the past 12 months, 2) Any prior intubation for an asthma exacerbation, or 3) Prior asthma-related hospitalization within the past 12 months. Corticosteroid Dependent Asthma (initial): Diagnosis of moderate to severe asthma. Patient is currently dependent on oral corticosteroids for the treatment of asthma. Eosinophilic Asthma, Corticosteroid Dependent Asthma (initial): Patient is currently being treated with one of the following unless there is a contraindication or intolerance to these medications: a) Both of the following: i) High-dose inhaled corticosteroid (ICS) [e.g., greater than 500 mcg fluticasone propionate equivalent/day] and ii) additional asthma controller medication [e.g., leukotriene receptor antagonist, long-acting beta-2 agonist (LABA), theophylline], OR b) One maximally-dosed combination ICS/LABA product [e.g., Advair (fluticasone propionate/salmeterol), Dulera (mometasone/formoterol), Symbicort (budesonide/formoterol)]. |
| Age Restrictions | Asthma (initial): Age greater than or equal to 12 years. Atopic dermatitis: no age restriction. |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|---------------------------------------|---|
| <p>Prescriber Restrictions</p> | <p>Atopic dermatitis (Initial): Prescribed by or in consultation with one of the following: dermatologist, allergist/immunologist. Asthma (initial, reauth): Prescribed by or in consultation with a pulmonologist or allergist/immunologist. CRSwNP (initial, reauth): Prescribed by or in consultation with an otolaryngologist or allergist/immunologist.</p> |
| <p>Coverage Duration</p> | <p>Atopic Dermatitis, CRSwNP (Init/Reauth): 12 months. Asthma (Init): 6 mo. Asthma (reauth): 12 mo.</p> |
| <p>Other Criteria</p> | <p>Chronic rhinosinusitis with nasal polyposis (CRSwNP) (initial): Diagnosis of CRSwNP. Unless contraindicated, the patient has had an inadequate response to 2 months of treatment with an intranasal corticosteroid (eg, fluticasone, mometasone). Used in combination with another agent for CRSwNP (eg, intranasal corticosteroid). Atopic dermatitis (reauth): Documentation of a positive clinical response to therapy (e.g., reduction in body surface area involvement, reduction in pruritus severity). Eosinophilic Asthma (reauth): Documentation of a positive clinical response to therapy (e.g., reduction in exacerbations, improvement in forced expiratory volume in 1 second [FEV1], decreased use of rescue medications). Patient is currently being treated with ONE of the following unless there is a contraindication or intolerance to these medications: 1) Inhaled corticosteroid (ICS) and additional asthma controller medication [e.g., leukotriene receptor antagonist, long-acting beta-2 agonist (LABA), theophylline] OR 2) A combination ICS/LABA product (e.g., Advair [fluticasone propionate/salmeterol], Dulera [mometasone/formoterol], Symbicort [budesonide/formoterol]). Corticosteroid Dependent Asthma (reauth): Documentation of a positive clinical response to therapy (e.g., reduction in exacerbations, improvement in FEV1, reduction in oral corticosteroid dose). Patient is currently being treated with ONE of the following unless there is a contraindication or intolerance to these medications: 1) Inhaled corticosteroid (ICS) and additional asthma controller medication [e.g., leukotriene receptor antagonist, long-acting beta-2 agonist (LABA), theophylline] OR 2) A combination ICS/LABA product (e.g., Advair [fluticasone propionate/salmeterol], Dulera [mometasone/formoterol], Symbicort [budesonide/formoterol]). CRSwNP (reauth): Documentation of a positive clinical response to therapy (e.g., reduction in nasal polyps score [NPS, 0-8 scale], improvement in nasal congestion/obstruction score [NC, 0-3 scale]). Used in combination with another agent for CRSwNP (eg, intranasal corticosteroid).</p> |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

EGRIFTA (S)

Products Affected

- Egrifta

- Egrifta Sv

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | HIV-associated lipodystrophy (initial): All of the following: 1) diagnosis of HIV-associated lipodystrophy, 2) one of the following: a) waist-circumference of greater than or equal to 95 cm (37.4 inches) in men, OR b) waist-circumference of greater than or equal to 94 cm (37 inches) for women, 3) one of the following: a) Waist-to-hip ratio of greater than or equal to 0.94 for men, OR b) waist-to-hip ratio of greater than or equal to 0.88 for women, 4) body mass index (BMI) greater than 20 kg/m ² , AND 5) fasting blood glucose (FBG) levels less than or equal to 150 mg/dL (8.33 mmol/L), AND 6) patient has been on a stable regimen of antiretrovirals (eg, NRTIs, NNRTI, Protease Inhibitors, Integrase Inhibitors) for at least 8 weeks. |
| Age Restrictions | Initial: 18 years of age or older. |
| Prescriber Restrictions | N/A |
| Coverage Duration | Initial, reauth: 6 months |
| Other Criteria | (reauth): documentation of clinical improvement (eg, improvement in visceral adipose tissue [VAT], decrease in waist circumference, belly appearance, etc) while on Egrifta therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ELAPRASE (S)

Products Affected

- Elaprase

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Diagnosis of Hunter Syndrome (Mucopolysaccharidosis II (MPS II)) |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ELIGARD (S)

Products Affected

- Eligard

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Prostate Cancer: Diagnosis of advanced or metastatic prostate cancer. Trial and failure, contraindication, or intolerance to any brand Lupron formulation. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ELZONRIS (S)

Products Affected

- Elzonris

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Diagnosis of blastic plasmacytoid dendritic cell neoplasm (BPDCN). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist/oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

EMFLAZA (S)

Products Affected

- Emflaza

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Initial: Diagnosis of Duchenne muscular dystrophy (DMD). Patient has received genetic testing for a mutation of the dystrophin gene. One of the following: A) Documentation of a confirmed mutation of the dystrophin gene or B) Muscle biopsy confirmed an absence of dystrophin protein. Patient has had a trial and failure or intolerance to prednisone or prednisolone. Dose will not exceed 0.9 milligrams per kilogram of body weight once daily. |
| Age Restrictions | Initial: Patient is 2 years of age or older |
| Prescriber Restrictions | Initial: Prescribed by or in consultation with a neurologist who has experience treating children |
| Coverage Duration | Initial, Reauth: 12 months |
| Other Criteria | Reauth: Patient has experienced a benefit from therapy (e.g., improvement or preservation of muscle strength). Dose will not exceed 0.9 milligrams per kilogram of body weight once daily. |

EMGALITY (S)

Products Affected

- Emgality

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Episodic Migraines (EM) (initial): Diagnosis of EM. Patient has 4 to 14 migraine days per month, but no more than 14 headache days per month. Chronic Migraines (CM) (initial): Diagnosis of CM. Medication overuse headache has been considered and potentially offending medication(s) have been discontinued. Patient has greater than or equal to 15 headache days per month, of which at least 8 must be migraine days for at least 3 months. Episodic Cluster Headache (ECH) (initial): Diagnosis of episodic cluster headache. Patient has experienced at least 2 cluster periods lasting from 7 days to 365 days, separated by pain-free periods lasting at least three months. |
| Age Restrictions | EM, CM, ECH (initial): 18 years of age or older. |
| Prescriber Restrictions | EM, CM, ECH (initial, reauth): Prescribed by or in consultation with a neurologist or pain specialist. |
| Coverage Duration | EM, CM (initial): 6 months. ECH (initial): 3 months. EM, CM, ECH (reauth): 12 months. |
| Other Criteria | EM, CM (reauth): Patient has experienced a positive response to therapy, demonstrated by a reduction in headache frequency and/or intensity. Use of acute migraine medications (e.g., non-steroidal anti-inflammatory drugs [NSAIDs], triptans) has decreased since the start of CGRP therapy. CM (reauth): Patient continues to be monitored for medication overuse headache. ECH (reauth): Patient has experienced a positive response to therapy, demonstrated by a reduction in headache frequency and/or intensity. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

EMPLICITI (S)

Products Affected

- Empliciti

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Multiple myeloma: Diagnosis of multiple myeloma. One of the following: 1) Both of the following: a) Patient has received at least one prior therapy for multiple myeloma [eg, Revlimid (lenalidomide), Thalomid (thalidomide), Velcade (bortezomib)] and b) Used in combination with both of the following: Revlimid (lenalidomide) and dexamethasone, OR 2) Both of the following: a) Patient has received at least two prior therapies including Revlimid (lenalidomide) and a proteasome inhibitor and b) Used in combination with both of the following: Pomalyst (pomalidomide) and dexamethasone. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist/oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

ENBREL (S)

Products Affected

- Enbrel
- Enbrel Mini
- Enbrel Sureclick

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Rheumatoid Arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. Trial and failure, contraindication, or intolerance to one disease-modifying antirheumatic drug (DMARD) [eg, methotrexate (Rheumatrex/Trexall), Arava (leflunomide), Azulfidine (sulfasalazine)]. Polyarticular Juvenile Idiopathic Arthritis (PJIA) (Initial): Diagnosis of moderately to severely active PJIA. Trial and failure, contraindication, or intolerance to one of the following DMARDs: Arava (leflunomide) or methotrexate (Rheumatrex/Trexall). Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. Plaque psoriasis (Initial): Diagnosis of moderate to severe chronic plaque psoriasis. Ankylosing Spondylitis (AS) (Initial): Diagnosis of active AS. Trial and failure, contraindication, or intolerance to two NSAIDs. |
| Age Restrictions | N/A |
| Prescriber Restrictions | RA (initial), PJIA (initial), AS (initial): Prescribed by or in consultation with a rheumatologist. PsA (initial): Prescribed by or in consultation with a rheumatologist or dermatologist. Plaque Psoriasis (initial): Prescribed by or in consultation with a dermatologist. |
| Coverage Duration | All uses (initial, reauth): 12 months |
| Other Criteria | All indications (Reauth): Documentation of positive clinical response to Enbrel therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

ENDARI (S)

Products Affected

- Endari

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Sickle cell disease (initial): Diagnosis of sickle cell disease. Used to reduce acute complications of sickle cell disease. One of the following: (a) Patient is using Endari with concurrent hydroxyurea therapy, OR (b) Patient has a contraindication or intolerance to hydroxyurea. Patient has had 2 or more painful sickle cell crises within the past 12 months. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Sickle cell disease (initial): Prescribed by or in consultation with a hematologist/oncologist |
| Coverage Duration | Sickle cell disease (initial, reauth): 12 months |
| Other Criteria | Sickle cell disease (reauth): Documentation of positive clinical response to Endari therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

ENTYVIO (S)

Products Affected

- Entyvio

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Ulcerative Colitis (UC) (init): Diagnosis (Dx) of moderately to severely active UC. Trial and failure, contraindication, or intolerance (F/C/I) to one of the following conventional therapies: 6-mercaptopurine (Purinethol), aminosalicylates [eg, mesalamine (Asacol, Pentasa, Rowasa), olsalazine (Dipentum), sulfasalazine (Azulfidine, Sulfazine)], azathioprine (Imuran), corticosteroids (eg, prednisone, methylprednisolone). F/C/I to one tumor necrosis factor (TNF) inhibitor [eg, Humira (adalimumab), infliximab]. Crohn's Disease (CD) (init): Dx of moderately to severely active CD. F/C/I to one of the following medications: 6-mercaptopurine (Purinethol), azathioprine (Imuran), corticosteroids (eg, prednisone, methylprednisolone), methotrexate (Rheumatrex, Trexall). F/C/I to one TNF inhibitor [eg, Humira (adalimumab), infliximab]. |
| Age Restrictions | N/A |
| Prescriber Restrictions | UC, CD (init): Prescribed by or in consultation with a gastroenterologist |
| Coverage Duration | UC, CD (init): 14 weeks. UC, CD (reauth): 12 months. |
| Other Criteria | UC, CD (reauth): Documentation of positive clinical response to Entyvio therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

EPCLUSA NON-PREFERRED (S)

Products Affected

- Epclusa

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Criteria will be applied consistent with current AASLD/IDSA guideline. Diagnosis of chronic hepatitis C. One of the following: 1) Trial and failure, contraindication or intolerance to Mavyret (except patients with decompensated cirrhosis) AND sofosbuvir/velpatasvir, OR 2) For continuation of prior brand Epclusa therapy. Patient is not receiving Epclusa in combination with another HCV direct acting antiviral agent [e.g., Sovaldi (sofosbuvir)]. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with one of the following: Hepatologist, Gastroenterologist, Infectious disease specialist, HIV specialist certified through the American Academy of HIV Medicine. |
| Coverage Duration | 12 to 24 weeks. Criteria will be applied consistent with current AASLD/IDSA guideline. |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

EPCLUSA PREFERRED (S)

Products Affected

- Sofosbuvir/velpatasvir

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Criteria will be applied consistent with current AASLD/IDSA guideline. Diagnosis of chronic hepatitis C. Patient is not receiving sofosbuvir/velpatasvir in combination with another HCV direct acting antiviral agent [e.g., Sovaldi (sofosbuvir)]. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with one of the following: Hepatologist, Gastroenterologist, Infectious disease specialist, HIV specialist certified through the American Academy of HIV Medicine. |
| Coverage Duration | 12 to 24 weeks. Criteria will be applied consistent with current AASLD/IDSA guideline. |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

EPIDIOLEX (S)

Products Affected

- Epidiolex

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Lennox-Gastaut syndrome (LGS): Diagnosis of seizures associated with LGS. Trial of, contraindication, or intolerance to two formulary anticonvulsants (e.g., topiramate, lamotrigine, valproate). Dravet syndrome (DS): Diagnosis of seizures associated with DS. |
| Age Restrictions | N/A |
| Prescriber Restrictions | LGS, DS: Prescribed by or in consultation with a neurologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

EPOPROSTENOL (S)

Products Affected

- Epoprostenol Sodium
- Veletri

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH. |
| Age Restrictions | N/A |
| Prescriber Restrictions | PAH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist. |
| Coverage Duration | PAH (Initial): 6 months. (Reauth): 12 months |
| Other Criteria | Subject to Part B vs D review. PAH (Reauth): Documentation of positive clinical response to therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ERBITUX (S)

Products Affected

- Erbitux

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Head and Neck Cancer: Diagnosis of locally or regionally advanced squamous cell head and neck cancer and used in combination with radiation therapy, or diagnosis of recurrent or metastatic squamous cell head and neck cancer and 1 of the following: trial and failure of platinum-based chemotherapy, or used in combination with 1 of the following: cisplatin (Platinol AQ), carboplatin (Paraplatin), cisplatin (Platinol AQ) plus 5-FU (Acrucil), or carboplatin (Paraplatin) plus 5-FU (Acrucil). Colorectal Cancer: Diagnosis of metastatic carcinoma of the colon or rectum. One of the following: Used in combination with either FOLFOX (fluorouracil, leucovorin, and oxaliplatin) or FOLFIRI (fluorouracil, leucovorin, and irinotecan), OR trial and failure or intolerance to irinotecan-based chemotherapy, oxaliplatin-based chemotherapy, or intensive therapy (eg, FOLFOX or FOLFIRI), OR used as monotherapy in patients not appropriate for intensive therapy. Tumor expresses wild-type KRAS gene and wild type NRAS gene. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months. |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ERIVEDGE (S)

Products Affected

- Erivedge

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Basal cell carcinoma: One of the following: A) Diagnosis of metastatic basal cell carcinoma OR B) Both of the following: 1) Diagnosis of locally advanced basal cell carcinoma AND 2) One of the following: a) Disease recurred following surgery or b) Patient is not a candidate for surgery and radiation. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or dermatologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ERLEADA (S)

Products Affected

- Erleada

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Non-metastatic castration-resistant or castration-recurrent prostate cancer (NM-CRPC): Diagnosis of non-metastatic, castration-resistant (chemical or surgical) or recurrent prostate cancer. Trial and failure or intolerance to Xtandi (enzalutamide) or Nubeqa (darolutamide). Metastatic castration-sensitive prostate cancer (M-CSPC): Diagnosis of metastatic, castration-sensitive prostate cancer. Trial and failure or intolerance to Xtandi (enzalutamide). |
| Age Restrictions | N/A |
| Prescriber Restrictions | NM-CRPC, M-CSPC: Prescribed by or in consultation with an oncologist or urologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

ESBRIET (S)

Products Affected

- Esbriet

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Idiopathic pulmonary fibrosis (IPF) (initial): Both of the following: 1) diagnosis of IPF as documented by all of the following: a) exclusion of other known causes of interstitial lung disease (ILD) (eg, domestic and occupational environmental exposures, connective tissue disease, drug toxicity), AND b) one of the following: i) in patients not subjected to surgical lung biopsy, the presence of a usual interstitial pneumonia (UIP) pattern on high-resolution computed tomography (HRCT) revealing IPF or probable IPF, OR ii) in patients subjected to a lung biopsy, both HRCT and surgical lung biopsy pattern revealing IPF or probable IPF. |
| Age Restrictions | N/A |
| Prescriber Restrictions | IPF (initial): Prescribed by or in consultation with a pulmonologist |
| Coverage Duration | initial, reauth: 12 months |
| Other Criteria | IPF (reauth): Documentation of positive clinical response to Esbriet therapy. |

EVENTITY (S)

Products Affected

- Evenity

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | <p>Diagnosis of postmenopausal osteoporosis. One of the following: Set I) Both of the following: A) Bone mineral density (BMD) T-score of -2.5 or lower in the lumbar spine, femoral neck, total hip, or radius (one-third radius site) AND B) One of the following: 1) history of low-trauma fracture of the hip, spine, proximal humerus, pelvis, or distal forearm, or 2) trial and failure, contraindication, or intolerance (TF/C/I) to one anti-resorptive treatment (e.g., alendronate, risedronate, zoledronic acid, Prolia [denosumab]), or Set II) Both of the following: A) BMD T-score between -1.0 and -2.5 in the lumbar spine, femoral neck, total hip, or radius (one-third radius site) AND B) One of the following: 1) history of low-trauma fracture of the hip, spine, proximal humerus, pelvis, or distal forearm, or 2) both of the following: i) TF/C/I to one anti-resorptive treatment (e.g., alendronate, risedronate, zoledronic acid, Prolia [denosumab]) and ii) one of the following FRAX (Fracture Risk Assessment Tool) 10-year probabilities: a) major osteoporotic fracture at 20% or more in the U.S., or the country-specific threshold in other countries or regions, or b) hip fracture at 3% or more in the U.S., or the country-specific threshold in other countries or regions. Trial of, contraindication, or intolerance to one of the following: Forteo (teriparatide) or Tymlos (abaloparatide).</p> <p>Treatment duration of Evenity (romosozumab-aqqg) has not exceeded a total of 12 months during the patient's lifetime.</p> |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months (max 12 months of therapy per lifetime) |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

EXONDYS 51 (S)

Products Affected

- Exondys 51

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Initial: Diagnosis of Duchenne muscular dystrophy (DMD). Documentation of a confirmed mutation of the dystrophin gene amenable to exon 51 skipping. Patient is ambulatory. Initial/Reauth: Exondys 51 dosing for DMD is in accordance with the United States Food and Drug Administration approved labeling: maximum dosing of 30 mg/kg infused once weekly. Patient's condition has been evaluated via the 6-minute walk test (6MWT) or North Star ambulatory assessment (NSAA) [documentation of the patient's most recent results must be provided]. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Initial/Reauth: Prescribed by or in consultation with a neurologist who has experience treating children |
| Coverage Duration | Initial: 6 months. Reauth: 12 months |
| Other Criteria | Reauth: One of the following: 1) All of the following: Patient has been on therapy for less than 12 months, patient is maintaining ambulatory status, and patient is tolerating therapy, OR 2) All of the following: Patient has been on therapy for 12 months or more, Patient is maintaining ambulatory status, patient has experienced a benefit from therapy (e.g., disease amelioration compared to untreated patients), and patient is tolerating therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

EYLEA (S)

Products Affected

- Eylea

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Diagnosis of one of the following: A) Neovascular (wet) age-related macular degeneration OR B) Macular edema following retinal vein occlusion, OR C) Diabetic macular edema OR D) Diabetic retinopathy in patients with diabetic macular edema. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

FABRAZYME (S)

Products Affected

- Fabrazyme

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Fabry Disease: Diagnosis of Fabry disease. Fabrazyme will not be used in combination with Galafold (migalastat). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | Fabry Disease: 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

FARYDAK (S)

Products Affected

- Farydak

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Multiple Myeloma (MM): Diagnosis of MM. Used in combination with both of the following: Velcade (bortezomib) and dexamethasone. Patient has received at least two prior treatment regimens which included both of the following: Velcade (bortezomib) and an immunomodulatory agent [eg, Revlimid (lenalidomide), Thalomid (thalidomide)]. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist/hematologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

FASENRA (S)

Products Affected

- Fasenra Pen

- Fasenra

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Asthma (initial): Diagnosis of severe asthma. Asthma is an eosinophilic phenotype as defined by a baseline (pre-treatment) peripheral blood eosinophil level greater than or equal to 150 cells per microliter. One of the following: 1) Patient has had at least one or more asthma exacerbations requiring systemic corticosteroids within the past 12 months, OR 2) Any prior intubation for an asthma exacerbation, OR 3) Prior asthma-related hospitalization within the past 12 months. Patient is currently being treated with one of the following unless there is a contraindication or intolerance to these medications: a) Both of the following: i) High-dose inhaled corticosteroid (ICS) [e.g., greater than 500 mcg fluticasone propionate equivalent/day] and ii) additional asthma controller medication [e.g., leukotriene receptor antagonist, long-acting beta-2 agonist (LABA), theophylline], OR b) One maximally-dosed combination ICS/LABA product [e.g., Advair (fluticasone propionate/salmeterol), Dulera (mometasone/formoterol), Symbicort (budesonide/formoterol)]. |
| Age Restrictions | Asthma (Initial): Patient is 12 years of age or older |
| Prescriber Restrictions | Asthma (Initial/Reauth): Prescribed by or in consultation with a pulmonologist or allergist/immunologist |
| Coverage Duration | Asthma (init): 6 months. Asthma (reauth): 12 months |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|-----------------------|---|
| Other Criteria | Asthma (Reauth): Documentation of positive clinical response to therapy (e.g., reduction in exacerbations, improvement in forced expiratory volume in 1 second [FEV1], decreased use of rescue medications). Patient is currently being treated with one of the following unless there is a contraindication or intolerance to these medications: a) Both of the following: i) inhaled corticosteroid (ICS) and ii) additional asthma controller medication [e.g., leukotriene receptor antagonist, long-acting beta-2 agonist (LABA), theophylline], OR b) A combination ICS/LABA product [e.g., Advair (fluticasone propionate/salmeterol), Dulera (mometasone/formoterol), Symbicort (budesonide/formoterol)]. |
|-----------------------|---|

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

FENTANYL (S)

Products Affected

- Fentanyl Citrate Oral Transmucosal

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | For the management of breakthrough cancer pain. Patient is currently taking a long-acting opioid around the clock for cancer pain. Patient must have at least a one week history of ONE of the following medications to demonstrate tolerance to opioids: Morphine sulfate at doses of greater than or equal to 60 mg/day, Fentanyl transdermal patch at doses greater than or equal to 25 µg/hr, Oxycodone at a dose of greater than or equal to 30 mg/day, Oral hydromorphone at a dose of greater than or equal to 8 mg/day, Oral oxymorphone at a dose of greater than or equal to 25 mg/day, or an alternative opioid at an equianalgesic dose (e.g., oral methadone greater than or equal to 20 mg/day). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with one of the following: Pain specialist, Oncologist, Hematologist, Hospice care specialist, or Palliative care specialist. |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

FERRIPROX (S)

Products Affected

- Ferriprox

- Ferriprox Twice-a-day

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Transfusional iron overload due to thalassemia syndromes (Initial): Diagnosis of transfusional iron overload due to thalassemia syndromes. Absolute neutrophil count (ANC) greater than $1.5 \times 10^9/L$. One of the following: A) Trial and failure, defined as serum ferritin greater than 2,500 mcg/L, to Desferal (deferoxamine), Exjade (deferasirox) or Jadenu (deferasirox) OR B) History of contraindication or intolerance to Desferal (deferoxamine), Exjade (deferasirox) or Jadenu (deferasirox). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | All uses (reauth): Patient has experienced greater than or equal to 20% decline in serum ferritin levels from baseline. ANC greater than $1.5 \times 10^9/L$. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

FIRAZYR (S)

Products Affected

- Firazyr

- Icatibant Acetate

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Treatment of hereditary angioedema (HAE) attacks: Diagnosis of HAE. For the treatment of acute HAE attacks. Not used in combination with other approved treatments for acute HAE attacks. |
| Age Restrictions | N/A |
| Prescriber Restrictions | HAE: Prescribed by or in consultation with an immunologist, allergist, or rheumatologist |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

FIRDAPSE (S)

Products Affected

- Firdapse

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Lambert-Eaton Myasthenic Syndrome (LEMS) (initial): Diagnosis of LEMS. |
| Age Restrictions | N/A |
| Prescriber Restrictions | LEMS (initial): Prescribed by or in consultation with a neurologist. |
| Coverage Duration | LEMS (initial): 3 months. LEMS (reauth): 12 months. |
| Other Criteria | LEMS (reauth): Documentation of positive clinical response to Firdapse therapy (e.g., improvement in dynamometry, Timed 25-Foot Walk Test, Timed Up and Go Test). |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

FIRMAGON (S)

Products Affected

- Firmagon INJ 120MG/VIAL, 80MG

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Diagnosis of advanced or metastatic prostate cancer. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

FOLOTYN (S)

Products Affected

- Folutyn

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Peripheral T-cell lymphoma: Diagnosis of relapsed or refractory PTCL. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist/oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

FORTEO (S)

Products Affected

- Forteo INJ 600MCG/2.4ML

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | <p>Postmenopausal osteoporosis or osteopenia or men with primary or hypogonadal osteoporosis or osteopenia: Diagnosis of one of the following: a) postmenopausal osteoporosis or osteopenia or b) primary or hypogonadal osteoporosis or osteopenia. One of the following: Set I) Both of the following: A) Bone mineral density (BMD) T-score of -2.5 or lower in the lumbar spine, femoral neck, total hip, or radius (one-third radius site) AND B) One of the following: 1) history of low-trauma fracture of the hip, spine, proximal humerus, pelvis, or distal forearm, or 2) trial and failure, contraindication, or intolerance (TF/C/I) to one osteoporosis treatment (e.g., alendronate, risedronate, zoledronic acid, Prolia [denosumab]), or Set II) Both of the following: A) BMD T-score between -1.0 and -2.5 in the lumbar spine, femoral neck, total hip, or radius (one-third radius site) AND B) One of the following: 1) history of low-trauma fracture of the hip, spine, proximal humerus, pelvis, or distal forearm, or 2) both of the following: i) TF/C/I to one osteoporosis treatment (e.g., alendronate, risedronate, zoledronic acid, Prolia [denosumab]) and ii) One of the following FRAX 10-year probabilities: a) Major osteoporotic fracture at 20% or more in the U.S., or the country-specific threshold in other countries or regions, or b) Hip fracture at 3% or more in the U.S., or the country-specific threshold in other countries or regions. Treatment duration of parathyroid hormones [e.g., Forteo (teriparatide), Tymlos (abaloparatide)] has not exceeded a total of 24 months during the patient's lifetime. Glucocorticoid-Induced Osteoporosis: See Other Criteria section.</p> |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|--------------------------|---|
| Coverage Duration | All uses: 24 months (max 24 months of therapy per lifetime) |
| Other Criteria | Glucocorticoid-Induced Osteoporosis: Diagnosis of glucocorticoid-induced osteoporosis. History of prednisone or its equivalent at a dose greater than or equal to 5mg/day for greater than or equal to 3 months. One of the following: 1) BMD T-score less than or equal to -2.5 based on BMD measurements from lumbar spine, femoral neck, total hip, or radius (one-third radius site), or 2) One of the following FRAX 10-year probabilities: a) Major osteoporotic fracture at 20% or more in the U.S., or the country-specific threshold in other countries or regions, or b) Hip fracture at 3% or more in the U.S., or the country-specific threshold in other countries or regions, or 3) History of one of the following fractures resulting from minimal trauma: vertebral compression fx, fx of the hip, fx of the distal radius, fx of the pelvis, or fx of the proximal humerus. TF/C/I to one bisphosphonate (e.g., alendronate). Treatment duration of parathyroid hormones [e.g., Forteo (teriparatide), Tymlos (abaloparatide)] has not exceeded a total of 24 months during the patient's lifetime. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

FULPHILA (S)

Products Affected

- Fulphila

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | <p>Febrile neutropenia (FN) prophylaxis: Patient will be receiving prophylaxis for FN due to one of the following: 1) Patient is receiving National Cancer Institute’s Breast Intergroup, INT C9741 dose dense chemotherapy protocol for primary breast cancer, 2) patient is receiving a dose-dense chemotherapy regimen for which the incidence of FN is unknown, 3) patient is receiving chemotherapy regimen(s) associated with greater than 20% incidence of FN, 4) both of the following: a) patient is receiving chemotherapy regimen(s) associated with 10-20% incidence of FN, AND b) patient has one or more risk factors associated with chemotherapy-induced infection, FN, or neutropenia, OR 5) both of the following: a) patient is receiving myelosuppressive anticancer drugs associated with neutropenia, AND b) patient has a history of FN or dose-limiting event during a previous course of chemotherapy (secondary prophylaxis). Treatment of FN (off-label): Patient has received or is receiving myelosuppressive anticancer drugs associated with neutropenia. Diagnosis of FN. Patient is at high risk for infection-associated complications. Acute radiation syndrome (ARS) (off-label): Patient was/will be acutely exposed to myelosuppressive doses of radiation (hematopoietic subsyndrome of ARS).</p> |
| Age Restrictions | N/A |
| Prescriber Restrictions | All uses: Prescribed by or in consultation with a hematologist/oncologist |
| Coverage Duration | ARS: 1 mo. FN (prophylaxis, treatment): 3 mo or duration of tx. |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

GALAFOLD (S)

Products Affected

- Galafold

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Fabry Disease (FD) (initial): Diagnosis of FD. Patient has an amenable galactosidase alpha gene (GLA) variant based on in vitro assay data. FD (initial, reauthorization): Galafold will not be used in combination with Fabrazyme (agalsidase beta). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | FD (initial, reauth): 12 months. |
| Other Criteria | FD (reauthorization): Documentation of positive clinical response to Galafold therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

GAMASTAN S/D (S)

Products Affected

- Gamastan
- Gamastan S/d

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | Contraindications to immune globulin therapy (i.e., IgA deficiency with antibodies to IgA and a history of hypersensitivity or product specific contraindication). |
| Required Medical Information | Immune globulin is being used intramuscularly. The immune globulin will be administered at the minimum effective dose and appropriate frequency for the prescribed diagnosis. Patient requires immunization for hepatitis A, measles, rubella, or varicella. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 3 months (Approve one dose only) |
| Other Criteria | Subject to Part B vs D review. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

GAMIFANT (S)

Products Affected

- Gamifant

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Primary Hemophagocytic Lymphohistiocytosis (HLH) (initial): Diagnosis of HLH. One of the following: 1) Disease is refractory, recurrent, or progressive, or 2) Trial and failure, contraindication, or intolerance to conventional HLH therapy (e.g., etoposide, dexamethasone, cyclosporine A, intrathecal methotrexate). HLH (initial, reauth): Patient has not received hematopoietic stem cell transplantation (HSCT). |
| Age Restrictions | N/A |
| Prescriber Restrictions | HLH (initial): Prescribed by or in consultation with a hematologist/oncologist. |
| Coverage Duration | HLH (initial, reauth): 6 months. |
| Other Criteria | HLH (reauth): Documentation of positive clinical response to Gamifant therapy (e.g., improvement in hemoglobin/lymphocyte/platelet counts, afebrile, normalization of inflammatory factors/markers). |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

GATTEX (S)

Products Affected

- Gattex

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Short Bowel Syndrome (SBS) (Initial): Diagnosis of SBS. Submission of medical records (e.g., chart notes, laboratory values) documenting that the patient is dependent on parenteral nutrition/intravenous (PN/IV) support for at least 12 months. |
| Age Restrictions | N/A |
| Prescriber Restrictions | SBS (Init, reauth): Prescribed by or in consultation with a gastroenterologist. |
| Coverage Duration | SBS (Init): 6 months. SBS (Reauth): 12 months. |
| Other Criteria | SBS (Reauth): Submission of medical records (e.g., chart notes, laboratory values) documenting that the patient has had a reduction in weekly parenteral nutrition/intravenous (PN/IV) support from baseline while on Gattex therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

GAZYVA (S)

Products Affected

- Gazyva

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Chronic lymphocytic leukemia (CLL): Diagnosis of CLL or small lymphocytic leukemia. Used in combination with chlorambucil. Patient is previously untreated for CLL. Follicular lymphoma (FL): One of the following: 1)All of the following: 1.1)Diagnosis of FL. 1.2) Patient has relapsed after or is refractory to a rituximab-containing regimen. 1.3) Both of the following: Used in combination with bendamustine and followed by Gazyva monotherapy. OR 2) All of the following: 2.1) Diagnosis of stage II bulky, III, or IV FL 2.2) Patient has not been treated with prior therapy 2.3) Both of the following: Used in combination with chemotherapy until patient has at least achieved a partial remission and followed by Gazyva monotherapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist/oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

GILENYA (S)

Products Affected

- Gilenya

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Multiple Sclerosis (MS): Diagnosis of a relapsing form of MS (eg, relapsing-remitting MS, secondary-progressive MS with relapses). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

GILOTRIF (S)

Products Affected

- Gilotrif

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Non-small cell lung cancer (NSCLC): A) Diagnosis of advanced or metastatic (stage IIIB or IV) NSCLC AND B) One of the following: 1) Both of the following: a) Tumors have non-resistant epidermal growth factor (EGFR) mutations as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) AND b) GILOTRIF will be used as first-line treatment, OR 2) All of the following: a) disease progressed after platinum-based chemotherapy and b) squamous NSCLC. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

GLATIRAMER ACETATE (S)

Products Affected

- Glatopra
- Glatiramer Acetate

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Multiple Sclerosis (MS): Diagnosis of a relapsing form of MS (eg, relapsing-remitting MS, secondary-progressive MS with relapses). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

GLEEVEC (S)

Products Affected

- Imatinib Mesylate

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | One of the following: A) Diagnosis of Philadelphia chromosome positive (Ph+)/BCR ABL-positive chronic myelogenous leukemia (CML) OR B) Ph+/BCR ABL+ acute lymphoblastic leukemia (ALL) OR C) Gastrointestinal stromal tumor (GIST) AND one of the following: 1) Patient has documented c-KIT (CD117) positive unresectable or metastatic malignant GIST, OR 2) Patient had resection of c-KIT (CD117) positive GIST and imatinib will be used as adjuvant therapy OR D) Dermatofibrosarcoma protuberans that is unresectable, recurrent, or metastatic OR E) Hypereosinophilic syndrome or chronic eosinophilic leukemia OR F) Myelodysplastic syndrome (MDS) or myeloproliferative disease associated with platelet-derived growth factor receptor gene rearrangements OR G) Aggressive systemic mastocytosis without the D816V c-KIT mutation or with c-KIT mutational status unknown. |
| Age Restrictions | N/A |
| Prescriber Restrictions | All uses: Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | All uses: 12 months |
| Other Criteria | All uses: Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

GOCOVRI (S)

Products Affected

- Gocovri

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Parkinson's disease (PD) (initial): Diagnosis of Parkinson's disease, patient is experiencing dyskinesia, patient is receiving levodopa-based therapy. Trial and failure or intolerance to amantadine immediate release and Osmolex (amantadine ER). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Initial: Prescribed by or in consultation with a neurologist |
| Coverage Duration | 12 months |
| Other Criteria | Parkinson's Disease (reauthorization): Documentation of positive clinical response to Gocovri therapy (e.g., decreased "off" periods or decreased "on" time with troublesome dyskinesia). |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

GROWTH HORMONE, NON-PREFERRED (S)

Products Affected

- Humatrope INJ 12MG, 24MG, 6MG
- Humatrope Combo Pack
- Norditropin Flexpro
- Norditropin Nordiflex Pen INJ 30MG/3ML
- Omnitrope INJ 10MG/1.5ML, 5MG/1.5ML
- Saizen
- Saizen Click.easy
- Saizenprep Reconstitutionkit

| PA Criteria | Criteria Details |
|---------------------------|-------------------------------------|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|--|--|
| <p>Required Medical Information</p> | <p>PGHD(initial):less than 4mo w/suspected GD based on clinical presentation (eg, persistent neonatal hypoglycemia, persistent/prolonged neonatal jaundice/elev bilirubin, male infant with microgenitalia, midline anatomical defects, failure to thrive),OR hx neonatal hypoglycemia assoc w/pituitary dz,or panhypopituitarism dx,or all of the following: PGHD dx [confrmd by ht (utilizing age and gender grwth charts related to ht) documented(doc) by ht more than 2.0SD below midparental ht or more than 2.25SD below population(pop) mean (below 1.2 percentile for age and gender),or grwth velocity more than 2SD below mean for age and gender, or delayed skeletal maturation more than 2SD below mean for age and gender (eg,delayed more than 2yrs compared w/chronological age)]. PWS(reauth):evidence of positive response to tx(eg,incr in total LBM, decr in fat mass) and expctd adult ht not attained and doc of expctd adult ht goal. GFSGA(initial):SGA dx based on catchup grwth failure in 1st 24mo of life using 0-36mo grwth chart confrmd by birth wt or length below 3rd percentile for gestational age(more than 2SD below pop mean) and ht remains at or below 3rd percentile (more than 2SD below pop mean). TS,NS(initial):ped grwth failure dx assoc w/TS w/doc female w/bone age less than 14yrs, or NS and ht below 5th percentile on grwth charts for age and gender. SHOX(initial):ped grwth failure dx w/SHOX gene deficiency confirmed by genetic testing. GFCRI(initial): ped grwth failure dx assoc w/CRI. ISS(initial):ISS dx, diagnostic eval excluded other causes assoc w/short stature(eg GHD, chronic renal insufficiency), doc ht at or below -2.25SD score below corresponding mean ht for age and gender assoc with growth rates unlikely to permit attainment of adult height in the normal range. PGHD,NS,SHOX,GFCRI,ISS (initial): doc male w/bone age less than 16yrs or female w/bone age less than 14yrs. PGHD,GFSGA,TS/NS,SHOX,GFCRI,ISS(reauth):expctd adult ht not attained and doc of expctd adult ht goal.</p> |
| <p>Age Restrictions</p> | <p>N/A</p> |
| <p>Prescriber Restrictions</p> | <p>PGHD, PWS, GFSGA, TS/NS, SHOX, AGHD, TPAP, IGHDA, ISS: prescribed by endocrinologist. GFCRI: prescribed by endocrinologist or nephrologist</p> |
| <p>Coverage Duration</p> | <p>All uses (initial, reauth): 12 months</p> |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|-----------------------|--|
| Other Criteria | <p>Trial and failure or intolerance to Genotropin and Nutropin.</p> <p>AGHD(initial):dx of AGHD with clin records supporting dx of childhood-onset GHD, or adult-onset GHD w/clin records doc hormone deficiency d/t hypothalamic-pituitary dz from organic or known causes (eg,damage from surgery, cranial irradiation, head trauma, subarachnoid hemorrhage) and pt has 1GH stim test (insulin tolerance test [ITT], arginine/GHRH,glucagon,arginine) to confirm adult GHD w/peak GH values ([ITT at or below 5mcg/L],[GHRH+ARG at or below 11mcg/L if BMI less than 25kg/m2, at or below 8mcg/L if BMI at or above 25 and below 30kg/m2, or at or below 4mcg/L if BMI at or above 30kg/m2],[glucagon at or below 3mcg/L],[Arg at or below 0.4mcg/L],[macimorelin less than 2.8 ng/mL 30, 45, 60 and 90 mins following macimorelin administration]) or doc deficiency of 3 anterior pituitary hormones (prolactin,ACTH,TSH,FSH/LH) and IGF-1/somatomedinC below age and gender adjstd nrml range as provided by physicians lab. AGHD,IGHDA(reauth):monitoring as demonstrated by doc w/in past 12mo of IGF-1/somatomedinC level.</p> <p>TransitionPhaseAdolescent Pts(TPAP)(initial): attained expctd adult ht or closed epiphyses on bone radiograph, and doc high risk of GHD d/t GHD in childhood (from embryopathic/congenital defects, genetic mutations, irreversible structural hypothalamic-pituitary dz, panhypopituitarism, or deficiency of 3 anterior pituitary hormones: ACTH,TSH,prolactin,FSH/LH), w/IGF-1/somatomedinC below age and gender adj nrml range as provided by physicians lab, or pt does not have low IGF-1/somatomedinC and d/c GH tx for at least 1mo, and pt has 1 GH stim test (ITT,GHRH+ARG,ARG,glucagon) after d/c of tx for at least 1mo w/peak GH value [ITT at or below 5mcg/L], [GHRH+ARG at or below 11mcg/L if BMI less than 25kg/m2, at or below 8mcg/L if BMI at or above 25 and below 30kg/m2, or at or below 4mcg/L if BMI at or above 30kg/m2], [glucagon at or below 3mcg/L], [Arg at or below 0.4mcg/L], or at low risk of severe GHD(eg d/t isolated and/or idiopathic GHD) and d/c GH tx for at least 1mo, and pt has 1 GH stim test (ITT, GHRH+ARG, ARG, glucagon) after d/c of tx for at least 1mo w/corresponding peak GH value [ITT at or below 5mcg/L], [GHRH+ARG at or below 11mcg/L if BMI less than 25kg/m2, at or below 8mcg/L if BMI at or above 25 and below 30kg/m2, or at or below 4mcg/L if BMI at or above 30kg/m2], [glucagon at or below 3mcg/L], [Arg at or below 0.4mcg/L]. TPAP(reauth): evidence of positive response to therapy (eg,incr in total lean body mass, exercise capacity or IGF-1 and IGFBP-3). IGHDA(initial):doc GHD by failure to produce peak serum GH greater than 5 mcg/L after 2 provocative pharmacol stim tests(insulin,L-ARG,glucagon).</p> |
|-----------------------|--|

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

GROWTH HORMONE, PREFERRED (S)

Products Affected

- Genotropin
- Genotropin Miniquick
- Nutropin Aq Nuspin 10
- Nutropin Aq Nuspin 20
- Nutropin Aq Nuspin 5
- Nutropin Aq Pen

| PA Criteria | Criteria Details |
|---------------------------|-------------------------------------|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|--|--|
| <p>Required Medical Information</p> | <p>PGHD(initial):less than 4mo w/suspected GD based on clinical presentation (eg, persistent neonatal hypoglycemia, persistent/prolonged neonatal jaundice/elev bilirubin, male infant with microgenitalia, midline anatomical defects, failure to thrive),OR hx neonatal hypoglycemia assoc w/pituitary dz,or panhypopituitarism dx,or all of the following: PGHD dx [confrmd by ht (utilizing age and gender grwth charts related to ht) documented(doc) by ht more than 2.0SD below midparental ht or more than 2.25SD below population(pop) mean (below 1.2 percentile for age and gender),or grwth velocity more than 2SD below mean for age and gender, or delayed skeletal maturation more than 2SD below mean for age and gender (eg,delayed more than 2yrs compared w/chronological age)]. PWS(reauth):evidence of positive response to tx(eg,incr in total LBM, decr in fat mass) and expctd adult ht not attained and doc of expctd adult ht goal. GFSGA(initial):SGA dx based on catchup grwth failure in 1st 24mo of life using 0-36mo grwth chart confrmd by birth wt or length below 3rd percentile for gestational age(more than 2SD below pop mean) and ht remains at or below 3rd percentile (more than 2SD below pop mean). TS,NS(initial):ped grwth failure dx assoc w/TS w/doc female w/bone age less than 14yrs, or NS and ht below 5th percentile on grwth charts for age and gender. SHOX(initial):ped grwth failure dx w/SHOX gene deficiency confirmed by genetic testing. GFCRI(initial): ped grwth failure dx assoc w/CRI. ISS(initial):ISS dx, diagnostic eval excluded other causes assoc w/short stature(eg GHD, chronic renal insufficiency), doc ht at or below -2.25SD score below corresponding mean ht for age and gender assoc with growth rates unlikely to permit attainment of adult height in the normal range. PGHD,NS,SHOX,GFCRI,ISS (initial): doc male w/bone age less than 16yrs or female w/bone age less than 14yrs. PGHD,GFSGA,TS/NS,SHOX,GFCRI,ISS(reauth):expctd adult ht not attained and doc of expctd adult ht goal.</p> |
| <p>Age Restrictions</p> | <p>N/A</p> |
| <p>Prescriber Restrictions</p> | <p>PGHD, PWS, GFSGA, TS/NS, SHOX, AGHD, TPAP, IGHDA, ISS: prescribed by endocrinologist. GFCRI: prescribed by endocrinologist or nephrologist</p> |
| <p>Coverage Duration</p> | <p>All uses (initial, reauth): 12 months</p> |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|-----------------------|--|
| Other Criteria | <p>AGHD(initial):dx of AGHD with clin records supporting dx of childhood-onset GHD, or adult-onset GHD w/clin records doc hormone deficiency d/t hypothalamic-pituitary dz from organic or known causes (eg,damage from surgery, cranial irradiation, head trauma, subarachnoid hemorrhage) and pt has 1GH stim test (insulin tolerance test [ITT], arginine/GHRH,glucagon,arginine) to confirm adult GHD w/peak GH values ([ITT at or below 5mcg/L],[GHRH+ARG at or below 11mcg/L if BMI less than 25kg/m2, at or below 8mcg/L if BMI at or above 25 and below 30kg/m2, or at or below 4mcg/L if BMI at or above 30kg/m2],[glucagon at or below 3mcg/L],[Arg at or below 0.4mcg/L],[macimorelin less than 2.8 ng/mL 30, 45, 60 and 90 mins following macimorelin administration]) or doc deficiency of 3 anterior pituitary hormones (prolactin,ACTH,TSH,FSH/LH) and IFG-1/somatomedinC below age and gender adjstd nrml range as provided by physicians lab. AGHD,IGHDA(reauth):monitoring as demonstrated by doc w/in past 12mo of IGF-1/somatomedinC level.</p> <p>TransitionPhaseAdolescent Pts(TPAP)(initial): attained expctd adult ht or closed epiphyses on bone radiograph, and doc high risk of GHD d/t GHD in childhood (from embryopathic/congenital defects, genetic mutations, irreversible structural hypothalamic-pituitary dz, panhypopituitarism, or deficiency of 3 anterior pituitary hormones: ACTH,TSH,prolactin,FSH/LH), w/IGF-1/somatomedinC below age and gender adj nrml range as provided by physicians lab, or pt does not have low IGF-1/somatomedinC and d/c GH tx for at least 1mo, and pt has 1 GH stim test (ITT,GHRH+ARG,ARG,glucagon) after d/c of tx for at least 1mo w/peak GH value [ITT at or below 5mcg/L], [GHRH+ARG at or below 11mcg/L if BMI less than 25kg/m2, at or below 8mcg/L if BMI at or above 25 and below 30kg/m2, or at or below 4mcg/L if BMI at or above 30kg/m2], [glucagon at or below 3mcg/L], [Arg at or below 0.4mcg/L], or at low risk of severe GHD(eg d/t isolated and/or idiopathic GHD) and d/c GH tx for at least 1mo, and pt has 1 GH stim test (ITT, GHRH+ARG, ARG, glucagon) after d/c of tx for at least 1mo w/corresponding peak GH value [ITT at or below 5mcg/L], [GHRH+ARG at or below 11mcg/L if BMI less than 25kg/m2, at or below 8mcg/L if BMI at or above 25 and below 30kg/m2, or at or below 4mcg/L if BMI at or above 30kg/m2], [glucagon at or below 3mcg/L], [Arg at or below 0.4mcg/L]. TPAP(reauth): evidence of positive response to therapy (eg,incr in total lean body mass, exercise capacity or IGF-1 and IGFBP-3). IGHDA(initial):doc GHD by failure to produce peak serum GH greater than 5 mcg/L after 2 provocative pharmacol stim tests(insulin,L-ARG,glucagon).</p> |
|-----------------------|--|

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

H.P. ACTHAR GEL (S)

Products Affected

- Acthar

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Infantile Spasm (West Syndrome): Diagnosis of infantile spasms (West Syndrome). Dosing for infantile spasms (West Syndrome) is in accordance with the United States Food and Drug Administration (FDA) approved labeling: not to exceed 150U/m ² daily. Multiple Sclerosis (MS): Acute exacerbations of MS. Dosing for multiple sclerosis is in accordance with the United States FDA approved labeling: not to exceed 120 units once daily. Rheumatic disorders: As adjunctive therapy for short-term administration in: psoriatic arthritis, rheumatoid arthritis, including juvenile rheumatoid arthritis (selected cases may require low-dose maintenance therapy), ankylosing spondylitis. Collagen diseases: During an exacerbation or as maintenance therapy in selected cases of: systemic lupus erythematosus, systemic dermatomyositis (polymyositis). Dermatologic diseases: Severe erythema multiforme, Stevens-Johnson syndrome. Allergic states: Serum sickness. Ophthalmic diseases: Severe acute and chronic allergic and inflammatory processes involving the eye and its adnexa, such as: keratitis, iritis and iridocyclitis, diffuse posterior uveitis and choroiditis, optic neuritis, chorioretinitis, anterior segment inflammation. Respiratory diseases: Symptomatic sarcoidosis. Edematous state: To induce a diuresis or a remission of proteinuria in the nephrotic syndrome without uremia of the idiopathic type or that due to lupus erythematosus. All indications except for infantile spasms: Trial and failure, contraindication, or intolerance to treatment with two corticosteroids. All indications (except infantile spasms, multiple sclerosis): Dosing is in accordance with the United States FDA approved labeling: not to exceed 80 units per day. |
| Age Restrictions | Infantile spasms: less than 2 years old |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|--------------------------------|---|
| Prescriber Restrictions | Infantile Spasm, MS: neurologist. Rheumatic disorder, collagen disease: rheumatologist. Dermatologic: dermatologist. Allergic state: allergist, immunologist. Ophthalmic disease: optometrist, ophthalmologist. Respiratory diseases: pulmonologist. Edematous state: nephrologist, rheumatologist. |
| Coverage Duration | Infantile Spasms: 4 weeks. MS: 3 weeks. All other FDA-approved uses: 3 months. |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

HAEGARDA (S)

Products Affected

- Haegarda

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Prophylaxis of hereditary angioedema (HAE) attacks: Diagnosis of HAE. For prophylaxis against HAE attacks. |
| Age Restrictions | N/A |
| Prescriber Restrictions | HAE: Prescribed by or in consultation with an immunologist, allergist, or rheumatologist |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

HALAVEN (S)

Products Affected

- Halaven

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Breast Cancer: Diagnosis of recurrent or metastatic breast cancer. Previous treatment with both of the following: one anthracycline [eg, doxorubicin, Ellence (epirubicin)] and one taxane [eg, paclitaxel, Taxotere (docetaxel)]. Liposarcoma: Diagnosis of unresectable or metastatic liposarcoma. Previous treatment with one anthracycline-containing regimen. |
| Age Restrictions | N/A |
| Prescriber Restrictions | All uses: prescribed by or in consultation with an oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

HARVONI (S)

Products Affected

- Harvoni
- Ledipasvir/sofosbuvir

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Criteria will be applied consistent with current AASLD/IDSA guideline. All (including patients with genotype 5 or 6 infection AND decompensated cirrhosis): A) Diagnosis of chronic hepatitis C, B) Patient is not receiving ledipasvir/sofosbuvir in combination with another HCV direct acting antiviral agent [eg, Sovaldi (sofosbuvir)]. ONE of the following: Trial and failure, intolerance, or contraindication to a) Mavyret (except patients with decompensated cirrhosis, or pediatric patients between 3 years and 11 years of age and weighing less than 45 kg) and b) sofosbuvir/velpatasvir (except pediatric patients 3 years to 5 years of age and weighing less than 17 kg), OR for continuation of prior ledipasvir/sofosbuvir therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with one of the following: Hepatologist, Gastroenterologist, Infectious disease specialist, HIV specialist certified through the American Academy of HIV Medicine. |
| Coverage Duration | 12 to 24 weeks. Criteria will be applied consistent with current AASLD/IDSA guideline. |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

HERCEPTIN (S)

Products Affected

- Herceptin

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Breast cancer: Diagnosis of human epidermal growth factor receptor 2 (HER2)-overexpressing breast cancer. One of the following treatment regimens: a) As adjuvant treatment, b) metastatic disease and one of the following: 1) used in combination with a taxane (eg, docetaxel, paclitaxel), or 2) used as a single agent in a patient who has received one or more chemotherapy regimens for metastatic disease, or c) used in combination with Perjeta (pertuzumab). Gastric Cancer: Diagnosis of HER2-overexpressing gastric or gastroesophageal junction adenocarcinoma (locally advanced, recurrent, or metastatic). Used in combination with one of the following treatment regimens: a) Adrucil (5-fluorouracil), or b) Platinol (cisplatin) and Xeloda (capecitabine). |
| Age Restrictions | N/A |
| Prescriber Restrictions | All uses: Prescribed by or in consultation with an oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

HETLIOZ (S)

Products Affected

- HetlioZ

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Non-24-Hour Sleep-Wake Disorder (Non-24) (initial): Both of the following: 1) Diagnosis of non-24-hour sleep-wake disorder (also known as free-running disorder, free-running or non-entrained type circadian rhythm sleep disorder, or hypernycthemeral syndrome), AND 2) patient is totally blind (has no light perception). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Non-24 (initial): Prescribed by or in consultation with a specialist in sleep disorders or neurologist |
| Coverage Duration | Non-24 (initial): 6 mo. (reauth): 12 mo |
| Other Criteria | Non-24 (reauth): Documentation of positive clinical response to HetlioZ therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

HIZENTRA (S)

Products Affected

- Hizentra

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | All uses (initial, reauth): Contraindications to immune globulin therapy (i.e., IgA deficiency with antibodies to IgA and a history of hypersensitivity or product specific contraindication). Patient does not have hyperprolinemia. |
| Required Medical Information | Initial: Immune globulin will be administered at the minimum effective dose and appropriate frequency for the prescribed diagnosis. Medication is being used subcutaneously. Diagnosis of chronic inflammatory demyelinating polyneuropathy (CIDP) OR one of the following FDA-approved or literature supported diagnoses: 1) Common variable immunodeficiency (CVID), OR 2) Congenital agammaglobulinemia (X-linked or autosomal recessive), OR 3) Severe combined immunodeficiencies (SCID), OR 4) Wiskott-Aldrich syndrome, OR 5) Other primary immunodeficiency with an immunologic evaluation including IgG levels below the normal laboratory value for the patient's age at the time of diagnosis and the patient lacks an adequate response to protein and polysaccharide antigens (i.e., tetanus toxoid or diphtheria toxoid and pneumovax or HiB vaccine). |
| Age Restrictions | N/A |
| Prescriber Restrictions | All uses (initial, reauth): Prescribed by or in consultation with a physician who has specialized expertise in managing patients on SCIG therapy (e.g., immunologist, hematologist, neurologist). |
| Coverage Duration | Initial, reauth: 12 months |
| Other Criteria | Subject to Part B vs. Part D review. All uses (reauth): Patient has experienced an objective improvement on immune globulin therapy and the immune globulin will be administered at the minimum effective dose (by decreasing the dose, increasing the frequency, or implementing both strategies) for maintenance therapy. |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

HRM - ANTIHISTAMINES

Products Affected

- Cyproheptadine Hcl SYRP
- Cyproheptadine Hydrochloride TABS
- Dexchlorpheniramine Maleate SOLN
- Dexchlorpheniramine Maleate SYRP
- Hydroxyzine Hcl INJ 25MG/ML
- Hydroxyzine Hcl SYRP
- Hydroxyzine Hcl TABS 50MG
- Hydroxyzine Hydrochloride INJ
- Hydroxyzine Hydrochloride TABS 10MG, 25MG
- Hydroxyzine Pamoate CAPS
- Phenadoz
- Phenergan SUPP
- Promethazine Hcl INJ
- Promethazine Hcl SUPP
- Promethazine Hcl TABS 12.5MG
- Promethazine Hcl Plain
- Promethazine Hydrochloride INJ
- Promethazine Hydrochloride TABS 25MG, 50MG
- Promethazine Vc Plain SOLN
- Promethazine/phenylephrine
- Promethegan

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication. |
| Age Restrictions | PA applies to patients 65 years or older |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

HRM - ANTIPSYCHOTICS

Products Affected

- Thioridazine Hcl TABS 100MG, 10MG, 25MG, 50MG

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | The drug is being used for an FDA-approved diagnosis AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication. Trial and failure, contraindication or intolerance to one of the following: haloperidol, fluphenazine, or an atypical antipsychotic. |
| Age Restrictions | PA applies to patients 65 years or older |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | Applies to New Starts only. |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

HRM - BUTALBITAL

Products Affected

- Ascomp/codeine
- Butalbital Compound/codeine CAPS 325MG; 50MG; 40MG; 30MG
- Butalbital/acetaminophen
- Butalbital/acetaminophen/caffeine CAPS
- Butalbital/acetaminophen/caffeine TABS 325MG; 50MG; 40MG
- Butalbital/acetaminophen/caffeine/codeine
- Butalbital/aspirin/caffeine CAPS
- Butalbital/aspirin/caffeine/codeine
- Capacet
- Cephadyn
- Esgic CAPS
- Margesic
- Marten-tab
- Phrenilin Forte CAPS 300MG; 50MG; 40MG
- Tencon TABS 325MG; 50MG
- Vanatol Lq
- Vtol Lq
- Zebutal CAPS 325MG; 50MG; 40MG

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication. |
| Age Restrictions | PA applies to patients 65 years or older |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

HRM - ENDOCRINE

Products Affected

- Megestrol Acetate SUSP
- Megestrol Acetate TABS

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication. |
| Age Restrictions | PA applies to patients 65 years or older |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | Applies to New Starts only. |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

HRM - PHENOBARBITAL, PENTOBARBITAL

Products Affected

- Pentobarbital Sodium INJ
- Phenobarbital ELIX 20MG/5ML
- Phenobarbital TABS 100MG, 15MG, 16.2MG, 30MG, 32.4MG, 60MG, 64.8MG, 97.2MG
- Phenobarbital Sodium INJ 130MG/ML, 65MG/ML

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | The drug is being used for an FDA-approved diagnosis AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication. |
| Age Restrictions | PA applies to patients 65 years or older |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | Applies to New Starts only. |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

HRM - SKELETAL MUSCLE RELAXANTS

Products Affected

- Chlorzoxazone TABS 250MG, 500MG
- Cyclobenzaprine Hydrochloride TABS
- Methocarbamol TABS
- Orphenadrine Citrate Er

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication. |
| Age Restrictions | PA applies to patients 65 years or older |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

HRM - TCA

Products Affected

- Amitriptyline Hcl TABS 100MG, 150MG, 25MG, 75MG
- Amitriptyline Hydrochloride TABS 10MG, 50MG
- Chlordiazepoxide/amitriptyline
- Doxepin Hcl CAPS 100MG, 10MG, 150MG, 50MG, 75MG
- Doxepin Hcl CONC
- Doxepin Hydrochloride CAPS 25MG
- Perphenazine/amitriptyline

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication . |
| Age Restrictions | PA applies to patients 65 years or older |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | Applies to New Starts only. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

HUMIRA (S)

Products Affected

- Humira
- Humira Pediatric Crohns Disease Starter Pack
- Humira Pen
- Humira Pen-cd/uc/hs Starter
- Humira Pen-ps/uv Starter

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Rheumatoid Arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. Trial and failure, contraindication, or intolerance (TF/C/I) to one disease modifying antirheumatic drug (DMARD) [eg, methotrexate (Rheumatrex/Trexall), Arava (leflunomide), Azulfidine (sulfasalazine)]. Juvenile Idiopathic Arthritis (JIA) (Initial): Diagnosis of moderately to severely active polyarticular JIA. TF/C/I to one of the following DMARDs: Arava (leflunomide) or methotrexate (Rheumatrex/Trexall). Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. Plaque psoriasis (Initial): Diagnosis of moderate to severe chronic plaque psoriasis. Ankylosing Spondylitis (AS) (Initial): Diagnosis of active AS. TF/C/I to two NSAIDs. Crohn's Disease (CD) (Initial): Diagnosis of moderately to severely active CD. TF/C/I to one of the following conventional therapies: 6-mercaptopurine, Imuran (azathioprine), corticosteroid (eg, prednisone, methylprednisolone), methotrexate (Rheumatrex/Trexall), or failure (ie, lost response) or intolerance to infliximab. Ulcerative Colitis (UC) (Initial): Diagnosis of moderately to severely active UC. TF/C/I to one of the following conventional therapies: 6-mercaptopurine, Imuran (azathioprine), corticosteroid (eg, prednisone, methylprednisolone), aminosalicylate [eg, mesalamine (Asacol, Pentasa, Rowasa), Dipentum (olsalazine), Azulfidine (sulfasalazine)]. Hidradenitis suppurativa (Initial): Diagnosis of moderate to severe hidradenitis suppurativa (ie, Hurley Stage II or III). Uveitis (initial): Diagnosis of non-infectious uveitis, classified as intermediate, posterior, or panuveitis. |
| Age Restrictions | N/A |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|--------------------------------|---|
| Prescriber Restrictions | RA, AS, JIA (initial): Prescribed by or in consultation with a rheumatologist. PsA (initial): Prescribed by or in consultation with a dermatologist or rheumatologist. Plaque Psoriasis, HS (initial): Prescribed by or in consultation with a dermatologist. CD, UC (initial): Prescribed by or in consultation with a gastroenterologist. Uveitis (initial): Prescribed by or in consultation with an ophthalmologist or rheumatologist. |
| Coverage Duration | UC (Initial): 12 wks. UC (reauth): 12 mo. All other indications (initial, reauth): 12 mo. |
| Other Criteria | RA, JIA, PsA, Plaque psoriasis, AS, CD, Hidradenitis suppurativa (HS), Uveitis (Reauth): Documentation of positive clinical response to Humira therapy. UC (Reauth): For patients who initiated Humira therapy within the past 12 weeks: Documentation of clinical remission or significant clinical benefit by eight weeks (Day 57) of therapy OR For patients who have been maintained on Humira therapy for longer than 12 weeks: Documentation of positive clinical response to Humira therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

HYDROXYPROGESTERONE (S)

Products Affected

- Hydroxyprogesterone Caproate INJ
1.25GM/5ML

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | All uses (initial): Pregnant patients. |
| Required Medical Information | Amenorrhea/Abnormal Uterine Bleeding: Diagnosis of one of the following: 1) primary or secondary amenorrhea or 2) abnormal uterine bleeding. Amenorrhea or abnormal uterine bleeding is due to hormonal imbalance in the absence of organic pathology (e.g., submucous fibroids or uterine cancer). Secretory endometrium and desquamation: Used for production of secretory endometrium and desquamation in patients with endometrial disorder. Adenocarcinoma: Diagnosis of Stage III or IV adenocarcinoma of the uterine corpus. Estrogen production test: Used for the testing of endogenous estrogen production. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Adenocarcinoma (initial): Prescribed by or in consultation with an oncologist |
| Coverage Duration | Amenorrhea/Abnormal Uterine Bleeding: 4 mo. Estrogen testing: 2 mo. All other uses: 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

IBRANCE (S)

Products Affected

- Ibrance

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Breast Cancer: Diagnosis of advanced or metastatic breast cancer. Disease is a) hormone receptor (HR)-positive, and b) human epidermal growth factor receptor 2 (HER2)-negative. One of the following: 1) used in combination with an aromatase inhibitor (e.g., anastrozole, letrozole, exemestane) and one of the following: a) patient is a male, or b) patient is a postmenopausal woman, OR 2) both of the following: used in combination with Faslodex (fulvestrant) and disease has progressed following endocrine therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ICLUSIG (S)

Products Affected

- Iclusig

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Chronic myelogenous leukemia: Diagnosis of chronic myelogenous leukemia AND One of the following: A) Trial and failure, resistance, relapse, contraindication, or intolerance to at least TWO other tyrosine kinase inhibitors (e.g., Gleevec [imatinib], Sprycel, Tassigna, and Bosulif) or B) Patient has the T315I mutation. Acute Lymphoblastic Leukemia: Diagnosis of Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph+ ALL) AND One of the following: A) Trial and failure, resistance relapse, contraindication, or intolerance to at least TWO other FDA-approved tyrosine kinase inhibitors (e.g., Gleevec [imatinib], Sprycel), or B) Patient has the T315I mutation. |
| Age Restrictions | N/A |
| Prescriber Restrictions | All uses: Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | All uses: 12 months |
| Other Criteria | All uses: Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

IDHIFA (S)

Products Affected

- Idhifa

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Acute Myeloid Leukemia (AML): Diagnosis of AML. Disease is relapsed or refractory. Patient has an isocitrate dehydrogenase-2 (IDH2) mutation as detected by a U.S. Food and Drug Administration (FDA)-approved test (e.g., Abbott RealTime IDH2 assay) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist/oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

ILARIS (S)

Products Affected

- Ilaris

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Periodic Fever Syndromes (Cryopyrin-Associated Periodic Syndromes (CAPS), Tumor Necrosis Factor Receptor Associated Periodic Syndrome (TRAPS), Hyperimmunoglobulin D Syndrome (HIDS)/Mevalonate Kinase Deficiency (MKD), Familial Mediterranean Fever (FMF)). (Initial) Diagnosis of one of the autoinflammatory Periodic Fever Syndromes: CAPS (including Familial Cold Auto-inflammatory Syndrome (FCAS) and/or Muckle-Wells Syndrome (MWS)), TRAPS, HIDS/MKD, or FMF, AND The medication will not be used in combination with another biologic agent. Systemic juvenile idiopathic arthritis (sJIA) (Initial): Diagnosis of active sJIA AND The medication will not be used in combination with another biologic. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Periodic Fever Syndromes (CAPS, TRAPS, HIDS/MKD, FMF) (initial): Prescribed by or in consultation with an immunologist, allergist, dermatologist, rheumatologist, neurologist or other medical specialist |
| Coverage Duration | All uses (initial, reauth): 12 months |
| Other Criteria | SJIA (initial): Trial and failure, contraindication, or intolerance to non-steroidal anti-inflammatory drugs (NSAIDs), methotrexate, or corticosteroids. Periodic Fever Syndrome (CAPS, TRAPS, HIDS/MKD, FMF) and SJIA (Reauth): Documentation of positive clinical response to therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

ILUMYA (S)

Products Affected

- Ilumya

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Plaque Psoriasis (initial): Diagnosis of moderate-to-severe plaque psoriasis. One of the following: set A) Both of the following: 1) Trial and failure, contraindication, or intolerance to Enbrel (etanercept) or Humira (adalimumab) AND 2) Trial and failure, contraindication, or intolerance to Cosentyx (secukinumab), or set B) For continuation of prior Ilumya therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Plaque Psoriasis (initial): Prescribed by or in consultation with a dermatologist |
| Coverage Duration | Initial, reauth: 12 months |
| Other Criteria | Plaque Psoriasis (Reauth): Documentation of positive clinical response to Ilumya therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

IMBRUVICA (S)

Products Affected

- Imbruvica

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Mantle cell lymphoma (MCL): Diagnosis of MCL AND patient has relapsed or is refractory to at least one prior therapy for the treatment of MCL. Chronic lymphocytic leukemia (CLL): Diagnosis of CLL. Waldenstrom's macroglobulinemia: Diagnosis of Waldenstrom's macroglobulinemia/lymphoplasmacytic lymphoma. Small lymphocytic lymphoma (SLL): Diagnosis of SLL. Marginal zone lymphoma (MZL): Diagnosis of MZL AND patient has received at least one prior anti-CD20-based therapy for MZL [e.g., Rituxan (rituximab), Zevalin (ibritumomab), Gazyva (obinutuzumab, etc.)]. Chronic graft versus host disease (cGVHD): Diagnosis of cGVHD AND trial and failure of one or more lines of systemic therapy (e.g., corticosteroids, mycophenolate). |
| Age Restrictions | N/A |
| Prescriber Restrictions | All uses (except chronic graft versus host disease): Prescribed by or in consultation with an oncologist or hematologist. Chronic graft versus host disease: Prescribed by or in consultation with a hematologist, oncologist, or physician experienced in the management of transplant patients. |
| Coverage Duration | All Uses: 12 months |
| Other Criteria | All Uses: Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

IMFINZI (S)

Products Affected

- Imfinzi

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Urothelial carcinoma: 1) Diagnosis of locally advanced or metastatic urothelial carcinoma AND 2) One of the following: a) Patient has experienced disease progression during or following platinum-containing chemotherapy OR b) Patient has experienced disease progression within 12 months of neoadjuvant or adjuvant treatment with platinum-containing chemotherapy. Non-Small Cell Lung Cancer (NSCLC): 1) Diagnosis of NSCLC AND 2) Disease is stage III and unresectable AND 3) Disease has not progressed following concurrent platinum-based chemotherapy and radiation therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

INBRIJA (S)

Products Affected

- Inbrija

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Parkinson's disease (PD) (initial): Diagnosis of PD. Patient is experiencing intermittent OFF episodes. Patient is receiving Inbrija in combination with carbidopa/levodopa. Trial and failure, contraindication or intolerance to one of the following: MAO-B inhibitor (e.g., rasagiline, selegiline), dopamine agonist (e.g., pramipexole, ropinirole), or COMT Inhibitor (e.g., entacapone). |
| Age Restrictions | N/A |
| Prescriber Restrictions | PD (initial): Prescribed by or in consultation with a neurologist |
| Coverage Duration | PD (initial, reauth): 12 months |
| Other Criteria | PD (reauth): Documentation of positive clinical response to Inbrija therapy. Patient is receiving Inbrija in combination with carbidopa/levodopa. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

INCRELEX (S)

Products Affected

- Increlex

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Insulin-like Growth Factor-1 (IGF-1) deficiency (initial): Diagnosis of severe primary IGF-1 deficiency. Height standard deviation score of -3.0 or less. Basal IGF-1 standard deviation score of -3.0 or less. Normal or elevated growth hormone (GH). GH gene deletion (initial): Diagnosis of GH gene deletion in patients who have developed neutralizing antibodies to GH. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Initial: Prescribed by or in consultation with a pediatric endocrinologist |
| Coverage Duration | Initial, reauth: 12 months |
| Other Criteria | (Reauth): Evidence of positive response to therapy. |

INFLECTRA (S)

Products Affected

- Inflectra

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Crohn's Disease (CD) and Fistulizing Crohn's Disease (FCD) (initial): Dx of moderately to severely active CD or FCD. Ulcerative colitis (UC) (initial): Dx of moderately to severely active UC. Rheumatoid arthritis (RA) (initial): Diagnosis (Dx) of moderately to severely active RA. Ankylosing spondylitis (AS) (initial): Dx of active AS. Psoriatic arthritis (PsA) (initial): Dx of active PsA. Plaque psoriasis (initial): Dx of chronic severe (ie, extensive and/or disabling) plaque psoriasis. CD, FCD (initial): Trial and failure, contraindication or intolerance (TF/C/I) to one of the following conventional therapies: 6-mercaptopurine (Purinethol), azathioprine (Imuran), corticosteroids (eg, prednisone, methylprednisolone), methotrexate (Rheumatrex, Trexall). UC (initial): TF/C/I to one of the following conventional therapies: corticosteroids, aminosalicylate [eg, mesalamine (Asacol/Pentasa/Rowasa), olsalazine (Dipentum), sulfasalazine (Azulfidine/Sulfazine)], azathioprine (Imuran), 6-mercaptopurine (Purinethol). RA (initial): Receiving concurrent therapy with methotrexate (Rheumatrex/Trexall), or TF/C/I to methotrexate. AS (initial): TF/C/I to two or more NSAIDs. All indications (initial): One of the following: 1) Trial and failure or intolerance to Remicade or 2) For continuation of prior Inflectra therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | RA, AS: Prescribed or recommended by a rheumatologist. PsA: Prescribed or recommended by rheumatologist or dermatologist. Crohn's Disease, Fistulizing Crohn's Disease, UC: Prescribed or recommended by a gastroenterologist. Plaque Psoriasis: Prescribed or recommended by a dermatologist. |
| Coverage Duration | All uses (initial, reauth): 12 months |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|-----------------------|--|
| Other Criteria | Reauth (all indications): Documentation of positive clinical response to infliximab therapy. |
|-----------------------|--|

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

INGREZZA (S)

Products Affected

- Ingrezza

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Tardive Dyskinesia (initial): Diagnosis of moderate to severe tardive dyskinesia. One of the following: a) patient has persistent symptoms of tardive dyskinesia despite a trial of dose reduction, tapering, or discontinuation of the offending medication OR b) patient is not a candidate for a trial of dose reduction, tapering, or discontinuation of the offending medication. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Initial: Prescribed by or in consultation with a neurologist or psychiatrist. |
| Coverage Duration | Initial: 3 months. Reauth: 12 months |
| Other Criteria | Tardive Dyskinesia (reauth): Documentation of positive clinical response to Ingrezza therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

INLYTA (S)

Products Affected

- Inlyta

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Renal cell carcinoma (RCC): Diagnosis of RCC. One of the following: (1) disease has relapsed or (2) diagnosis of stage IV disease. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

INREBIC (S)

Products Affected

- Inrebic

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Myelofibrosis: Diagnosis of one of the following: primary myelofibrosis, post-polycythemia vera myelofibrosis, or post-essential thrombocythemia myelofibrosis. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist/oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

INTRON A (S)

Products Affected

- Intron A
- Intron A W/diluent INJ 10MU

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Chronic hepatitis B: Diagnosis of chronic hepatitis B infection, and patient is without decompensated liver disease. Chronic Hepatitis C: Diagnosis of chronic hepatitis C, patient without decompensated liver disease, patient has not previously been treated with interferon, and one of the following - 1) used in combination with ribavirin or 2) contraindication or intolerance to ribavirin. Metastatic renal cell carcinoma (RCC): diagnosis of metastatic RCC, used in combination with Avastin (bevacizumab). Other: diagnosis of condylomata acuminata (genital or perianal), diagnosis of hairy cell leukemia, diagnosis of AIDS-related Kaposi's sarcoma, diagnosis of malignant melanoma, diagnosis of Stage III or IV follicular Non-Hodgkin's Lymphoma, as maintenance therapy for the treatment of multiple myeloma. |
| Age Restrictions | N/A |
| Prescriber Restrictions | RCC: Prescribed by or in consultation with an oncologist. |
| Coverage Duration | HepB, HepC: 48 wks. Condylomata acuminata (genital or perianal): 6 wks. Other: 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

IRESSA (S)

Products Affected

- Iressa

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Non-small cell lung cancer (NSCLC): Diagnosis of metastatic NSCLC AND Patient has known active epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutations as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

ISOTRETINOIN (S)

Products Affected

- Amnesteem
- Claravis
- Isotretinoin CAPS
- Myorisan
- Zenatane

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Acne (initial): Diagnosis of acne. One of the following: A) Prescribed by a dermatologist or, B) Trial and failure, contraindication or intolerance to an adequate trial (at least 6 weeks) on both of the following conventional therapy regimens: a) topical retinoid or retinoid-like agent [eg, Retin-A/Retin-A Micro (tretinoin)] AND b) combination therapy with benzoyl peroxide and one of the following: 1) oral antibiotic [eg, Ery-Tab (erythromycin), Minocin (minocycline)] OR 2) topical antibiotic [eg, Cleocin-T (clindamycin), erythromycin, BenzaClin (benzoyl peroxide/clindamycin), Benzamycin (benzoyl peroxide/erythromycin)]. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | Acne (initial): 5 months. Acne (reauth): Retreatment - 5 months, Dose Titration - 1 month |
| Other Criteria | Acne, Retreatment (reauth): After more than 2 months off therapy, persistent or recurring acne is still present. Acne, Dose Titration (reauth): Confirmation that the total cumulative dose is less than 150 mg/kg. |

ISTODAX (S)

Products Affected

- Istodax
- Istodax (overfill)
- Romidepsin

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Cutaneous T-cell lymphoma (CTCL): Diagnosis of CTCL. Trial and failure, contraindication, or intolerance to at least one systemic therapy for the treatment of CTCL [e.g., Trexall (methotrexate), Targretin (bexarotene), Cytoxan (cyclophosphamide)] Peripheral T-cell lymphoma (PTCL): Diagnosis of PTCL. Trial and failure, contraindication, or intolerance to at least one therapy for the treatment of PTCL (e.g., conventional chemotherapy such as CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone), CHOEP (cyclophosphamide, doxorubicin, vincristine, etoposide, prednisone), Dose-adjusted EPOCH (etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin), HyperCVAD (cyclophosphamide, vincristine, doxorubicin, dexamethasone) alternating with high-dose methotrexate and cytarabine). |
| Age Restrictions | N/A |
| Prescriber Restrictions | CTCL, PTCL: Prescribed by or in consultation with an oncologist/hematologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ISTURISA (S)

Products Affected

- Isturisa

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Cushing's disease (initial): Diagnosis of Cushing's disease. One of the following: a) Patient is not a candidate for pituitary surgery, OR b) Pituitary surgery has not been curative for the patient. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Cushing's disease (initial): Prescribed by or in consultation with an endocrinologist. |
| Coverage Duration | Cushing's disease (initial, reauth): 12 months |
| Other Criteria | Cushing's disease (reauth): Documentation of positive clinical response to therapy (e.g., a clinically meaningful reduction in 24-hour urinary free cortisol levels, improvement in signs or symptoms of the disease). |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

IVIG (S)

Products Affected

- Asceniv
- Bivigam
- Carimune Nanofiltered INJ 12GM, 6GM
- Flebogamma Dif
- Gammagard Liquid
- Gammagard S/d Iga Less Than 1mcg/ml
- Gammaked
- Gammaplex
- Gamunex-c
- Octagam INJ 10GM/100ML, 10GM/200ML, 1GM/20ML, 2.5GM/50ML, 20GM/200ML, 25GM/500ML, 2GM/20ML, 5GM/100ML, 5GM/50ML
- Panzyga
- Privigen

| PA Criteria | Criteria Details |
|---------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | All uses (initial, reauth): Contraindications to immune globulin therapy (i.e., IgA deficiency with antibodies to IgA and a history of hypersensitivity or product specific contraindication). Privigen only: Hyperprolinemia. Octagam only: Allergy to corn. Gammaplex only: Hereditary intolerance to fructose. Infants for whom sucrose or fructose tolerance has not been established. |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|--|--|
| <p>Required Medical Information</p> | <p>Initial: Immune globulin (Ig) will be administered at the minimum effective dose and appropriate frequency for the prescribed diagnosis. For IVIG – Ig is being used intravenously (IV) AND One of the following diagnoses: [A] Primary Immunodeficiency 1) Common variable immunodeficiency. 2) Congenital agammaglobulinemia (X-linked or autosomal recessive). 3) Severe combined immunodeficiencies. 4) Wiskott-Aldrich syndrome. OR 5) Other PI with an immunologic evaluation including IgG levels below the normal laboratory value for the patient’s age at the time of diagnosis and the patient lacks an adequate response to protein and polysaccharide antigens (i.e., tetanus toxoid or diphtheria toxoid and pneumovax or HiB vaccine). [B] Secondary Acquired Antibody Deficiency 1) B-cell chronic lymphocytic leukemia with an Ig level less than 500 mg/dL OR history of recurrent bacterial infections. 2) HIV infection with an Ig level less than 400 mg/dL OR Patient has active bleeding or a platelet count less than 10 x 10⁹/L. 3) Multiple myeloma in plateau phase and patient has hypogammaglobulinemia. [C] Hematological Autoimmune Disorders 1) Acquired (pure) red cell aplasia (PRCA) that is immunologic and patient had a trial and failure, contraindication, or intolerance (TF/C/I) to a corticosteroid and an immunosuppressant (i.e., cyclophosphamide, cyclosporine) OR patient has viral PRCA caused by parvovirus B19. 2) Fetal alloimmune thrombocytopenia. 3) Hemolytic disease of the newborn and the patient has established hyperbilirubinemia. 4) Idiopathic thrombocytopenic purpura and patient had a TF/C/I to a corticosteroid OR a platelet count less than 30,000 cells/mm³. Continued in Other Criteria Section.</p> |
| <p>Age Restrictions</p> | <p>HIV (initial): patient is less than or equal to 12 years of age.</p> |
| <p>Prescriber Restrictions</p> | <p>All uses (initial, reauth): Prescribed by or in consultation with a physician who has specialized expertise in managing patients on immune globulin therapy (e.g., immunologist, hematologist, neurologist).</p> |
| <p>Coverage Duration</p> | <p>4 months: Solid organ transplant. 12 months: all other diagnoses.</p> |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|-----------------------|---|
| Other Criteria | <p>[D] Neuromuscular Autoimmune Disorders 1) Chronic inflammatory demyelinating polyneuropathy. 2) Guillain-Barré syndrome. 3) Inflammatory myopathies (dermatomyositis or polymyositis) AND Patient had a TF/C/I to a corticosteroid AND an immunosuppressant (i.e., azathioprine, methotrexate, cyclosporine A, cyclophosphamide, or tacrolimus). 4) Lambert-Eaton myasthenic syndrome AND Patient had a TF/C/I to a corticosteroid AND an immunosuppressant (e.g., azathioprine). 5) Multifocal motor neuropathy. 6) Myasthenia gravis with severe exacerbations or myasthenic crises AND Patient had a TF/C/I to a corticosteroid AND an immunosuppressant (i.e., azathioprine, cyclosporine, cyclophosphamide, or mycophenolate mofetil). 7) Stiff person syndrome AND Patient had a TF/C/I to at least 2 standard therapies (i.e., benzodiazepines, muscle relaxants, or anti-convulsants).</p> <p>[E] Other Disorders 1) Autoimmune blistering disease AND Patient had a TF/C/I to a corticosteroid AND an immunosuppressant (i.e., cyclophosphamide, dapsone, methotrexate, azathioprine, or mycophenolate mofetil). 2) Kawasaki syndrome. 3) Solid organ transplant and IVIG is being used for CMV prophylaxis, or patient is a kidney transplant recipient and has donor specific antibodies, or patient has steroid-resistant rejection and had a TF/C/I to standard therapies. For SCIG (Gamunex-C, Gammagard Liquid, Gammaked only)- Immune globulin is being used subcutaneously AND One of the following PI diagnoses: 1) Common variable immunodeficiency. 2) Congenital agammaglobulinemia (X-linked or autosomal recessive). 3) Severe combined immunodeficiencies. 4) Wiskott-Aldrich syndrome. OR 5) Other PI with an immunologic evaluation including IgG levels below the normal laboratory value for the patient's age at the time of diagnosis and patient lacks an adequate response to protein and polysaccharide antigens (i.e., tetanus toxoid or diphtheria toxoid and pneumovax or HiB vaccine). All products: Subject to Part B vs. Part D review. For non-oncology renewal, the patient has experienced an objective improvement on immune globulin therapy and the immune globulin will be administered at the minimum effective dose (by decreasing the dose, increasing the frequency, or implementing both strategies) for maintenance therapy.</p> |
|-----------------------|---|

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

JAKAFI (S)

Products Affected

- Jakafi

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Myelofibrosis: Diagnosis of primary myelofibrosis, OR post-polycythemia vera myelofibrosis, OR post-essential thrombocythemia myelofibrosis. Polycythemia vera: Diagnosis of polycythemia vera, AND trial and failure, contraindication, or intolerance to hydroxyurea. Acute graft versus host disease (aGVHD): Diagnosis of aGVHD. Disease is steroid-refractory. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Myelofibrosis, Polycythemia vera: Prescribed by or in consultation with a hematologist/oncologist. Acute graft versus host disease: Prescribed by or in consultation with one of the following: hematologist, oncologist, physician experienced in the management of transplant patients. |
| Coverage Duration | 12 months. |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

JEVTANA (S)

Products Affected

- Jevtana

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Prostate Cancer: Diagnosis of castration-resistant metastatic prostate cancer AND patient has been previously treated with a docetaxel-containing regimen AND patient is receiving concurrent prednisone. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

JUXTAPID (S)

Products Affected

- Juxtapid

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Homozygous familial hypercholesterolemia (HoFH) (initial): Submission of medical records (eg, chart notes, laboratory values) documenting diagnosis of HoFH as confirmed by one of the following: a) genetic confirmation of 2 mutations in the LDL receptor, ApoB, PCSK9, or LDL receptor adaptor protein 1 (ie, LDLRAP1 or ARH), or b) both of the following: 1) either untreated LDL-C greater than 500 mg/dL or treated LDL-C greater than 300 mg/dL AND 2) either xanthoma before 10 years of age or evidence of heterozygous FH in both parents. Patient is receiving other lipid-lowering therapy. Trial and failure, contraindication, or intolerance to Repatha therapy. Not used in combination with a proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor. |
| Age Restrictions | N/A |
| Prescriber Restrictions | HoFH (initial, reauth): Prescribed by a cardiologist, endocrinologist, or lipid specialist. |
| Coverage Duration | HoFH (initial): 6 months. (reauth): 12 months |
| Other Criteria | HoFH (reauthorization): Patient continues to receive other lipid-lowering therapy. Submission of medical records (eg, chart notes, laboratory values) documenting LDL-C reduction while on Juxtapid therapy. Not used in combination with a proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

KADCYLA (S)

Products Affected

- Kadcyła

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Breast cancer: A) Diagnosis of human epidermal growth factor receptor 2 (HER2)-positive recurrent or metastatic breast cancer AND B) Patient has been previously treated with Herceptin (trastuzumab) and a taxane (eg, docetaxel, paclitaxel). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

KALBITOR (S)

Products Affected

- Kalbitor

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Treatment of hereditary angioedema (HAE) attacks: Diagnosis of HAE. For the treatment of acute HAE attacks. Not used in combination with other approved treatments for acute HAE attacks. |
| Age Restrictions | 12 years of age or older |
| Prescriber Restrictions | HAE: Prescribed by or in consultation with an immunologist, allergist, or rheumatologist |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

KALYDECO (S)

Products Affected

- Kalydeco

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Cystic Fibrosis (CF) (Initial): Diagnosis of cystic fibrosis. Patient has one of the following mutations on at least one allele in the cystic fibrosis transmembrane conductance regulator gene as detected by a U.S. Food and Drug Administration (FDA)-cleared cystic fibrosis mutation test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA): A455E, A1067T, D110E, D110H, D579G, D1152H, D1270N, E56K, E193K, E831X, F1052V, F1074L, G178R, G551D, G551S, G1069R, G1244E, G1349D, K1060T, L206W, P67L, R74W, R117C, R117H, R347H, R352Q, R1070Q, R1070W, S549N, S549R, S945L, S977F, S1251N, S1255P, 711+3A-G, 2789+5G-A, 3272-26A-G, or 3849+10kbC-T. |
| Age Restrictions | CF (Initial): 6 months of age or older |
| Prescriber Restrictions | CF (initial): Prescribed by or in consultation with a specialist affiliated with a CF care center or pulmonologist |
| Coverage Duration | CF (initial, reauth): 12 months |
| Other Criteria | CF (Reauth): Documentation of positive clinical response (i.e. improvement in lung function [percent predicted forced expiratory volume in one second (PPFEV1)], decreased number of pulmonary exacerbations) to Kalydeco therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

KANUMA (S)

Products Affected

- Kanuma

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Lysosomal acid lipase deficiency: Diagnosis of lysosomal acid lipase deficiency (LAL-D). Diagnosis was confirmed by an enzymatic blood (e.g., dried blood spot test) or genetic test. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a specialist experienced in the treatment of inborn errors of metabolism, gastroenterologist, or lipidologist |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

KEVEYIS (S)

Products Affected

- Keveyis

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Periodic paralysis (Initial): Diagnosis of one of the following: Primary hyperkalemic periodic paralysis, Primary hypokalemic periodic paralysis, or Paramyotonia Congenita with periodic paralysis. |
| Age Restrictions | N/A |
| Prescriber Restrictions | All uses (initial): Prescribed by or in consultation with a neurologist |
| Coverage Duration | All uses (Initial): 3 months. (Reauth): 12 months |
| Other Criteria | All uses (Reauth): Documentation of positive clinical response to Keveyis therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

KEVZARA (S)

Products Affected

- Kevzara

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Rheumatoid Arthritis (RA) (initial): Diagnosis of moderately to severely active RA. One of the following: a) Trial and failure, contraindication, or intolerance (TF/C/I) to both Enbrel (etanercept) and Humira (adalimumab), b) or attestation demonstrating a trial may be inappropriate, OR c) For continuation of prior Kevzara therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Initial: Prescribed by or in consultation with a rheumatologist |
| Coverage Duration | Initial, Reauth: 12 months |
| Other Criteria | RA (reauth): Documentation of positive clinical response to Kevzara therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

KEYTRUDA (S)

Products Affected

- Keytruda

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | N/A |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

KINERET (S)

Products Affected

- Kineret

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Rheumatoid Arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. One of the following: Trial and failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR for continuation of prior Kineret therapy. Neonatal-Onset Multisystem Inflammatory Disease (NOMID) (initial): Diagnosis of NOMID AND dx of NOMID has been confirmed by one of the following: 1) NLRP-3 (nucleotide-binding domain, leucine rich family (NLR), pyrin domain containing 3) gene (also known as Cold-Induced Auto-inflammatory Syndrome-1 [CIAS1]) mutation OR 2) Both of the following: a) two of the following clinical symptoms: urticaria-like rash, cold/stress triggered episodes, sensorineural hearing loss, musculoskeletal symptoms (e.g., arthralgia, arthritis, myalgia), chronic aseptic meningitis, or skeletal abnormalities (e.g., epiphyseal overgrowth, frontal bossing) AND b) elevated acute phase reactants (eg, erythrocyte sedimentation rate [ESR], C-reactive protein [CRP], serum amyloid A [SAA]). |
| Age Restrictions | N/A |
| Prescriber Restrictions | RA (initial): Prescribed by or in consultation with a rheumatologist. NOMID (initial): Prescribed by or in consultation with allergist/immunologist or rheumatologist or pediatrician. |
| Coverage Duration | All Uses (initial, reauth): 12 months |
| Other Criteria | All Uses (Reauth): Documentation of positive clinical response to Kineret therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

KISQALI (S)

Products Affected

- Kisqali

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Breast cancer: Diagnosis of advanced or metastatic breast cancer. Disease is hormone receptor (HR)-positive and human epidermal growth factor receptor 2 (HER2)-negative. One of the following: A) Kisqali is used in combination with an aromatase inhibitor [e.g., Femara (letrozole)] OR B) Used in combination with Faslodex (fulvestrant). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

KISQALI-FEMARA PACK (S)

Products Affected

- Kisqali Femara 200 Dose
- Kisqali Femara 400 Dose
- Kisqali Femara 600 Dose

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Breast cancer: Diagnosis of advanced or metastatic breast cancer. Cancer is hormone receptor (HR)-positive and human epidermal growth factor receptor 2 (HER2)-negative. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

KORLYM (S)

Products Affected

- Korlym

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Cushing's syndrome (Initial): Diagnosis of endogenous Cushing's syndrome (i.e., hypercortisolism is not a result of chronic administration of high dose glucocorticoids). Diagnosis of either type 2 diabetes mellitus or diagnosis of glucose intolerance. Patient has either failed surgery or patient is not a candidate for surgery. Patient is not pregnant. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Initial: Prescribed by or in consultation with an endocrinologist. |
| Coverage Duration | Initial, reauth: 6 months |
| Other Criteria | Reauth: Documentation of one of the following: patient has improved glucose tolerance while on Korlym therapy or patient has stable glucose tolerance while on Korlym therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

KOSELUGO (S)

Products Affected

- Koselugo

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Neurofibromatosis Type 1 (NF1): Diagnosis of NF1. Patient has plexiform neurofibromas that are both of the following: inoperable and causing significant morbidity (e.g., disfigurement, motor dysfunction, pain, airway dysfunction, visual impairment). Patient is able to swallow a capsule whole. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with one of the following: oncologist or neurologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

KRYSTEXXA (S)

Products Affected

- Krystexxa

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Gout (initial): Diagnosis of treatment failure gout characterized by one of the following symptoms: a) greater than or equal to 3 flares in previous 18 months, b) greater than or equal to 1 gout tophus or c) gouty arthritis. History of failure or intolerance to both of the following conventional therapies: xanthine oxidase inhibitor (ie., allopurinol, febuxostat) and uricosuric agent (e.g., probenecid). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | Gout (initial, reauth): 12 months |
| Other Criteria | Gout (reauth): Serum urate level has decreased since initiating therapy. Clinical improvement in the signs and symptoms of gout (e.g., decrease in tophi size or frequency of gouty flares per year from baseline or improvement in chronic arthropathy or quality of life). |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

KUVAN (S)

Products Affected

- Kuvan

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Phenylketonuria (PKU) (initial): Diagnosis of PKU. Patient will have blood Phe levels measured after 1 week of therapy (new starts to therapy only) and periodically for up to 2 months of therapy to determine response. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | PKU (Init): 2 months (Reauth): 12 months |
| Other Criteria | PKU (reauth): Patient has had an objective response to therapy, defined as a 30% or greater reduction in phenylalanine (Phe) blood levels from baseline. Patient will continue to have blood Phe levels measured periodically during therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

KYNAMRO (S)

Products Affected

- Kynamro

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Homozygous familial hypercholesterolemia (HoFH) (initial): Submission of medical records (eg, chart notes, laboratory values) documenting diagnosis of HoFH as confirmed by one of the following: a) genetic confirmation of 2 mutations in the LDL receptor, ApoB, PCSK9, or LDL receptor adaptor protein 1 (ie, LDLRAP1 or ARH) , or b) both of the following: 1) either untreated LDL-C greater than 500 mg/dL or treated LDL-C greater than 300 mg/dL AND 2) either xanthoma before 10 years of age or evidence of heterozygous FH in both parents. Patient is receiving other lipid-lowering therapy. Trial and failure, contraindication, or intolerance to Repatha therapy. Not used in combination with Juxtapid (lomitapide). Not used in combination with a proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor. |
| Age Restrictions | N/A |
| Prescriber Restrictions | HoFH (initial, reauth): Prescribed by a cardiologist, endocrinologist, or lipid specialist. |
| Coverage Duration | HoFH (initial): 6 months. (reauth): 12 months |
| Other Criteria | HoFH (reauthorization): Patient continues to receive other lipid-lowering therapy. Submission of medical records (eg, chart notes, laboratory values) documenting LDL-C reduction while on Kynamro therapy. Not used in combination with Juxtapid (lomitapide). Not used in combination with a proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

KYPROLIS (S)

Products Affected

- Kyprolis

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Multiple Myeloma (MM): Diagnosis of MM. Disease is relapsed or refractory. Patient has received at least one prior therapy for MM. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist/hematologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

LARTRUVO (S)

Products Affected

- Lartruvo

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Diagnosis of soft tissue sarcoma (STS). One of the following: Disease is not amenable to curative treatment with radiotherapy OR Disease is not amenable to curative treatment with surgery. Used in combination with doxorubicin for the first 8 cycles of treatment AND Physician is aware that the phase 3 ANNOUNCE trial does not support initiating treatment with Lartruvo. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

LEMTRADA (S)

Products Affected

- Lemtrada

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Multiple Sclerosis (MS): Diagnosis of a relapsing form of MS (eg, relapsing-remitting MS, secondary-progressive MS with relapses). One of the following: 1) Patient has not been previously treated with alemtuzumab, and failure after a trial of at least 4 weeks, contraindication, or intolerance to two of the following disease-modifying therapies for MS: Aubagio (teriflunomide), Avonex (interferon beta-1a), Betaseron (interferon beta-1b), Copaxone/Glatopa (glatiramer acetate), Extavia (interferon beta-1b), Gilenya (fingolimod), Mavenclad (cladribine), Mayzent (siponimod), Ocrevus (ocrelizumab), Plegridy (peginterferon beta-1a), Rebif (interferon beta-1a), Tecfidera (dimethyl fumarate), Tysabri (natalizumab), Zinbryta (daclizumab), or 2) Patient has previously received treatment with alemtuzumab, and at least 12 months have or will have elapsed since the most recent treatment course with alemtuzumab. Not used in combination with another disease-modifying therapy for MS. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | MS: 12 months, max 2 yrs of therapy. |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

LENVIMA (S)

Products Affected

- Lenvima 10 Mg Daily Dose
- Lenvima 12mg Daily Dose
- Lenvima 14 Mg Daily Dose
- Lenvima 18 Mg Daily Dose
- Lenvima 20 Mg Daily Dose
- Lenvima 24 Mg Daily Dose
- Lenvima 4 Mg Daily Dose
- Lenvima 8 Mg Daily Dose

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Differentiated thyroid cancer (DTC): Diagnosis of DTC. Disease is locally recurrent or metastatic. Patient has symptomatic or progressive disease. Disease is refractory to radioactive iodine treatment. Renal Cell Carcinoma (RCC): Diagnosis of advanced RCC. Treatment follows one prior anti-angiogenic therapy. Used in combination with everolimus. Hepatocellular Carcinoma (HCC): Diagnosis of HCC. One of the following: patient has metastatic disease, or patient has extensive liver tumor burden, or patient is inoperable by performance status or comorbidity (local disease or local disease with minimal extrahepatic disease only), or disease is unresectable. Endometrial Carcinoma (EC): Diagnosis of advanced endometrial carcinoma that is not microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR). Patient has disease progression following systemic therapy. Used in combination with Keytruda (pembrolizumab) therapy. Patient is not a candidate for curative surgery or radiation. |
| Age Restrictions | N/A |
| Prescriber Restrictions | DTC/RCC/EC: Prescribed by or in consultation with an oncologist. HCC: Prescribed by or in consultation with one of the following: oncologist, hepatologist, or gastroenterologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

LETAIRIS (S)

Products Affected

- Ambrisentan
- Letairis

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH. |
| Age Restrictions | N/A |
| Prescriber Restrictions | PAH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist. |
| Coverage Duration | PAH (Initial): 6 months. PAH (Reauth): 12 months |
| Other Criteria | PAH (Reauth): Documentation of positive clinical response to therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

LEUKINE (S)

Products Affected

- Leukine INJ 250MCG

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | <p>Bone marrow/stem cell transplant (BMSCT): One of the following: 1) patient has non-myeloid malignancies undergoing myeloablative chemotherapy followed by autologous or allogeneic BMT, OR 2) used for mobilization of hematopoietic progenitor cells into the peripheral blood for collection by leukapheresis, OR 3) patient has had a peripheral stem cell transplant (PSCT) and has received myeloablative chemotherapy.</p> <p>Acute myeloid leukemia (AML): Diagnosis of AML. Patient has completed induction or consolidation chemotherapy. Age greater than or equal to 55 years.</p> <p>Febrile Neutropenia (FN) Prophylaxis: Patient will be receiving prophylaxis for FN due to one of the following: 1) Patient is receiving National Cancer Institute’s Breast Intergroup, INT C9741 dose dense chemotherapy protocol for primary breast cancer, 2) patient is receiving a dose-dense chemotherapy regimen for which the incidence of FN is unknown, 3) patient is receiving chemotherapy regimen(s) associated with a greater than 20% incidence of FN, 4) both of the following: a) patient is receiving chemotherapy regimen(s) associated with 10-20% incidence of FN, AND b) patient has one or more risk factors associated with chemotherapy-induced infection, FN, or neutropenia, OR 5) Both of the following: a) patient is receiving myelosuppressive anticancer drugs associated with neutropenia, AND b) patient has a history of FN or dose-limiting event during a previous course of chemotherapy (secondary prophylaxis).</p> <p>Acute radiation syndrome (ARS): Patient was/will be acutely exposed to myelosuppressive doses of radiation (hematopoietic subsyndrome of ARS). Treatment of High-Risk FN: Patient has received or is receiving myelosuppressive anticancer drugs associated with neutropenia.</p> <p>Diagnosis of FN. Patient is at high risk for infection-associated complications.</p> <p>HIV-related neutropenia (HIVN): Patient is infected with HIV, and ANC less than or equal to 1000 (cells/mm³).</p> |
| Age Restrictions | N/A |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|--------------------------------|--|
| Prescriber Restrictions | HIVN: Prescribed by or in consultation with a hematologist/oncologist or infectious disease specialist. All other uses: Prescribed by or in consultation with a hematologist/oncologist. |
| Coverage Duration | BMSCT, AML, FN (prophylaxis, treatment): 3mo or duration of tx. HIVN: 6mo. ARS: 1 mo. |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

LIBTAYO (S)

Products Affected

- Libtayo

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Cutaneous Squamous Cell Carcinoma (CSCC): Diagnosis of CSCC. Disease is metastatic or locally advanced. Patient is not a candidate for curative surgery or curative radiation. |
| Age Restrictions | N/A |
| Prescriber Restrictions | CSCC: Prescribed by or in consultation with an oncologist or dermatologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

LIDOCAINE TOPICAL (S)

Products Affected

- 7t Lido Gel
- Glydo
- Lidocaine OINT
- Lidocaine And Tetracaine Cream
- Lidocaine Hcl EXTERNAL SOLN 4%
- Lidocaine Hcl GEL
- Lidocaine Hcl PRSY
- Lidocaine Hcl Jelly
- Lidocaine/prilocaine CREA
- Lidocaine-prilocaine-cream Base CREA 2.5%; 2.5%
- Pliaglis
- Premium Lidocaine

| PA Criteria | Criteria Details |
|------------------------------|-------------------------------------|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | N/A |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 3 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

LIDODERM (S)

Products Affected

- Lidocaine PTCH 5%

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Post-herpetic neuralgia: Diagnosis of post-herpetic neuralgia. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

LONSURF (S)

Products Affected

- Lonsurf

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Colorectal Cancer: Diagnosis of metastatic colorectal cancer AND trial and failure, contraindication, or intolerance to at least one component in the following: fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy (e.g., FOLFOX, FOLFIRI, FOLFOXIRI) AND trial and failure, contraindication, or intolerance to at least one anti-VEGF therapy (e.g., Avastin) AND One of the following: A) patient has KRAS wild-type tumors and trial and failure, contraindication, or intolerance to at least one anti-EGFR therapy (e.g., Vectibix, Erbitux) OR Patient has KRAS mutant tumors. Gastric/Gastroesophageal Junction Adenocarcinoma: Diagnosis of metastatic gastric cancer or diagnosis of metastatic gastroesophageal junction adenocarcinoma. Trial and failure, contraindication or intolerance to at least two of the following: fluopyrimidine-based chemotherapy (e.g. fluorouracil), Platinum-based chemotherapy (e.g., carboplatin, cisplatin, oxaliplatin), Taxane (e.g., docetaxel, paclitaxel) or irinotecan-based chemotherapy, HER2/neu-targeted therapy (e.g., trastuzumab) (if HER2 overexpression). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

LORBRENA (S)

Products Affected

- Lorbrena

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Non-small cell lung cancer (NSCLC): Diagnosis of NSCLC. Disease is metastatic and anaplastic lymphoma kinase (ALK)-positive. Metastatic disease has progressed on one of the following: 1) Xalkori (crizotinib) and at least one other ALK inhibitor [e.g., Alunbrig (brigatinib)], 2) Alecensa (alectinib) as the first ALK inhibitor therapy, or 3) Zykadia (ceritinib) as the first ALK inhibitor therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

LOTRONEX (S)

Products Affected

- Alosetron Hydrochloride

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Severe Diarrhea-Predominant Irritable Bowel Syndrome (IBS) in Women (initial): All of the following: 1) diagnosis of severe diarrhea-predominant IBS, 2) symptoms for at least 6 months, 3) female patient, AND 4) trial and failure, contraindication, or intolerance to an antidiarrheal agent [eg, loperamide]. |
| Age Restrictions | Initial: 18 years of age or older |
| Prescriber Restrictions | N/A |
| Coverage Duration | IBS (initial): 12 weeks. IBS (reauth): 6 mo. |
| Other Criteria | IBS (reauthorization): Symptoms of IBS continue to persist, AND documentation of positive clinical response to therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

LUMIZYME (S)

Products Affected

- Lumizyme
- Myozyme

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Pompe disease: Diagnosis of Pompe disease [acid alpha-glucosidase (GAA) deficiency]. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

LUMOXITI (S)

Products Affected

- Lumoxiti

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Hairy cell leukemia (HCL): Diagnosis of HCL. Disease is relapsed or refractory. Patient has received at least two prior systemic therapies, including treatment with a purine nucleoside analog (PNA). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist/hematologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

LUPANETA PACK (S)

Products Affected

- Lupaneta Pack

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Endometriosis (initial): Diagnosis of endometriosis. One of the following: Patient has had surgical ablation to prevent recurrence, or trial and failure, contraindication, or intolerance to one NSAID or one oral contraceptive. Trial and failure, contraindication, or intolerance to Lupron Depot (7.5 mg, 22.5 mg, 30 mg, and 45 mg). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | Endomet (init, reauth): 6 months |
| Other Criteria | Endometriosis (reauthorization): Patient is experiencing recurrence of symptoms after an initial course of therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

LUPRON (S)

Products Affected

- Leuprolide Acetate INJ

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Prostate Cancer: Diagnosis of advanced or metastatic prostate cancer. Central Precocious Puberty (CPP) (initial): Diagnosis of CPP (idiopathic or neurogenic). Early onset of secondary sexual characteristics in females less than age 8 or males less than age 9. Advanced bone age of at least one year compared with chronologic age. One of the following: a) patient has undergone gonadotropin-releasing hormone agonist (GnRHa) testing AND Peak luteinizing hormone (LH) level above pre-pubertal range, or b) patient has a random LH level in the pubertal range. |
| Age Restrictions | N/A |
| Prescriber Restrictions | CPP (initial, reauth): Prescribed by or in consultation with a pediatric endocrinologist. |
| Coverage Duration | CPP (initial, reauth), Prostate CA: 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

LUPRON DEPOT (S)

Products Affected

- Lupron Depot (1-month)
- Lupron Depot (3-month)
- Lupron Depot (4-month)
- Lupron Depot (6-month)

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Prostate Cancer (7.5 mg, 22.5 mg, 30 mg, 45 mg): Diagnosis of advanced or metastatic prostate cancer. Endometriosis (3.75 mg, 11.25 mg) (initial): Diagnosis of endometriosis. One of the following: Patient has had surgical ablation to prevent recurrence, or trial and failure, contraindication, or intolerance to one NSAID and one oral contraceptive. Uterine Leiomyomata (UL) (3.75 mg, 11.25 mg): a) For use prior to surgery to reduce size of fibroids to facilitate a surgical procedure (eg, myomectomy, hysterectomy) OR b) all of the following: treatment of anemia, anemia is caused by uterine leiomyomata (fibroids), and for use prior to surgery. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | Prostate CA: 12 mo. Endomet (init, reauth):6mo. UL (anemia):3 mo (fibroids):4 mo |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

LUPRON DEPOT PED (S)

Products Affected

- Lupron Depot-ped (1-month)
- Lupron Depot-ped (3-month)

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Central Precocious Puberty (CPP) (initial): Diagnosis of CPP (idiopathic or neurogenic). Early onset of secondary sexual characteristics in females less than age 8 or males less than age 9. Advanced bone age of at least one year compared with chronologic age. One of the following: a) patient has undergone gonadotropin-releasing hormone agonist (GnRHa) testing AND Peak luteinizing hormone (LH) level above pre-pubertal range, or b) patient has a random LH level in the pubertal range. |
| Age Restrictions | N/A |
| Prescriber Restrictions | CPP (initial, reauth): Prescribed by or in consultation with a pediatric endocrinologist. |
| Coverage Duration | CPP (init, reauth): 12 months |
| Other Criteria | CPP (reauthorization): LH levels have been suppressed to pre-pubertal levels. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

LYNPARZA (S)

Products Affected

- Lynparza CAPS

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Ovarian Cancer: Diagnosis of advanced ovarian cancer. Presence of deleterious or suspected deleterious germline BRCA-mutations as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Trial and failure, contraindication, or intolerance to three or more prior lines of chemotherapy (e.g., paclitaxel with cisplatin). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

LYNPARZA TABLET (S)

Products Affected

- Lynparza TABS

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Ovarian cancer, advanced disease with known or suspected BRCA mutation with 3 or more prior lines of chemotherapy: Diagnosis of advanced ovarian cancer. Presence of deleterious or suspected deleterious germline BRCA-mutations as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Trial and failure, contraindication, or intolerance to three or more prior lines of chemotherapy (e.g., paclitaxel with cisplatin). Maintenance treatment of recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer: Diagnosis of one of the following: recurrent epithelial ovarian cancer, recurrent fallopian tube cancer, or recurrent primary peritoneal cancer. Used for maintenance treatment in patients who are in a complete or partial response to platinum-based chemotherapy (e.g., cisplatin, carboplatin). Breast cancer: Diagnosis of metastatic breast cancer. Presence of a deleterious or suspected deleterious germline BRCA-mutation as detected by an FDA-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Disease is human epidermal growth factor receptor 2 (HER2)-negative. Patient has been previously treated with chemotherapy (e.g., anthracycline, taxane) in the neoadjuvant, adjuvant, or metastatic setting. One of the following: a) Disease is hormone receptor (HR) negative, or b) Disease is hormone receptor (HR)-positive and one of the following: i) patient has been treated with prior endocrine therapy or ii) patient is considered an inappropriate candidate for endocrine therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | All uses (except prostate cancer): Prescribed by or in consultation with an oncologist. Prostate cancer: Prescribed by or in consultation with an oncologist or urologist. |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|---------------------------------|--|
| <p>Coverage Duration</p> | <p>12 months</p> |
| <p>Other Criteria</p> | <p>First-line maintenance treatment of BRCA-mutated advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer: Diagnosis of one of the following: advanced epithelial ovarian cancer, advanced fallopian tube cancer, or advanced primary peritoneal cancer. Presence of deleterious or suspected deleterious BRCA-mutation as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Patient has had a complete or partial response to first-line platinum-based chemotherapy (e.g., carboplatin, cisplatin). Lynparza will be used as first-line maintenance treatment. Pancreatic adenocarcinoma: Diagnosis of metastatic pancreatic adenocarcinoma. Presence of a deleterious or suspected deleterious germline BRCA-mutation as detected by an FDA-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Disease has not progressed while receiving at least 16 weeks of a first-line platinum-based chemotherapy regimen (e.g., FOLFIRINOX, FOLFOX, etc.). First-line maintenance treatment of HRD-positive advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer in combination with bevacizumab: Diagnosis of advanced epithelial ovarian cancer, advanced fallopian tube cancer, or advanced primary peritoneal cancer. Cancer is associated with homologous recombination deficiency (HRD)-positive status (defined by either: a deleterious or suspected deleterious BRCA mutation, and/or genomic instability) as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Patient has had a complete or partial response to first-line platinum-based chemotherapy (e.g., carboplatin, cisplatin). Used in combination with bevacizumab (e.g., Avastin, Mvasi). Lynparza will be used as first-line maintenance treatment. Prostate cancer: Diagnosis of metastatic castration-resistant prostate cancer. Presence of deleterious or suspected deleterious homologous recombination repair (HRR) gene mutation as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Disease has progressed following prior treatment with one of the following: a) enzalutamide (Xtandi) or b) abiraterone (e.g., Zytiga, Yonsa). All indications: Approve for continuation of prior therapy.</p> |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

MAKENA (S)

Products Affected

- Hydroxyprogesterone Caproate INJ 250MG/ML
- Makena INJ 275MG/1.1ML

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Preterm birth prophylaxis: Patient had a previous singleton (single offspring) spontaneous preterm birth. Patient is having a singleton pregnancy. Therapy will be started between 16 weeks, 0 days and 20 weeks, 6 days of gestation. Therapy will be continued until week 37 (through 36 weeks, 6 days) of gestation or delivery, whichever occurs first. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Preterm birth prophylaxis: Prescribed by or in consultation with a specialist in obstetrics and gynecology |
| Coverage Duration | Preterm birth prophylaxis: 21 weeks |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

MARINOL (S)

Products Affected

- Dronabinol

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Nausea and Vomiting Associated with Cancer Chemotherapy (CINV): Patient is receiving cancer chemotherapy. Trial and failure, contraindication, or intolerance to one 5HT-3 receptor antagonist (eg, Anzemet [dolasetron], Kytril [granisetron], or Zofran [ondansetron]). Trial and failure, contraindication, or intolerance to one of the following: Compazine (prochlorperazine), Decadron (dexamethasone), Haldol (haloperidol), Zyprexa (olanzapine). AIDS anorexia: Diagnosis of anorexia with weight loss in patients with AIDS. Patient is on antiretroviral therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | CINV: 6 months. AIDS anorexia: 3 months. |
| Other Criteria | Subject to Part B vs. Part D review. CINV: Approve for continuation of therapy for treatment covered under Part B when patient is receiving cancer chemotherapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

MAVENCLAD (S)

Products Affected

- Mavenclad

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Multiple Sclerosis (MS): Diagnosis of a relapsing form of MS (e.g., relapsing-remitting MS, secondary-progressive MS with relapses). One of the following: 1) Patient has not been previously treated with cladribine AND Failure after a trial of at least 4 weeks, contraindication, or intolerance to two of the following disease-modifying therapies for MS: Aubagio (teriflunomide), Avonex (interferon beta-1a), Betaseron (interferon beta-1b), Copaxone/Glatopa (glatiramer acetate), Extavia (interferon beta-1b), Gilenya (fingolimod), Lemtrada (alemtuzumab), Mayzent (siponimod), Ocrevus (ocrelizumab), Plegridy (peginterferon beta-1a), Rebif (interferon beta-1a), Tecfidera (dimethyl fumarate), Tysabri (natalizumab), Zinbryta (daclizumab), OR 2) Patient has previously received treatment with cladribine AND Patient has not already received the FDA-recommended lifetime limit of 2 treatment courses (or 4 treatment cycles total) of cladribine. Not used in combination with another disease-modifying therapy for MS. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | MS: 1 month |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

MAVYRET (S)

Products Affected

- Mavyret

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Criteria will be applied consistent with current AASLD/IDSA guideline. All patients: Diagnosis of chronic hepatitis C, patient is without decompensated liver disease (defined as Child-Pugh Class B or C), and patient is not receiving Mavyret in combination with another HCV direct acting antiviral agent [e.g., Harvoni, Zepatier]. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with one of the following: Hepatologist, Gastroenterologist, Infectious disease specialist, HIV specialist certified through the American Academy of HIV Medicine. |
| Coverage Duration | 8 to 16 weeks. Criteria will be applied consistent with current AASLD/IDSA guideline. |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

MAYZENT (S)

Products Affected

- Mayzent

- Mayzent Starter Pack

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Multiple Sclerosis (MS): Diagnosis of a relapsing form of MS (eg, relapsing-remitting MS, secondary-progressive MS with relapses). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

MEKINIST (S)

Products Affected

- Mekinist

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | <p>Melanoma: Diagnosis of unresectable or metastatic melanoma AND cancer is BRAF V600E or V600K mutant type as detected by a U.S. Food and Drug Administration (FDA)-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Adjuvant Treatment for Melanoma: Diagnosis of melanoma. Cancer is BRAF V600E or V600K mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Involvement of lymph nodes following complete resection. Used as adjunctive therapy. Medication is used in combination with Tafenlar (dabrafenib). Non-small Cell Lung Cancer (NSCLC): All of the following: diagnosis of metastatic non-small cell lung cancer AND cancer is BRAF V600E mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) AND medication is used in combination with Tafenlar (dabrafenib). Anaplastic Thyroid Cancer (ATC): Diagnosis of locally advanced or metastatic ATC. Cancer is BRAF V600E mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Cancer may not be treated with standard locoregional treatment options. Medication is used in combination with Tafenlar (dabrafenib).</p> |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist. |
| Coverage Duration | 12 months |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

| | |
|-----------------------|--|
| Other Criteria | Approve for continuation of prior therapy. |
|-----------------------|--|

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

MEKTOVI (S)

Products Affected

- Mektovi

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Melanoma: Diagnosis of unresectable melanoma or metastatic melanoma. Cancer is BRAF V600E or V600K mutant type (MT) as detected by a U.S. Food and Drug Administration (FDA)-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Used in combination with Braftovi (encorafenib). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

METADATE ER-RITALIN SR (S)

Products Affected

- Metadate Er TBCR 20MG
- Methylphenidate Hcl Sr
- Methylphenidate Hydrochloride Er TBCR 10MG, 20MG

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | One of the following: a) diagnosis of attention deficit hyperactivity disorder (ADHD), OR b) diagnosis of attention deficit disorder (ADD), OR c) diagnosis of narcolepsy as confirmed by a sleep study (unless the prescriber provides justification confirming that a sleep study would not be feasible). |
| Age Restrictions | PA applies to members 19 years of age or older |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

METHOTREXATE INJECTION (S)

Products Affected

- Rasuvo

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Rheumatoid Arthritis (RA) (initial): Diagnosis of severe, active RA. Polyarticular juvenile idiopathic arthritis (PJIA) (initial): Diagnosis of active PJIA. Psoriasis (initial): Diagnosis of severe psoriasis. All Indications (initial): Trial and failure or intolerance to oral methotrexate. |
| Age Restrictions | N/A |
| Prescriber Restrictions | RA, PJIA (initial): Prescribed by or in consultation with a rheumatologist. Psoriasis (initial): Prescribed by or in consultation with a dermatologist. |
| Coverage Duration | All Indications (Initial, reauth): 12 months |
| Other Criteria | All Indications (reauth): Documentation of positive clinical response to therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

METHYLIN CHEW (S)

Products Affected

- Methylphenidate Hydrochloride CHEW

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | One of the following: a) diagnosis of attention deficit hyperactivity disorder (ADHD), OR b) diagnosis of attention deficit disorder (ADD), OR c) diagnosis of narcolepsy as confirmed by a sleep study (unless the prescriber provides justification confirming that a sleep study would not be feasible). |
| Age Restrictions | PA applies to members 19 years of age or older |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

METHYLPHENIDATE (S)

Products Affected

- Methylphenidate Hydrochloride TABS

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | One of the following: a) diagnosis of attention deficit hyperactivity disorder (ADHD), OR b) diagnosis of attention deficit disorder (ADD), OR c) diagnosis of narcolepsy as confirmed by a sleep study (unless the prescriber provides justification confirming that a sleep study would not be feasible). |
| Age Restrictions | PA applies to members 19 years of age or older |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

METHYLPHENIDATE ER (S)

Products Affected

- Methylphenidate Hydrochloride CD
CPCR 10MG, 20MG, 50MG, 60MG
- Methylphenidate Hydrochloride Er
CP24
- Methylphenidate Hydrochloride Er
CPCR 20MG, 30MG, 40MG
- Methylphenidate Hydrochloride Er
TB24
- Methylphenidate Hydrochloride Er
TBCR 18MG, 27MG, 36MG, 54MG,
72MG
- Methylphenidate Hydrochloride Er (1a)
- Relexxii

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | One of the following: a) diagnosis of attention deficit hyperactivity disorder (ADHD), OR b) diagnosis of attention deficit disorder (ADD). |
| Age Restrictions | PA applies to members 19 years of age or older |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

METHYLTESTOSTERONE (S)

Products Affected

- Methitest

- Methyltestosterone CAPS

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Hypogonadism (HG) (Initial): Diagnosis (dx) of HG AND male patient at birth AND one of the following: 1) Two pre-treatment serum total testosterone (T) levels less than 300 ng/dL (10.4 nmol/L) or less than the reference range for the lab OR 2) Both of the following: a) Has a condition that may cause altered sex-hormone binding globulin (SHBG) (eg, thyroid disorder, HIV disease, liver disorder, diabetes, obesity), and b) one pre-treatment calculated free or bioavailable T level less than 5 ng/dL (0.17 nmol/L) or less than reference range for the lab, OR 3) History of bilateral orchiectomy, panhypopituitarism, or a genetic disorder known to cause HG (eg, congenital anorchia, Klinefelter's syndrome). Delayed puberty (DP): Dx of DP AND male patient at birth. Breast cancer (BC): Dx of inoperable BC AND used for palliative treatment AND female patient at birth. Gender Dysphoria (GD) (off-label): Dx of GD. Patient is a female-to-male transsexual. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | HG(init): (New to T tx:6 mo. Cont T tx:12 mo), (reauth): 12 mo. BC, GD: 12 mo. DP: 6 mo. |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|-----------------------|--|
| Other Criteria | HG (Reauth): 1) Follow-up total serum T level within or below the normal limits of the reporting lab, or 2) Follow-up total serum T level outside of upper limits of normal for the reporting lab and the dose is adjusted, OR 3) Has a condition that may cause altered SHBG (eg, thyroid disorder, HIV disease, liver disorder, diabetes, obesity), and one of the following: Follow-up calculated free or bioavailable T level within or below the normal limits of the reporting lab, or follow-up calculated free or bioavailable T level outside of upper limits of normal for the reporting lab and the dose is adjusted. |
|-----------------------|--|

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

MIRVASO (S)

Products Affected

- Mirvaso

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Rosacea (init): Diagnosis of rosacea. Patient has moderate to severe persistent (nontransient) facial erythema. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | Rosacea (init, reauth): 12 months |
| Other Criteria | Rosacea (reauth) Documentation of positive clinical response to Mirvaso therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

MOZOBIL (S)

Products Affected

- Mozobil

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Hematopoietic Stem Cell (HSC) Mobilization: Patient with non-Hodgkin's lymphoma (NHL) or multiple myeloma (MM) who will be undergoing autologous HSC transplantation. Used in combination with granulocyte-colony stimulating factor (G-CSF) [e.g., Neupogen (filgrastim), Zarxio (filgrastim)]. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist/oncologist |
| Coverage Duration | One course of therapy up to 4 days |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

MS INTERFERONS (NON-PREFERRED) (S)

Products Affected

- Avonex
- Avonex Pen
- Betaseron
- Extavia
- Plegridy
- Plegridy Starter Pack

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Multiple Sclerosis (MS): Diagnosis of a relapsing form of MS (eg, relapsing-remitting MS, secondary-progressive MS with relapses). One of the following: 1) Trial and failure, contraindication, or intolerance (TF/C/I) to Rebif, or 2) for continuation of prior therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

MS INTERFERONS (PREFERRED) (S)

Products Affected

- Rebif
- Rebif Rebidose
- Rebif Rebidose Titration Pack
- Rebif Titration Pack

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Multiple Sclerosis (MS): Diagnosis of a relapsing form of MS (eg, relapsing-remitting MS, secondary-progressive MS with relapses). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

MULPLETA (S)

Products Affected

- Mulpleta

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Thrombocytopenia Prior to Planned Procedure (TPPP): Diagnosis (dx) of thrombocytopenia. Baseline platelet count is less than 50,000/mcL. Patient has chronic liver disease and is scheduled to undergo a procedure. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 1 month |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

MYALEPT (S)

Products Affected

- Myalept

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Lipodystrophy (initial): Diagnosis of congenital or acquired generalized lipodystrophy AND one of the following: 1) Diabetes mellitus or insulin resistance despite optimized insulin therapy at maximum tolerated doses OR 2) Hypertriglyceridemia despite optimized therapy with at least two triglyceride-lowering agents from different classes (e.g., fibrates, statins). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Initial: Prescribed by or in consultation with an endocrinologist |
| Coverage Duration | Initial, reauth: 12 months |
| Other Criteria | Lipodystrophy (reauth): Patient has experienced an objective response to therapy, such as A) Sustained reduction in hemoglobin A1c (HbA1c) level from baseline OR B) Sustained reduction in triglyceride (TG) levels from baseline. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

MYLOTARG (S)

Products Affected

- Mylotarg

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Acute myeloid leukemia (AML): One of the following diagnoses: Newly diagnosed AML or relapsed/refractory (R/R) AML. Disease is CD33-positive. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist/oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

NAGLAZYME (S)

Products Affected

- Naglazyme

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Mucopolysaccharidosis (MPS VI): Diagnosis of MPS VI (Maroteaux-Lamy Syndrome). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | MPS VI: 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

NATPARA (S)

Products Affected

- Natpara

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Hypocalcemia (Initial): Diagnosis of hypocalcemia due to chronic hypoparathyroidism. NATPARA is not being used in the setting of acute post-surgical hypoparathyroidism. Patient does not have a known calcium-sensing receptor mutation. Patient has a documented parathyroid hormone concentration that is inappropriately low for the level of calcium, recorded on at least two occasions within the previous 12 months. Patient has normal thyroid-stimulating hormone concentrations if not on thyroid hormone replacement therapy (or if on therapy, the dose had to have been stable for greater than or equal to 3 months). Patient has normal magnesium and serum 25-hydroxyvitamin D concentrations. NATPARA will be used as an adjunct treatment. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Hypocalcemia (initial): Prescribed by or in consultation with an endocrinologist. |
| Coverage Duration | Initial: 6 months. Reauth: 12 months |
| Other Criteria | Hypocalcemia (Reauth): One of the following: A) Patient has achieved and maintained serum calcium levels in the ideal range (7.5 - 10.6 mg/dL), OR B) Patient has experienced a 50% or greater reduction in oral calcium intake, OR C) Patient has experienced a 50% or greater reduction in oral vitamin D intake. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

NERLYNX (S)

Products Affected

- Nerlynx

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Early Stage Breast cancer: Diagnosis (dx) of early stage breast cancer. Disease is human epidermal growth factor receptor 2 (HER2)-positive. Patient has received adjuvant trastuzumab-based therapy. Advanced or Metastatic Breast Cancer: Dx of advanced or metastatic breast cancer. Disease is human epidermal growth factor receptor 2 (HER2)-positive. Patient has received two or more prior anti-HER2 based regimens (e.g., trastuzumab + pertuzumab + docetaxel, ado-trastuzumab emtansine, etc.). Used in combination with capecitabine. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

NEULASTA (S)

Products Affected

- Neulasta

- Neulasta Onpro Kit

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | <p>Febrile neutropenia (FN) prophylaxis: Patient will be receiving prophylaxis for FN due to one of the following: 1) Patient is receiving National Cancer Institute’s Breast Intergroup, INT C9741 dose dense chemotherapy protocol for primary breast cancer, 2) patient is receiving a dose-dense chemotherapy regimen for which the incidence of FN is unknown, 3) patient is receiving chemotherapy regimen(s) associated with greater than 20% incidence of FN, 4) both of the following: a) patient is receiving chemotherapy regimen(s) associated with 10-20% incidence of FN, AND b) patient has one or more risk factors associated with chemotherapy-induced infection, FN, or neutropenia, OR 5) Both of the following: a) patient is receiving myelosuppressive anticancer drugs associated with neutropenia, AND b) patient has a history of FN or dose-limiting event during a previous course of chemotherapy (secondary prophylaxis). Acute radiation syndrome (ARS): Patient was/will be acutely exposed to myelosuppressive doses of radiation (hematopoietic subsyndrome of ARS). Treatment of FN: Patient has received or is receiving myelosuppressive anticancer drugs associated with neutropenia. Diagnosis of FN. Patient is at high risk for infection-associated complications.</p> |
| Age Restrictions | N/A |
| Prescriber Restrictions | All uses: Prescribed by or in consultation with a hematologist/oncologist |
| Coverage Duration | ARS: 1 mo. FN (prophylaxis, treatment): 3 mo or duration of tx. |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

NEXAVAR (S)

Products Affected

- Nexavar

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Renal cell carcinoma (RCC): Diagnosis of RCC. One of the following: Relapsed disease OR both medically/surgically unresectable tumor and dx of Stage IV disease. Hepatocellular carcinoma (HCC): Diagnosis of HCC. One of the following: patient has metastatic disease, or patient has extensive liver tumor burden, or patient is inoperable by performance status or comorbidity (local disease or local disease with minimal extrahepatic disease only), or disease is unresectable. Differentiated thyroid carcinoma (DTC): Diagnosis of DTC (ie, follicular carcinoma, Hurthle cell carcinoma, or papillary carcinoma). One of the following: locally recurrent disease, metastatic disease, or unresectable disease. One of the following: patient has symptomatic disease or patient has progressive disease. Disease is refractory to radioactive iodine (RAI) treatment. Medullary thyroid carcinoma (MTC): Diagnosis of MTC. One of the following: 1) Disease is progressive or 2) Disease is symptomatic with distant metastases. Trial and failure, contraindication, or intolerance to Caprelsa (vandetanib) or Cometriq (cabozantinib). |
| Age Restrictions | N/A |
| Prescriber Restrictions | DTC, MTC: Prescribed by or in consultation with an oncologist. RCC: Prescribed by or in consultation with an oncologist or nephrologist. HCC: Prescribed by or in consultation with an oncologist, hepatologist, or gastroenterologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

NEXLETOL (S)

Products Affected

- Nexletol

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Initial: One of the following diagnoses: A) Heterozygous familial hypercholesterolemia (HeFH) as confirmed by one of the following: (1) Both of the following: a) Untreated/pre-treatment LDL-C greater than 190 mg/dL (greater than 155 mg/dL if less than 16 years of age), AND b) One of the following: i) Family history (hx) of myocardial infarction in 1st-degree relative less than 60 years of age, ii) Family hx of myocardial infarction in 2nd-degree relative less than 50 years of age, iii) Family hx of LDL-C greater than 190 mg/dL in 1st- or 2nd-degree relative, iv) Family hx of FH in 1st- or 2nd-degree relative, or v) Family hx of tendinous xanthomas and/or arcus cornealis in 1st- or 2nd-degree relative, or (2) Both of the following: a) Untreated/pre-treatment LDL-C greater than 190 mg/dL (greater than 155 mg/dL if less than 16 years of age), AND b) One of the following: i) Functional mutation in the LDL receptor, ApoB, or PCSK9 gene, ii) Tendinous xanthomata, or iii) Arcus cornealis before age 45 OR B) Atherosclerotic cardiovascular disease (ASCVD) as confirmed by acute coronary syndromes, hx of myocardial infarction, stable or unstable angina, coronary or other arterial revascularization (eg, percutaneous coronary intervention [PCI] or coronary artery bypass graft [CABG] surgery), stroke, transient ischemic attack, peripheral arterial disease presumed to be of atherosclerotic origin, or clinically significant coronary heart disease diagnosed by invasive or noninvasive testing (e.g., coronary angiography, stress test using treadmill, stress echocardiography or nuclear imaging). One of the following LDL-C values while on max tolerated statin tx within the last 120 days: (1) LDL-C greater than or equal to 70 mg/dL with ASCVD or (2) LDL-C greater than or equal to 100 mg/dL without ASCVD. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|---------------------------------|---|
| <p>Coverage Duration</p> | <p>Initial: 6 months. Reauth: 12 months</p> |
| <p>Other Criteria</p> | <p>Initial, cont: One of the following: (1) Pt has been receiving at least 12 weeks of one high-intensity statin (HIS) therapy (tx) [ie, atorvastatin 40-80 mg, rosuvastatin 20-40 mg] and will continue to receive a HIS at max tolerated dose, OR (2) Both of the following: a) Pt is unable to tolerate HIS as evidenced by one of the following intolerable and persistent (ie, more than 2 wks) symptoms: myalgia (muscle symptoms w/o CK elevations) or myositis (muscle symptoms w/ CK elevations less than 10 times ULN) AND b) Pt has been receiving at least 12 weeks of one moderate-intensity statin (MIS) [i.e., atorvastatin 10-20 mg, rosuvastatin 5-10 mg, simvastatin 20-40 mg, pravastatin 40-80 mg, lovastatin 40 mg, Lescol XL (fluvastatin XL) 80 mg, fluvastatin 40 mg twice daily, or Livalo (pitavastatin) 2-4 mg] or one low-intensity statin (LIS) [i.e., simvastatin 10 mg, pravastatin 10-20 mg, lovastatin 20 mg, fluvastatin 20-40 mg, Livalo (pitavastatin) 1 mg] tx and will continue to receive a MIS or LIS at max tolerated dose, OR (3) Pt is unable to tolerate low-, moderate-, or high-intensity statins as evidenced by one of the following intolerable and persistent (ie, more than 2 weeks) symptoms: myalgia (muscle symptoms without CK elevations) or myositis (muscle symptoms with CK elevations less than 10 times ULN), OR (4) Pt has a labeled contraindication to all statins, OR (5) Pt has experienced rhabdomyolysis or muscle symptoms w/ statin tx with CK elevations greater than 10 times ULN. AND Pt has been receiving at least 12 weeks of generic ezetimibe tx as adjunct to max tolerated statin tx or pt has a history of contraindication or intolerance to ezetimibe. Reauth: Documentation of positive clinical response to therapy (eg reduction in LDL-C levels). Pt continues to receive other lipid-lowering tx (eg statins, ezetimibe) at the max tolerated dose or pt has a documented inability to take other lipid-lowering therapy (eg statins, ezetimibe).</p> |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

NINLARO (S)

Products Affected

- Ninlaro

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Multiple myeloma: Diagnosis of multiple myeloma. Used in combination with Revlimid (lenalidomide) and dexamethasone. Patient has received at least one prior therapy for multiple myeloma [eg, Revlimid (lenalidomide), Thalomid (thalidomide), Velcade (bortezomib)]. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist/oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

NON-PREFERRED TIRF (S)

Products Affected

- Abstral
- Fentanyl Citrate TABS
- Lazanda

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | For the management of breakthrough cancer pain. Patient is currently taking a long-acting opioid around the clock for cancer pain. Patient must have at least a one week history of ONE of the following medications to demonstrate tolerance to opioids: Morphine sulfate at doses of greater than or equal to 60 mg/day, Fentanyl transdermal patch at doses greater than or equal to 25 µg/hr, Oxycodone at a dose of greater than or equal to 30 mg/day, Oral hydromorphone at a dose of greater than or equal to 8 mg/day, Oral oxymorphone at a dose of greater than or equal to 25 mg/day, or an alternative opioid at an equianalgesic dose (e.g., oral methadone greater than or equal to 20 mg/day). Trial and failure or intolerance to generic fentanyl lozenge. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with one of the following: Pain specialist, Oncologist, Hematologist, Hospice care specialist, or Palliative care specialist. |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

NORTHERA (S)

Products Affected

- Northera

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Neurogenic orthostatic hypotension (NOH) (init): Diagnosis of symptomatic NOH. NOH is caused by one of the following conditions: primary autonomic failure (eg, Parkinson's disease, multiple system atrophy, pure autonomic failure), dopamine beta-hydroxylase deficiency, non-diabetic autonomic neuropathy. Trial and failure, contraindication, or intolerance to one of the following agents: fludrocortisone acetate, midodrine. |
| Age Restrictions | N/A |
| Prescriber Restrictions | NOH (init): Prescribed by or in consultation with a cardiologist, neurologist, or nephrologist |
| Coverage Duration | NOH (init): 1 month (reauth): 12 months |
| Other Criteria | NOH (reauth): Documentation of positive clinical response to therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020
NOVANTRONE (S)

Products Affected

- Mitoxantrone Hcl INJ 2MG/ML

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Multiple Sclerosis (MS): Diagnosis (dx) of one of the following: secondary progressive MS: gradually worsening disability with or without superimposed relapses, progressive relapsing MS: progression of disability from the onset with superimposed relapses, or worsening relapsing-remitting MS: neurological status remains significantly abnormal in between MS relapses. Trial and failure, contraindication, or intolerance to one of the following disease-modifying therapies for MS: Avonex (interferon beta-1a), Aubagio (teriflunomide), Betaseron (interferon beta-1b), Copaxone/Glatopa (glatiramer acetate), Extavia (interferon beta-1b), Gilenya (fingolimod), Lemtrada (alemtuzumab), Mavenclad (cladribine), Mayzent (siponimod), Ocrevus (ocrelizumab), Plegridy (peginterferon beta-1a), Rebif (interferon beta-1a), Tecfidera (dimethyl fumarate), Tysabri (natalizumab), Zinbryta (daclizumab). Left ventricular ejection fraction (LVEF) greater than or equal to 50%. Neutrophil count greater than or equal to 1500 cell/mm ³ . Lifetime cumulative dose less than 140 mg/m ² . Prostate Cancer (PC): Dx of advanced hormone-refractory (castration-resistant) PC. Used in combination with corticosteroids (eg, prednisone, methylprednisolone). LVEF greater than or equal to 50%. Neutrophil count greater than or equal to 1500 cell/mm ³ . Acute Non-Lymphocytic Leukemia (ANLL): Dx of ANLL (eg, myelogenous, promyelocytic, monocytic, and erythroid). Used in combination with other medications used for the treatment of ANLL. LVEF greater than or equal to 50%. |
| Age Restrictions | N/A |
| Prescriber Restrictions | PC: Prescribed by or in consultation with an oncologist. ANLL: Prescribed by or in consultation with a hematologist/oncologist. |
| Coverage Duration | All Uses: 6 months |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

| | |
|-----------------------|--|
| Other Criteria | Approve for continuation of prior therapy. |
|-----------------------|--|

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

NPLATE (S)

Products Affected

- Nplate

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Immune (idiopathic) thrombocytopenic purpura (ITP): All of the following: A) Diagnosis of one of the following: a) chronic ITP or b) relapsed/refractory ITP AND B) Baseline platelet count is less than 30,000/mcL AND C) Patient's degree of thrombocytopenia and clinical condition increase the risk of bleeding AND D) One of the following: 1) Patient had an insufficient response, intolerance, or contraindication to corticosteroids or immune globulin or 2) Patient had an inadequate response or contraindication to splenectomy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | ITP (initial): Prescribed by or in consultation with a hematologist/oncologist. |
| Coverage Duration | ITP (initial, reauth): 12 months |
| Other Criteria | ITP (reauth): After at least 4 weeks of therapy at the maximum weekly dose (10 mcg/kg) the platelet count increased to a sufficient level to avoid clinically important bleeding. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

NUBEQA (S)

Products Affected

- Nubeqa

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Non-metastatic castration-resistant or castration-recurrent prostate cancer (nmCRPC): Diagnosis of non-metastatic castration-resistant (chemical or surgical) or castration-recurrent prostate cancer. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or urologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

NUCALA (S)

Products Affected

- Nucala

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | <p>Asthma (init): Diagnosis of severe asthma. Asthma is an eosinophilic phenotype as defined by one of the following: baseline (pre-treatment) peripheral blood eosinophil level is greater than or equal to 150 cells/microliter or peripheral blood eosinophil levels were greater than or equal to 300 cells/microliter within the past 12 months. Patient has had at least one or more asthma exacerbations requiring systemic corticosteroids within the past 12 months or Patient has had any prior intubation for an asthma exacerbation or Patient has had a prior asthma-related hospitalization within the past 12 months. Patient is currently being treated with one of the following unless there is a contraindication or intolerance to these medications: a) Both of the following: i) High-dose inhaled corticosteroid (ICS) [e.g., greater than 500 mcg fluticasone propionate equivalent/day] and ii) additional asthma controller medication [e.g., leukotriene receptor antagonist, long-acting beta-2 agonist (LABA), theophylline], OR b) One maximally-dosed combination ICS/LABA product [e.g., Advair (fluticasone propionate/salmeterol), Dulera (mometasone/formoterol), Symbicort (budesonide/formoterol)].</p> <p>Eosinophilic Granulomatosis with Polyangiitis (EGPA) (init): Diagnosis of EGPA. Patient's disease has relapsed or is refractory to standard of care therapy (i.e., corticosteroid treatment with or without immunosuppressive therapy). Patient is currently receiving corticosteroid therapy (e.g., prednisolone, prednisone).</p> |
| Age Restrictions | Asthma (init): Age greater than or equal to 6 years |
| Prescriber Restrictions | Asthma (init, reauth): Prescribed by or in consultation with a pulmonologist or allergist/immunologist. EGPA (init): Prescribed by or in consultation with a pulmonologist, rheumatologist or allergist/immunologist. |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|--------------------------|---|
| Coverage Duration | Asthma (init): 6 mo, Asthma (reauth): 12 months. EGPA (init, reauth): 12 months |
| Other Criteria | Asthma (reauth): Documentation of positive clinical response to therapy (eg, reduction in exacerbations, improvement in forced expiratory volume in 1 second (FEV1), decreased use of rescue medications). Patient is currently being treated with one of the following unless there is a contraindication or intolerance to these medications: a) Both of the following: i) inhaled corticosteroid (ICS) and ii) additional asthma controller medication [e.g., leukotriene receptor antagonist, long-acting beta-2 agonist (LABA), theophylline], OR b) A combination ICS/LABA product [e.g., Advair (fluticasone propionate/salmeterol), Dulera (mometasone/formoterol), Symbicort (budesonide/formoterol)]. EGPA (reauth): Documentation of positive clinical response to therapy (e.g., increase in remission time). |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

NUEDEXTA (S)

Products Affected

- Nuedexta

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Pseudobulbar affect (PBA) (initial): Diagnosis of PBA. (Reauthorization): Documentation of clinical benefit from ongoing therapy with Nuedexta. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | PBA (initial/reauth): 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

NULOJIX (S)

Products Affected

- Nulojix

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Kidney transplant: The medication is being used for prevention of kidney transplant organ rejection AND The patient is immune to the Epstein-Barr virus (i.e. EBV seropositive) AND The patient is prescribed concurrent therapy with mycophenolate and corticosteroids. |
| Age Restrictions | Kidney transplant: 18 years of age or older |
| Prescriber Restrictions | Kidney transplant: Prescriber is experienced in immunosuppressive therapy and management of transplant patients |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

NUPLAZID (S)

Products Affected

- Nuplazid

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Parkinson's disease psychosis: Diagnosis of Parkinson's disease. Patient has at least one of the following: hallucinations or delusions. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

NURTEC (S)

Products Affected

- Nurtec

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Initial: Diagnosis of migraine with or without aura. Will be used for the acute treatment of migraine. Will not be used for preventive treatment of migraine. Patient has fewer than 15 headache days per month. Trial and failure or intolerance to one triptan (e.g., eletriptan, rizatriptan, sumatriptan) or a contraindication to all triptans. If patient has 4 or more headache days per month, patient must meet one of the following: a) currently being treated with Elavil (amitriptyline) or Effexor (venlafaxine) unless there is a contraindication or intolerance to these medications, OR b) currently being treated with Depakote/Depakote ER (divalproex sodium) or Topamax (topiramate) unless there is a contraindication or intolerance to these medications, OR c) currently being treated with a beta blocker (i.e., atenolol, propranolol, nadolol, timolol, or metoprolol) unless there is a contraindication or intolerance to these medications. Medication will not be used in combination with another CGRP inhibitor. |
| Age Restrictions | Initial: 18 years of age or older. |
| Prescriber Restrictions | Initial, Reauth: Prescribed by or in consultation with a neurologist or pain specialist. |
| Coverage Duration | Initial: 3 months. Reauth: 12 months. |
| Other Criteria | Reauth: Patient has experienced a positive response to therapy (e.g., reduction in pain, photophobia, phonophobia, nausea). Will not be used for preventive treatment of migraine. Medication will not be used in combination with another CGRP inhibitor. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

NUVIGIL (S)

Products Affected

- Armodafinil

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Obstructive sleep apnea/hypopnea syndrome (OSAHS) (Initial): Diagnosis (dx) of OSAHS defined by one of the following: a) 15 or more obstructive respiratory events per hour of sleep confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible), or b) both of the following: 5 or more obstructive respiratory events per hour of sleep confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible), AND 1 of the following symptoms: unintentional sleep episodes during wakefulness, daytime sleepiness, unrefreshing sleep, fatigue, insomnia, waking up breath holding/gasping/choking, loud snoring, or breathing interruptions during sleep. Shift-work sleep disorder (SWSD) (Initial):Dx of SWSD confirmed by one of the following: 1) symptoms of excessive sleepiness or insomnia for at least 3 months, which is associated with a work period (usually night work) that occurs during the normal sleep period, OR 2) A sleep study demonstrating loss of a normal sleep-wake pattern (ie, disturbed chronobiologic rhythmicity). No other medical condition or medication accounts for the symptoms. Narcolepsy (initial): Dx of narcolepsy as confirmed by sleep study (unless prescriber provides justification confirming that a sleep study is not feasible). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | OSAHS, SWSD: Initial, Reauth: 6 mo. Narcolepsy: Initial, Reauth: 12 mo |
| Other Criteria | OSAHS, Narcolepsy (Reauth): Documentation of positive clinical response to armodafinil therapy. SWSD (Reauth): Documentation of positive clinical response to armodafinil therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

Ocaliva (S)

Products Affected

- Ocaliva

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Primary Biliary Cholangitis (PBC) (initial): Diagnosis of PBC (aka primary biliary cirrhosis). One of the following: a) patient has failed to achieve an alkaline phosphatase (ALP) level of less than 1.67 times the upper limit of normal (ULN) after treatment with ursodeoxycholic acid (UDCA) (e.g., Urso, Urso Forte, ursodiol) AND used in combination with UDCA, OR b) contraindication or intolerance to UDCA. Patients with moderate to severe hepatic impairment (Child-Pugh class B or C) will be subject to a quantity limit of 5 mg or 10 mg twice weekly (MDD = 0.34). |
| Age Restrictions | N/A |
| Prescriber Restrictions | PBC (initial): Prescribed by or in consultation with a hepatologist or gastroenterologist. |
| Coverage Duration | PBC (initial): 6 months, (reauth): 12 months |
| Other Criteria | PBC (reauthorization): Submission of medical records (eg, laboratory values) documenting a reduction in ALP level from pre-treatment baseline (ie, prior Ocaliva therapy) while on Ocaliva therapy. Patients with moderate to severe hepatic impairment (Child-Pugh class B or C) will be subject to a quantity limit of 5 mg or 10 mg twice weekly (MDD = 0.34). |

OCREVUS (S)

Products Affected

- Ocrevus

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Relapsing forms of multiple sclerosis (initial): Diagnosis of a relapsing form of multiple sclerosis (MS) (e.g., relapsing-remitting MS, secondary-progressive MS with relapses). One of the following: a) Failure after a trial of at least 4 weeks, contraindication, or intolerance to one of the following: Aubagio (teriflunomide), Avonex (interferon beta-1a), Betaseron (interferon beta-1b), Copaxone/Glatopa (glatiramer acetate), Extavia (interferon beta-1b), Gilenya (fingolimod), Lemtrada (alemtuzumab), Mavenclad (cladribine), Mayzent (siponimod), Plegridy (peginterferon beta-1a), Rebif (interferon beta-1a), Tecfidera (dimethyl fumarate), Tysabri (natalizumab), Zinbryta (daclizumab) OR b) Patient is not a candidate for any of the drugs listed as prerequisites due to the severity of their multiple sclerosis, OR c) For continuation of prior Ocrevus therapy. Primary progressive MS (initial): Diagnosis of primary progressive multiple sclerosis (PPMS). All indications (initial, reauth): Not used in combination with another disease-modifying therapy for MS. Not used in combination with another B-cell targeted therapy (e.g., rituximab [Rituxan], belimumab [Benlysta], ofatumumab [Arzerra]). Not used in combination with another lymphocyte trafficking blocker (e.g., alemtuzumab [Lemtrada], mitoxantrone). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | All uses (initial, reauth): 12 months |
| Other Criteria | All indications (reauth): Documentation of positive clinical response to Ocrevus therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ODOMZO (S)

Products Affected

- Odomzo

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Basal cell carcinoma: Diagnosis of locally advanced basal cell carcinoma AND One of the following: 1) Cancer has recurred following surgery or radiation therapy or 2) Patient is not a candidate for surgery or radiation therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or dermatologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

OFEV (S)

Products Affected

- Ofev

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | <p>Idiopathic pulmonary fibrosis (IPF) (initial): Both of the following: 1) diagnosis of IPF as documented by all of the following: a) exclusion of other known causes of interstitial lung disease (ILD) (eg, domestic and occupational environmental exposures, connective tissue disease, drug toxicity) AND b) one of the following: i) in patients not subjected to surgical lung biopsy, the presence of a usual interstitial pneumonia (UIP) pattern on high-resolution computed tomography (HRCT) revealing IPF or probable IPF, OR ii) in patients subjected to a lung biopsy, both HRCT and surgical lung biopsy pattern revealing IPF or probable IPF, AND 2) not used in combination with Esbriet (pirfenidone). Systemic sclerosis-associated interstitial lung disease (SSc-ILD) (initial): Both of the following: 1) Diagnosis of SSc-ILD as documented by all of the following: a) exclusion of other known causes of ILD (eg, domestic and occupational environmental exposures, connective tissue disease, drug toxicity) AND b) One of the following: i) In patients not subjected to surgical lung biopsy, the presence of idiopathic interstitial pneumonia (eg, fibrotic nonspecific interstitial pneumonia [NSIP], usual interstitial pneumonia [UIP] and centrilobular fibrosis) pattern on HRCT revealing SSc-ILD or probable SSc-ILD, OR ii) in patients subjected to a lung biopsy, both HRCT and surgical lung biopsy pattern revealing SSc-ILD or probable SSc-ILD, AND 2) not used in combination with Esbriet (pirfenidone). Chronic Fibrosing Interstitial Lung Diseases (ILDs) with a Progressive Phenotype (initial): 1) diagnosis of chronic fibrosing interstitial lung disease, AND 2) patient has a high-resolution computed tomography (HRCT) showing at least 10% of lung volume with fibrotic features, AND 3) disease has a progressive phenotype as observed by one of the following: decline of forced vital capacity (FVC), worsening of respiratory symptoms, or increased extent of fibrosis seen on imaging.</p> |
| Age Restrictions | N/A |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|--------------------------------|--|
| Prescriber Restrictions | IPF, SSc-ILD, Chronic Fibrosing ILDs with a Progressive Phenotype (initial): Prescribed by or in consultation with a pulmonologist |
| Coverage Duration | Initial, reauth: 12 months |
| Other Criteria | IPF, SSc-ILD, Chronic Fibrosing ILDs with a Progressive Phenotype (reauth): Documentation of positive clinical response to Ofev therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

OLUMIANT (S)

Products Affected

- Olumiant

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Rheumatoid arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. One of the following: Trial and failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR patient has a documented needle-phobia to the degree that the patient has previously refused any injectable therapy or medical procedure (refer to DSM-IV-TR F40.2 for specific phobia diagnostic criteria), OR for continuation of prior Olumiant therapy. Patient is not receiving Olumiant in combination with a potent immunosuppressant (e.g., azathioprine, cyclosporine). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Initial: Prescribed by or in consultation with a rheumatologist. |
| Coverage Duration | Initial, reauth: 12 months |
| Other Criteria | Reauth: Documentation of positive clinical response to Olumiant therapy. Patient is not receiving Olumiant in combination with a potent immunosuppressant (e.g., azathioprine, cyclosporine). |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ONMEL (S)

Products Affected

- Onmel

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | All of the following: 1) Diagnosis of onychomycosis of the toenail as confirmed by one of the following: a) positive potassium hydroxide (KOH) preparation, OR b) fungal culture, OR c) nail biopsy, AND 2) patient has had a trial and failure, contraindication, or intolerance to oral terbinafine. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 3 months |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

ONPATTRO (S)

Products Affected

- Onpattro

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Hereditary transthyretin-mediated amyloidosis (hATTR amyloidosis) (initial): Diagnosis of hATTR amyloidosis with polyneuropathy. Patient has a transthyretin (TTR) mutation (e.g., V30M). Patient has a baseline polyneuropathy disability (PND) score less than or equal to IIIb, a baseline familial amyloidotic polyneuropathy (FAP) stage of 1 or 2, or a baseline neuropathy impairment score (NIS) between 10 and 130. Presence of clinical signs and symptoms of the disease (e.g., peripheral/autonomic neuropathy). |
| Age Restrictions | N/A |
| Prescriber Restrictions | hATTR amyloidosis (initial): Prescribed by or in consultation with a neurologist |
| Coverage Duration | hATTR amyloidosis (initial, reauth): 12 months |
| Other Criteria | Subject to Part B vs D review. hATTR amyloidosis (reauth): Patient has demonstrated a benefit from therapy (e.g., improved neurologic impairment, slowing of disease progression, quality of life assessment). Patient continues to have a PND score less than or equal to IIIb, a FAP stage of 1 or 2, or a NIS between 10 and 130. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

OPDIVO (S)

Products Affected

- Opdivo

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | N/A |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

OPSUMIT (S)

Products Affected

- Opsumit

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH. |
| Age Restrictions | N/A |
| Prescriber Restrictions | PAH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist. |
| Coverage Duration | PAH: Initial: 6 months. Reauth: 12 months. |
| Other Criteria | PAH (Reauth): Documentation of positive clinical response to therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ORENCIA IV (S)

Products Affected

- Orencia INJ 250MG

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Rheumatoid Arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. Juvenile Idiopathic Arthritis (JIA) (Initial): Diagnosis of moderately to severely active polyarticular JIA. Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. All indications (Initial): One of the following: Trial and failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR for continuation of prior Orencia therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | RA, JIA (initial): Prescribed by or in consultation with a rheumatologist. PsA (initial): Prescribed by or in consultation with a dermatologist or rheumatologist. |
| Coverage Duration | All uses (initial, reauth): 12 months |
| Other Criteria | All indications (Reauth): Documentation of positive clinical response to Orencia therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ORENCIA SC (S)

Products Affected

- Orencia INJ 125MG/ML, 50MG/0.4ML, 87.5MG/0.7ML
- Orencia Clickject

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Rheumatoid Arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. Juvenile Idiopathic Arthritis (JIA) (Initial): Diagnosis of moderately to severely active polyarticular JIA. Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. All indications (Initial): One of the following: Trial and failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR for continuation of prior Orencia therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | RA, JIA (initial): Prescribed by or in consultation with a rheumatologist. PsA (initial): Prescribed by or in consultation with a dermatologist or rheumatologist. |
| Coverage Duration | All uses (initial, reauth): 12 months |
| Other Criteria | All indications (Reauth): Documentation of positive clinical response to Orencia therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ORENITRAM (S)

Products Affected

- Orenitram

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH. |
| Age Restrictions | N/A |
| Prescriber Restrictions | PAH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist. |
| Coverage Duration | PAH: Initial: 6 months. Reauth: 12 months. |
| Other Criteria | PAH (Reauth): Documentation of positive clinical response to therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

ORIAHNN (S)

Products Affected

- Oriahnn

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Initial: Diagnosis of heavy menstrual bleeding associated with uterine leiomyomas (fibroids). Patient is premenopausal. One of the following: 1) History of inadequate control of bleeding following a trial of at least 3 months, or history of intolerance or contraindication to one of the following: combination (estrogen/progesterone) oral contraceptive, progestins, or tranexamic acid or 2) Patient has had a previous interventional therapy to reduce bleeding. Treatment duration of therapy has not exceeded a total of 24 months. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | Reauth: Patient has improvement in bleeding associated with uterine leiomyomas (fibroids) (e.g., significant/sustained reduction in menstrual blood loss per cycle, improved quality of life, etc.). Treatment duration of therapy has not exceeded a total of 24 months. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

ORILISSA (S)

Products Affected

- Orilissa

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Endometriosis (EM) (initial - 150 mg): Diagnosis of moderate to severe pain associated with EM. One of the following: 1) History of inadequate pain control response following a trial of at least 3 months, or history of intolerance or contraindication to one of the following: danazol, combination (estrogen/progesterone) oral contraceptive, or progestins, or 2) Patient has had surgical ablation to prevent recurrence. EM (200 mg): Diagnosis of moderate to severe pain associated with EM. One of the following: 1) History of inadequate pain control response following a trial of at least 3 months, or history of intolerance or contraindication to one of the following: danazol, combination (estrogen/progesterone) oral contraceptive, or progestins, or 2) Patient has had surgical ablation to prevent recurrence. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | EM (init, reauth-150mg): 6 mo. EM (200mg): 6 mo. |
| Other Criteria | EM (reauthorization - 150 mg): Patient has improvement in pain associated with endometriosis (e.g., improvement in dysmenorrhea and nonmenstrual pelvic pain). Treatment duration of Orilissa has not exceeded a total of 24 months. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ORKAMBI (S)

Products Affected

- Orkambi TABS

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Cystic Fibrosis (CF) (Initial): Diagnosis of CF. Patient is homozygous for the F508del mutation in the CF transmembrane conductance regulator (CFTR) gene as detected by a U.S. Food and Drug Administration (FDA)-cleared cystic fibrosis mutation test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). |
| Age Restrictions | CF (Initial): Patient is 6 years of age or older |
| Prescriber Restrictions | CF (initial): Prescribed by or in consultation with a specialist affiliated with a CF care center or pulmonologist |
| Coverage Duration | CF (initial, reauth): 12 months |
| Other Criteria | CF (Reauth): Patient is benefiting from treatment (i.e. improvement in lung function [forced expiratory volume in one second (FEV1)], decreased number of pulmonary exacerbations). |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

ORKAMBI GRANULES (S)

Products Affected

- Orkambi PACK

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Cystic Fibrosis (CF) (Initial): Diagnosis of CF. Patient is homozygous for the F508del mutation in the CF transmembrane conductance regulator (CFTR) gene as detected by a U.S. Food and Drug Administration (FDA)-cleared cystic fibrosis mutation test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). One of the following: A) Patient is 2 through 5 years of age, OR B) Both of the following: Patient is 6 years of age or greater AND Patient is unable to swallow oral tablets. |
| Age Restrictions | N/A |
| Prescriber Restrictions | CF (Initial): Prescribed by or in consultation with a specialist affiliated with a CF care center or pulmonologist |
| Coverage Duration | CF (initial, reauth): 12 months |
| Other Criteria | CF (Reauth): Patient is benefiting from treatment (i.e., improvement in lung function [forced expiratory volume in one second (FEV1)], decreased number of pulmonary exacerbations). One of the following: A) Patient is 2 through 5 years of age, OR B) Both of the following: Patient is 6 years of age or greater AND Patient is unable to swallow oral tablets. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

OSPHERA (S)

Products Affected

- Ospheña

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Dyspareunia (initial): Diagnosis of moderate to severe dyspareunia due to vulvar and vaginal atrophy associated with menopause. Vaginal dryness (initial): Diagnosis of moderate to severe vaginal dryness due to vulvar and vaginal atrophy associated with menopause. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | All uses (Initial, reauth): 12 months |
| Other Criteria | Dyspareunia, Vaginal dryness (reauth): Documentation of positive clinical response to therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

OTEZLA (S)

Products Affected

- Otezla

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Psoriatic arthritis (PsA, initial): Diagnosis of active PsA. Plaque psoriasis (Initial): Diagnosis of moderate to severe plaque psoriasis. PsA, plaque psoriasis (initial): Trial and failure, contraindication, or intolerance to both Humira and Enbrel, or attestation demonstrating a trial may be inappropriate, OR for continuation of prior Otezla therapy. Oral ulcers associated with Behcet's Disease (Initial): Diagnosis of Behcet's Disease. Patient has active oral ulcers. |
| Age Restrictions | N/A |
| Prescriber Restrictions | PsA (init): Prescribed by or in consultation with a dermatologist or rheumatologist. Plaque psoriasis (init): Prescribed by or in consultation with a dermatologist. |
| Coverage Duration | Initial, Reauth: 12 months |
| Other Criteria | Reauth (PsA, plaque psoriasis): Documentation of positive clinical response to Otezla therapy. Reauth (oral ulcers associated with Behcet's Disease): Documentation of positive clinical response to Otezla therapy (eg, reduction in pain from oral ulcers or reduction in number of oral ulcers). |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

OXANDRIN (S)

Products Affected

- Oxandrolone TABS

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Promote weight gain (initial): Used as adjunctive therapy to promote weight gain AND Diagnosis of one of the following: Extensive surgery, Chronic infections, Severe trauma, Failure to gain or maintain at least 90% of ideal body weight without definite pathophysiologic reasons AND a nutritional consult was performed. Counterbalance protein catabolism (initial): Used to counterbalance protein catabolism associated with chronic corticosteroid administration. Bone pain: Diagnosis of bone pain associated with osteoporosis. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | bone pain: 1 month. Others (initial, reauth): 3 months |
| Other Criteria | All diagnoses except bone pain (reauth): Documentation of a positive clinical response to therapy as evidenced by an improvement in weight gain or increase in lean body mass. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

OXBRYTA (S)

Products Affected

- Oxbryta

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Initial: Diagnosis of Sickle Cell Disease. Documentation of hemoglobin level that does not exceed 10.5 g/dL prior to therapy initiation. Trial and failure, contraindication, or intolerance to hydroxyurea. |
| Age Restrictions | Initial: Patient is 12 years of age or older. |
| Prescriber Restrictions | Initial: Prescribed by or in consultation with one of the following: 1) Hematologist/Oncologist or 2) Specialist w/ expertise in the diagnosis and management of sickle cell disease. |
| Coverage Duration | Initial, Reauth: 12 months. |
| Other Criteria | Reauth: Documentation of positive clinical response to Oxbryta therapy (e.g., an increase in hemoglobin level of 1 g/dL or greater from baseline, decreased annualized incidence rate of VOCs). Documentation of hemoglobin level that does not exceed 10.5 g/dL. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

OXERVATE (S)

Products Affected

- Oxervate

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Neurotrophic keratitis (NK): Diagnosis of NK. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an ophthalmologist or optometrist. |
| Coverage Duration | 8 weeks. |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

PEGASYS (S)

Products Affected

- Pegasys

- Pegasys Proclick

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Chronic hepatitis B: Diagnosis of chronic hepatitis B infection, and patient is without decompensated liver disease. Chronic Hepatitis C: Criteria will be applied consistent with current AASLD-IDSA guidance. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | HepB: 48 wks. HepC: Initial: 28 wks. Reauth: 20 wks. |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

PEG-INTRON (S)

Products Affected

- Pegintron INJ 50MCG/0.5ML
- Peg-intron Redipen INJ 120MCG/0.5ML

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Chronic Hepatitis C:Criteria will be applied consistent with current AASLD-IDSA guidance. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | HepC: Initial: 28 wks. Reauth: 20 wks. |
| Other Criteria | HepC (reauth): patient has an undetectable HCV RNA at week 24, additional treatment weeks of peginterferon are required to complete treatment regimen, and patient has not exceeded 48 wks of therapy with peginterferon. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

PEMAZYRE (S)

Products Affected

- Pemazyre

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Diagnosis of cholangiocarcinoma. Disease is one of the following: unresectable locally advanced or metastatic. Disease has presence of a fibroblast growth factor receptor 2 (FGFR2) fusion or other rearrangement. Patient has been previously treated. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hepatologist, gastroenterologist, or oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

PENNSAID (S)

Products Affected

- Diclofenac Sodium TRANSDERMAL SOLN 1.5%
- Klofensaid II
- Pennsaid SOLN 2%

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Osteoarthritis of the knees (initial): Diagnosis of osteoarthritis of the knees. Patient meets one of the following: 1) Treatment failure with at least two prescription strength oral non-steroidal anti-inflammatory drugs (NSAIDs) OR 2) Documented swallowing disorder OR 3) History of peptic ulcer disease/gastrointestinal bleed OR 4) Patient is older than 65 years of age with one additional risk factor for gastrointestinal adverse events (e.g. use of anticoagulants, chronic corticosteroids). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | Initial, reauth: 12 months |
| Other Criteria | Osteoarthritis of the knees (reauth): Documentation of positive clinical response to therapy (e.g., improvement in pain symptoms of osteoarthritis). |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

PERJETA (S)

Products Affected

- Perjeta

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Metastatic breast cancer: Diagnosis of human epidermal growth factor receptor 2 (HER2)-positive metastatic breast cancer. One of the following: a) patient has not received prior anti-HER2 therapy or chemotherapy for metastatic disease AND used in combination with Herceptin (trastuzumab) and a taxane (eg, docetaxel, paclitaxel), OR b) patient was previously treated with chemotherapy and Herceptin (trastuzumab) without Perjeta AND used in combination with Herceptin (trastuzumab). Non-metastatic breast cancer: One of the following diagnoses: HER2-positive early stage breast cancer, HER2-positive locally advanced breast cancer, or HER2-positive inflammatory breast cancer. Used in combination with both Herceptin (trastuzumab) and chemotherapy. Early Breast Cancer Adjuvant Treatment: Diagnosis of HER2-positive early breast cancer. Patient is at high risk of recurrence. Used in combination with both of the following: Herceptin (trastuzumab) and chemotherapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | All uses: Prescribed by or in consultation with an oncologist. |
| Coverage Duration | Non-metastatic breast cancer: 6 months. All others: 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

PIQRAY (S)

Products Affected

- Piqray 200mg Daily Dose
- Piqray 250mg Daily Dose
- Piqray 300mg Daily Dose

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Breast Cancer (BC): Diagnosis of advanced or metastatic BC. Disease is hormone receptor (HR)-positive, and human epidermal growth factor receptor 2 (HER2)-negative. Cancer is PIK3CA-mutated as detected by an FDA-approved test (therascreen PIK3CA RGQ PCR Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Patient is a postmenopausal woman or male. Used in combination with fulvestrant. Disease has progressed on or after an endocrine-based regimen. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

POMALYST (S)

Products Affected

- Pomalyst

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Multiple Myeloma (MM): Diagnosis of MM. Used in combination with dexamethasone. Patient has received two prior therapies, including Revlimid (lenalidomide) and a proteasome inhibitor [eg, Velcade (bortezomib) or Kyprolis (carfilzomib)] or has a contraindication or intolerance to Revlimid and proteasome inhibitors. Patient has experienced disease progression on or within 60 days of completion of last therapy. Kaposi sarcoma (KS): One of the following: 1) Both of the following: a) Diagnosis of AIDS-related KS and b) Patient has failed highly active antiretroviral therapy (HAART), OR 2) Both of the following: a) Diagnosis of KS and b) Patient is HIV-negative. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

PORTRAZZA (S)

Products Affected

- Portrazza

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | N/A |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

POTELIGEO (S)

Products Affected

- Poteligeo

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Mycosis fungoides (MF)/Sezary syndrome (SS): Diagnosis of one of the following: MF or SS. Disease is relapsed or refractory. Patient has received at least one prior systemic therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist/oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

PROCYSBI (S)

Products Affected

- Procysbi

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Nephropathic cystinosis: Diagnosis of nephropathic cystinosis, confirmed by elevated leukocyte cystine levels (LCL) or genetic analysis of the CTNS gene or demonstration of cysteine corneal crystals by slit lamp examination AND Trial and failure or intolerance to therapy with Cystagon (immediate-release cysteamine bitartrate). |
| Age Restrictions | 1 year of age or older |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

PROMACTA (S)

Products Affected

- Promacta

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Chronic immune (idiopathic) thrombocytopenic purpura (ITP) (initial): Diagnosis of one of the following: relapsed/refractory ITP or chronic ITP. Baseline platelet count is less than 30,000/mcL. Patient's degree of thrombocytopenia and clinical condition increase the risk of bleeding. Trial and failure, intolerance, contraindication to corticosteroids or immunoglobulins or splenectomy. Chronic hepatitis C (initial): Diagnosis of chronic hepatitis C-associated thrombocytopenia. One of the following: 1) Planning to initiate and maintain interferon-based treatment, or 2) currently receiving interferon-based treatment. First-line for severe aplastic anemia (SAA): Diagnosis of SAA. Used for first-line treatment (i.e., patient has not received prior immunosuppressive therapy). Used in combination with standard immunosuppressive therapy. Patient meets at least two of the following: 1) absolute neutrophil count less than 500/mcL, 2) platelet count less than 20,000/mcL, 3) absolute reticulocyte count less than 60,000/mcL. Refractory SAA (initial): Diagnosis of refractory severe aplastic anemia. Patient has a platelet count less than 30,000/mcL. Insufficient response to immunosuppressive therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Chronic ITP and SAA: Prescribed by or in consultation with a hematologist/oncologist. Chronic hepatitis C associated thrombocytopenia: Prescribed by or in consultation with a hematologist/oncologist, gastroenterologist, hepatologist, infectious disease specialist, or HIV specialist certified through the American Academy of HIV Medicine. |
| Coverage Duration | ITP(init, reauth): 12mo. HepC: 3mo(init), 12mo(reauth). 1stline SAA: 6mo. RefractSAA: 16wk-init, 12mo-reauth |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|-----------------------|---|
| Other Criteria | ITP (reauth): Documentation of positive clinical response to Promacta therapy as evidenced by an increase in platelet count to a level sufficient to avoid clinically important bleeding. Hepatitis C (reauth): One of the following: 1) For patients that started treatment with Promacta prior to initiation of treatment with interferon, Promacta will be approved when both of the following are met: a) Currently on antiviral interferon therapy for treatment of chronic hepatitis C and b) Documentation that the patient reached a threshold platelet count that allows initiation of antiviral interferon therapy with Promacta treatment by week 9, OR 2) For patients that started treatment with Promacta while on concomitant treatment with interferon, Promacta will be approved based on the following: Currently on antiviral interferon therapy for treatment of chronic hepatitis C. Refractory SAA (reauth): Documentation of positive clinical response to Promacta therapy as evidenced by an increase in platelet count. |
|-----------------------|---|

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

PROVIGIL (S)

Products Affected

- Modafinil

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | <p>Obstructive sleep apnea/hypopnea syndrome (OSAHS) (Initial): Diagnosis (dx) of OSAHS defined by one of the following: 15 or more obstructive respiratory events per hour of sleep confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible), or both of the following: 5 or more obstructive respiratory events per hour of sleep confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible), and 1 of the following symptoms: unintentional sleep episodes during wakefulness, daytime sleepiness, unrefreshing sleep, fatigue, insomnia, waking up breath holding/gasping/choking, loud snoring, or breathing interruptions during sleep. Shift-work sleep disorder (SWSD) (Initial):Dx of SWSD confirmed by one of the following: 1) Symptoms of excessive sleepiness or insomnia for at least 3 months, which is associated with a work period (usually night work) that occurs during the normal sleep period, OR 2) A sleep study demonstrating loss of a normal sleep-wake pattern (ie, disturbed chronobiologic rhythmicity). No other medical condition or medication accounts for the symptoms. Narcolepsy (initial): Dx of narcolepsy as confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible). MS Fatigue (initial): Dx of multiple sclerosis (MS). Patient is experiencing fatigue. Depression (initial): Treatment-resistant depression defined as diagnosis of major depressive disorder (MDD) or bipolar depression, AND trial and failure, contraindication, or intolerance to at least two antidepressants from different classes (eg, SSRIs, SNRIs, bupropion). Used as adjunctive therapy. Idiopathic Hypersomnia (Initial): Diagnosis of idiopathic hypersomnia as confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible).</p> |
| Age Restrictions | N/A |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|--------------------------------|--|
| Prescriber Restrictions | N/A |
| Coverage Duration | Narcolepsy: Init, Reauth: 12 mo. All other indications: Init, Reauth: 6 mo. |
| Other Criteria | OSAHS, Narcolepsy, Idiopathic Hypersomnia (Reauth): Documentation of positive clinical response to modafinil therapy. SWSD (Reauth): Documentation of positive clinical response to modafinil therapy. MS Fatigue (reauth): Patient is experiencing relief of fatigue with modafinil therapy. Depression (reauth): Documentation of positive clinical response to modafinil therapy. Used as adjunctive therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

PULMOZYME (S)

Products Affected

- Pulmozyme

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Cystic Fibrosis (CF) (Initial, Reauth): Diagnosis of CF. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | CF (initial, reauth): 12 months |
| Other Criteria | Part B vs D determination applies. CF (reauth): Patient is benefiting from treatment (i.e. improvement in lung function [forced expiratory volume in one second (FEV1)], decreased number of pulmonary exacerbations). |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

QINLOCK (S)

Products Affected

- Qinlock

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Gastrointestinal Stromal Tumor (GIST): Diagnosis of gastrointestinal stromal tumor (GIST). Disease is advanced. Patient has received prior treatment with three or more kinase inhibitors (e.g., sunitinib, regorafenib), one of which must include imatinib. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

QUALAQUIN (S)

Products Affected

- Quinine Sulfate CAPS 324MG

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Malaria: Diagnosis of uncomplicated malaria. One of the following: 1) Treatment in areas of chloroquine-sensitive malaria, and trial and failure, contraindication, or intolerance to chloroquine or hydroxychloroquine, OR 2) Treatment in areas of chloroquine-resistant malaria. Not used solely for the treatment or prevention of nocturnal leg cramps. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 7 days |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

RADICAVA (S)

Products Affected

- Radicava

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Amyotrophic lateral sclerosis (ALS) (initial): Submission of medical records (e.g., chart notes, previous medical history, diagnostic testing including: imaging, nerve conduction studies, laboratory values) to support a diagnosis of “definite” or “probable” ALS per the revised El Escorial diagnostic criteria. Patient has scores of greater than or equal to 2 in all items of the ALS Functional Rating Scale-Revised (ALSFRRS-R) criteria at the start of treatment. Patient has a percent forced vital capacity (%FVC) of greater than or equal to 80% at the start of treatment. |
| Age Restrictions | N/A |
| Prescriber Restrictions | ALS (initial): Prescribed by or in consultation with a neurologist. |
| Coverage Duration | Initial, reauth: 6 months |
| Other Criteria | ALS (reauthorization): Documentation of a benefit from therapy (e.g., slowing in the decline of functional abilities), and Patient is not dependent on invasive ventilation or tracheostomy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

RAVICTI (S)

Products Affected

- Ravicti

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Urea cycle disorders (UCDs) (Initial): Diagnosis of UCDs. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | UCDs (Initial, reauth): 12 months |
| Other Criteria | UCDs (reauth): Documentation of positive clinical response to Ravicti therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

REGRANEX (S)

Products Affected

- Regranex

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Diabetic neuropathic ulcers: Patient has a lower extremity diabetic neuropathic ulcer. Treatment will be given in combination with ulcer wound care (eg, debridement, infection control, and/or pressure relief). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 5 months |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

REMICADE (S)

Products Affected

- Remicade

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Crohn's Disease (CD) and Fistulizing Crohn's Disease (FCD) (Initial): Diagnosis (Dx) of moderately to severely active CD or FCD. Failure, contraindication, or intolerance to one of the following conventional therapies: 6-mercaptopurine, Imuran (azathioprine), corticosteroid (eg, prednisone, methylprednisolone), methotrexate (Rheumatrex/Trexall). Ulcerative colitis (UC) (Initial): Dx of moderately to severely active UC. Trial and failure, contraindication, or intolerance (TF/C/I) to one of the following conventional therapies: 6-mercaptopurine, Imuran (azathioprine), corticosteroid (eg, prednisone, methylprednisolone), aminosalicylate [eg, mesalamine (Asacol/Pentasa/Rowasa), Dipentum (olsalazine), Azulfidine (sulfasalazine)]. Rheumatoid arthritis (RA) (Initial): Dx of moderately to severely active RA. One of the following: Receiving concurrent therapy with methotrexate (Rheumatrex/Trexall), OR TF/C/I to methotrexate (Rheumatrex/Trexall). Ankylosing spondylitis (AS) (Initial): Dx of active AS. TF/C/I to two NSAIDs. Psoriatic arthritis (PsA) (Initial): Dx of active PsA. Plaque psoriasis (Initial): Dx of chronic severe (ie, extensive and/or disabling) plaque psoriasis. Sarcoidosis (initial): TF/C/I to one immunosuppressant [eg, methotrexate (Rheumatrex/Trexall), cyclophosphamide, or azathioprine (Imuran)] AND TF/C/I to one corticosteroid (eg, prednisone). |
| Age Restrictions | N/A |
| Prescriber Restrictions | CD, FCD, UC (initial): Prescribed by or in consultation with a gastroenterologist. RA, AS (initial): Prescribed by or in consultation with a rheumatologist. PsA (initial): Prescribed by or in consultation with rheumatologist or dermatologist. Plaque Psoriasis (initial): Prescribed by or in consultation with a dermatologist. Sarcoidosis (initial): Prescribed by or in consultation with a pulmonologist, dermatologist, ophthalmologist. |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|--------------------------|--|
| Coverage Duration | All uses (initial, reauth): 12 months |
| Other Criteria | Reauth (all indications): Documentation of positive clinical response to infliximab therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

REMODULIN (S)

Products Affected

- Remodulin

- Treprostinil

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH. |
| Age Restrictions | N/A |
| Prescriber Restrictions | PAH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist. |
| Coverage Duration | PAH: Initial: 6 months. Reauth: 12 months. |
| Other Criteria | Subject to Part B vs. D Review. PAH (Reauth): Documentation of positive clinical response to therapy. |

RENFLEXIS (S)

Products Affected

- Renflexis

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Crohn's Disease (CD) and Fistulizing Crohn's Disease (FCD) (initial): Dx of moderately to severely active CD or FCD. Ulcerative colitis (UC) (initial): Dx of moderately to severely active UC. Rheumatoid arthritis (RA) (initial): Diagnosis (Dx) of moderately to severely active RA. Ankylosing spondylitis (AS) (initial): Dx of active AS. Psoriatic arthritis (PsA) (initial): Dx of active PsA. Plaque psoriasis (initial): Dx of chronic severe (ie, extensive and/or disabling) plaque psoriasis. CD, FCD (initial): Trial and failure, contraindication or intolerance (TF/C/I) to one of the following conventional therapies: 6-mercaptopurine (Purinethol), azathioprine (Imuran), corticosteroids (eg, prednisone, methylprednisolone), methotrexate (Rheumatrex, Trexall). UC (initial): TF/C/I to one of the following conventional therapies: corticosteroids, aminosalicylate [eg, mesalamine (Asacol/Pentasa/Rowasa), olsalazine (Dipentum), sulfasalazine (Azulfidine/Sulfazine)], azathioprine (Imuran), 6-mercaptopurine (Purinethol). RA (initial): Receiving concurrent therapy with methotrexate (Rheumatrex/Trexall), or TF/C/I to methotrexate. AS (initial): TF/C/I to two or more NSAIDs. All indications (initial): One of the following: 1) Trial and failure or intolerance to Remicade or 2) For continuation of prior Renflexis therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Initial: RA, AS: Prescribed by or in consultation with a rheumatologist. PsA: Prescribed or in consultation with a rheumatologist or dermatologist. Crohn's Disease, Fistulizing Crohn's Disease, UC: Prescribed by or in consultation with a gastroenterologist. Plaque Psoriasis: Prescribed by or in consultation with a dermatologist. |
| Coverage Duration | All indications (initial, reauth): 12 months |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|-----------------------|---|
| Other Criteria | Reauthorization for all indications: Documentation of positive clinical response to infliximab therapy. |
|-----------------------|---|

REPATHA (S)

Products Affected

- Repatha
- Repatha Pushtronex System
- Repatha Sureclick

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | <p>HeFH/ASCVD/Primary HLD (init): One of the following dx: A)HeFH as confirmed by one of the following: 1)Both of the following: a)Untreated/pre-treatment LDL greater than 190 mg/dL in an adult, AND b)One of the following: i) Family hx of tendinous xanthomas and/or arcus cornealis in 1st degree relative, or 2nd degree relative, ii)Hx of myocardial infarction (MI) in 1st-degree relative less than 60 years of age, iii)Family hx of MI in 2nd-degree relative less than 50 years of age, iv)Family hx of LDL-C greater than 190 mg/dL in 1st- or 2nd-degree relative, v)Family hx of FH in 1st- or 2nd-degree relative, or 2)Untreated/pre-treatment LDL-C greater than 190 mg/dL in an adult AND submission of MR (eg chart notes, lab values) documenting one of the following: presence of tendinous xanthoma in pt, arcus cornealis before age 45, or functional mutation in the LDL receptor, ApoB, or PCSK9 gene. OR B)ASCVD as confirmed by ACS, hx of MI, stable or unstable angina, coronary or other arterial revascularization, stroke,TIA, or peripheral arterial disease presumed to be of atherosclerotic origin. OR C)Primary hyperlipidemia (HLD). HoFH (init): Sub of MR (eg, chart notes, lab values) documenting dx of HoFH as confirmed by one of the following: 1)Gen confirmation of 2 mutations in LDL receptor, ApoB, PCSK9, or LDLRAP1 or ARH, or 2)either untreated LDL greater than 500 or treated LDL greater than 300, AND either xanthoma before 10 yo or evidence of HeFH in both parents. HeFH/ASCVD/Primary HLD (init): One of the following: set A)Both of the following: a)One of the following LDL values while on max tolerated lipid lowering regimen w/in the last 120 days: 1)LDL greater than or equal to 100 mg/dL w/ASCVD, or 2)LDL greater than or equal to 130 mg/dL w/o ASCVD. AND b)One of the following: 1)Pt has been receiving at least 12 wks of one high-intensity (HI) statin therapy (tx) and will continue to receive a HI statin [ie, atorvastatin 40-80 mg, rosuvastatin 20-40 mg] at max tolerated dose.</p> |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

| | |
|--------------------------------|--|
| Age Restrictions | N/A |
| Prescriber Restrictions | Initial/Reauth: Prescribed by or in consultation with a cardiologist, endocrinologist, or lipid specialist |
| Coverage Duration | Initial: 6 months. Reauth: 12 months |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|-----------------------|--|
| Other Criteria | <p>Set A (cont, initial): OR (2) Both of the following: A) Pt unable to tolerate HI statin as evidenced by intolerable and persistent (ie, more than 2 wks) myalgia (muscle symptoms w/o CK elevations) or myositis (muscle symptoms w/ CK elevations less than 10 times ULN) AND B) One of the following: a) Pt has been receiving at least 12 wks of one moderate-intensity (MI) or low-intensity (LI) statin tx and will continue to receive a MI or LI statin [ie, atorvastatin 10-20 mg, rosuvastatin 5-10 mg, simvastatin 10-40 mg, pravastatin 10-80 mg, lovastatin 20-40 mg, Lescol XL (fluvastatin XL) 80 mg, fluvastatin 20-40 mg, or Livalo (pitavastatin) 1-4 mg] at max tolerated dose, OR b) Pt is unable to tolerate MI or LI statin as evidenced by intolerable and persistent (ie, more than 2 weeks) myalgia (muscle symptoms w/o CK elevations) or myositis (muscle symptoms w/ CK elevations less than 10 times ULN), OR (3) Submission of MR documenting pt has a labeled contraindication to all statins, OR (4) Pt has experienced rhabdomyolysis or muscle symptoms w/ statin treatment w/ CK elevations greater than 10 times ULN on one statin tx. OR set B) Both of the following: a) One of the following LDL values while on max tolerated statin tx w/in the last 120 days: (1) LDL b/t 70 and 99 mg/dL w/ ASCVD. (2) LDL b/t 100 and 129 mg/dL w/o ASCVD. AND b) Both of the following: (1) One of the following: i) Pt has been receiving at least 12 wks of one max-tolerated statin tx and will continue to receive a statin at max tolerated dose, ii) pt is unable to tolerate statin tx as evidenced by intolerable and persistent (ie, more than 2 wks) myalgia (muscle symptoms w/o CK elevations) or myositis (muscle symptoms w/ CK elevations less than 10 times ULN, iii) Submission of MR documenting patient has a labeled contraindication to all statins, or iv) Pt has experienced rhabdomyolysis or muscle symptoms w/ statin tx w/ CK elevations greater than 10 times ULN on one statin tx and (2) Pt has been receiving at least 12 weeks of ezetimibe (Zetia) tx as adjunct to max tolerated statin tx OR Pt has a hx of contraindication or intolerance to ezetimibe. HoFH (init): Pt is receiving other lipid-lowering tx (eg statin, ezetimibe).HeFH/ASCVD/Primary HLD (reauth): Pt continues to receive other lipid-lowering tx (eg statins, ezetimibe) at max tolerated dose (unless pt has documented inability to take these medications). HoFH (reauth): Pt continues to receive other lipid-lowering tx (eg statin, ezetimibe). HeFH/ASCVD/Primary HLD/HoFH (reauth): Sub of MR (eg, lab values) documenting LDL reduction while on Repatha tx. HeFH/ASCVD/Primary HLD/HoFH (Init, reauth): Prescriber attests that the info provided is true and accurate to the best of their knowledge and they understand that OptumRx may perform a routine audit and request the medical info necessary to verify the accuracy of the info provided. HoFH (Init, reauth): Not used in combo w/ Juxtapid.</p> |
|-----------------------|--|

RETACRIT (S)

Products Affected

- Retacrit

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | <p>Anemia with Chronic Kidney Disease (CKD) (Initial): Diagnosis (Dx) of CKD. Anemia by lab values (Hct less than 30% or Hgb less than 10 g/dL) collected within 30 days of request. One of the following: a) both of the following: Patient is on dialysis, patient is without ESRD OR b) all of the following: patient is not on dialysis, the rate of hemoglobin decline indicates the likelihood of requiring a red blood cell (RBC) transfusion, and reducing the risk of alloimmunization and/or other RBC transfusion-related risks is a goal. Anemia with chemo (Initial): Other causes of anemia have been ruled out. Anemia by lab values (Hct less than 30%, Hgb less than 10 g/dL) collected within the prior 2 weeks of request. Cancer is a non-myeloid malignancy. Patient is concurrently on chemo, or will receive concomitant chemo for a minimum of 2 mos, or anemia is caused in part by cancer chemo. Preoperative for reduction of allogeneic blood transfusion: Patient is scheduled to undergo elective, non-cardiac, non-vascular surgery. Hgb is greater than 10 to less than or equal to 13 g/dL. Patient is at high risk for perioperative transfusions. Patient is unwilling or unable to donate autologous blood pre-operatively. Anemia in hepatitis C virus (HCV)-infected pts due to ribavirin in combination with interferon/peg-interferon (Initial): Dx of HCV infection. Anemia by labs (Hct less than 36% or Hgb less than 12 g/dL) collected within 30 days of request. Patient is receiving ribavirin and one of the following: interferon alfa or peginterferon alfa. Anemia with HIV (Initial): Anemia by lab values (Hgb less than 12 g/dL or Hct less than 36%) collected within 30 days of request. Serum erythropoietin level less than or equal to 500 mU/mL. Receiving zidovudine therapy or dx of HIV. Anemia in Myelodysplastic Syndrome (MDS) (Initial): Dx of MDS. Serum erythropoietin level is 500 mU/mL or less, or dx of transfusion-dependent MDS.</p> |
| Age Restrictions | N/A |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|--------------------------------|---|
| Prescriber Restrictions | N/A |
| Coverage Duration | CKD,HIV(Init):6mo. CKD,HIV(reauth):12mo. Chemo,HCV(all):3mo. MDS:(init) 3mo,(reauth)12mo. Preop:1mo. |
| Other Criteria | <p>Subject to ESRD review. CKD (Reauth): Dx of CKD. One of the following: 1) Most recent or average (avg) Hct over 3 months is 33% or less (Hgb is 11 g/dL or less) for patients on dialysis, without ESRD, 2) Most recent or avg Hct over 3 mo is 30% or less (Hgb 10 g/dL or less) for patients not on dialysis, OR 3) Most recent or avg Hct over 3 mo is 36% or less (Hgb 12 g/dL or less) for pediatric patients. Decrease in the need for blood transfusion or Hgb increased by 1 g/dL or more from pre-treatment level. HIV (Reauth): Most recent or avg Hct over 3 months is below 36% or most recent or avg Hgb over 3 months is below 12 g/dl. Decrease in the need for blood transfusion or Hgb increased by 1 g/dL or more from pre-treatment level. Chemo (Reauth): Anemia by lab values (Hgb less than 10 g/dl or Hct less than 30%) collected within the prior 2 weeks of request. Decrease in the need for blood transfusion or Hgb increased by 1 g/dL or more from pre-treatment level. Patient is concurrently on chemo, or will receive concomitant chemo for a minimum of 2 months, or anemia is caused in part by cancer chemo. HCV (Reauth): Most recent or avg Hct over 3 months is 36% or less, OR most recent or avg Hgb over 3 months is 12 g/dl or less. Decrease in the need for blood transfusion or Hgb increased by 1 g/dL or more from pre-treatment level. If patient has demonstrated response to therapy, authorization will be issued for the full course of ribavirin therapy. MDS (Reauth): Most recent or avg Hct over 3 months is 36% or less, OR most recent or avg Hgb over 3 months is 12 g/dl or less. Decrease in the need for blood transfusion or Hgb increased by 1 g/dL or more from pre-treatment level. Other Off-label uses (except MDS, HCV): Will not be approved if patient has Hgb greater than 10 g/dL or Hct greater than 30%. CKD (init, reauth), HIV (init), Chemo (init), Preop, MDS (init), HCV (init): Verify iron evaluation for adequate iron stores.</p> |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

RETEVMO (S)

Products Affected

- Retevmo

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Lung Cancer: Diagnosis of metastatic non-small cell lung cancer (NSCLC). Disease has presence of RET gene fusion-positive tumor(s). Medullary Thyroid Cancer (MTC): Diagnosis of medullary thyroid cancer (MTC). Disease is advanced or metastatic. Disease has presence of RET gene mutation tumor(s). Disease requires treatment with systemic therapy. Thyroid Cancer: Diagnosis of thyroid cancer. Disease is advanced or metastatic. Disease has presence of RET gene fusion-positive tumor(s). Disease requires treatment with systemic therapy. Patient is radioactive iodine-refractory or radioactive iodine therapy is not appropriate. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Lung Cancer, MTC: Prescribed by or in consultation with an oncologist. Thyroid Cancer: Prescribed by or in consultation with an endocrinologist or an oncologist. |
| Coverage Duration | Lung Cancer, MTC, Thyroid Cancer: 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

REVATIO (S)

Products Affected

- Sildenafil Citrate TABS 20MG

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH. |
| Age Restrictions | N/A |
| Prescriber Restrictions | PAH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist. |
| Coverage Duration | PAH: Initial: 6 months. Reauth: 12 months. |
| Other Criteria | PAH (Reauth): Documentation of positive clinical response to therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

REVATIO INJECTION (S)

Products Affected

- Sildenafil INJ

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH. Patient is unable to take oral medications. |
| Age Restrictions | N/A |
| Prescriber Restrictions | PAH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist. |
| Coverage Duration | PAH: Initial: 6 months. Reauth: 12 months. |
| Other Criteria | PAH (Reauth): Documentation of positive clinical response to therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

REVATIO SUSPENSION (S)

Products Affected

- Sildenafil Citrate SUSR

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH. One of the following: A) Intolerance to generic Revatio tablets, OR B) Patient is unable to ingest a solid dosage form (e.g., an oral tablet or capsule) due to one of the following: age, oral-motor difficulties, or dysphagia. |
| Age Restrictions | N/A |
| Prescriber Restrictions | PAH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist. |
| Coverage Duration | PAH: Initial: 6 months. Reauth: 12 months. |
| Other Criteria | PAH (Reauth): Documentation of positive clinical response to therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

REVCovi (S)

Products Affected

- Renvovi

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Diagnosis of adenosine deaminase deficiency (ADA) with severe combined immunodeficiency (SCID). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

REVLIMID (S)

Products Affected

- Revlimid

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Multiple myeloma (MM): Diagnosis of MM. Either used as 1) combination therapy with dexamethasone, or 2) maintenance therapy following autologous hematopoietic stem cell transplantation (auto-HSCT). Myelodysplastic syndromes (MDS): Diagnosis of transfusion-dependent anemia due to low- or intermediate-1-risk MDS associated with a deletion 5q. Mantle cell lymphoma (MCL): Diagnosis of MCL. Disease has relapsed after, is refractory to, or progressed after at least one prior therapy (eg, bortezomib, bendamustine, cladribine, rituximab). Follicular Lymphoma (FL): Diagnosis of FL that has been previously treated. Used in combination with a rituximab product. Marginal Zone Lymphoma (MZL): Diagnosis of MZL that has been previously treated. Used in combination with a rituximab product. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist/hematologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

REYVOW (S)

Products Affected

- Reyvow

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Initial: Diagnosis of migraine with or without aura. Will be used for the acute treatment of migraine. Will not be used for preventive treatment of migraine. Patient has less than 15 headache days per month. Trial and failure or intolerance to one triptan (e.g., eletriptan, rizatriptan, sumatriptan) or a contraindication to all triptans. If patient has 4 or more headache days per month, patient must meet one of the following: a) currently being treated with Elavil (amitriptyline) or Effexor (venlafaxine) unless there is a contraindication or intolerance to these medications, OR b) currently being treated with Depakote/Depakote ER (divalproex sodium) or Topamax (topiramate) unless there is a contraindication or intolerance to these medications, OR c) currently being treated with a beta blocker (i.e., atenolol, propranolol, nadolol, timolol, or metoprolol) unless there is a contraindication or intolerance to these medications. Will not be used concomitantly with central nervous system (CNS) depressants (e.g., alprazolam, phenobarbital, alcohol). Prescriber attests that the patient has been counseled and has agreed to adhere to the following: Will follow instructions to not drive or operate machinery until at least 8 hours after taking each dose of Reyvow. |
| Age Restrictions | Initial: 18 years of age or older. |
| Prescriber Restrictions | Initial, Reauth: Prescribed by or in consultation with a neurologist or pain specialist. |
| Coverage Duration | Initial: 3 months. Reauth: 12 months. |
| Other Criteria | Reauth: Patient has experienced a positive response to therapy (e.g., reduction in pain, photophobia, phonophobia, nausea). Will not be used for preventive treatment of migraine. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

RILUTEK (S)

Products Affected

- Riluzole

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Amyotrophic lateral sclerosis (ALS): Diagnosis of amyotrophic lateral sclerosis (ALS). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | ALS: 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

RINVOQ (S)

Products Affected

- Rinvoq

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Rheumatoid arthritis (RA) (initial): Diagnosis of moderately to severely active RA. One of the following: Trial and failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR patient has a documented needle-phobia to the degree that the patient has previously refused any injectable therapy or medical procedure (refer to DSM-V-TR 300.29/F40.2 for specific phobia diagnostic criteria), OR for continuation of prior Rinvoq therapy. Patient is not receiving Rinvoq in combination with a potent immunosuppressant (e.g., azathioprine, cyclosporine). |
| Age Restrictions | N/A |
| Prescriber Restrictions | RA (initial): Prescribed by or in consultation with a rheumatologist. |
| Coverage Duration | RA (initial, reauth): 12 months. |
| Other Criteria | RA (reauth): Documentation of positive clinical response to therapy. Patient is not receiving Rinvoq in combination with a potent immunosuppressant (e.g., azathioprine, cyclosporine). |

RITUXAN (S)

Products Affected

- Rituxan

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | <p>Non-Hodgkin's Lymphoma (NHL): As first-line treatment of diffuse large B-cell, CD20-positive, NHL in combination with CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone) or other anthracycline-based chemotherapy regimens, or as first-line treatment of follicular, CD20-positive, B-cell NHL in combination with chemotherapy, or for the treatment of follicular, CD20-positive, B-cell NHL in patients achieving a complete or partial response to Rituxan in combination with chemotherapy, used as monotherapy for maintenance therapy, or for the treatment of low-grade, CD20-positive, B-cell NHL in patients with stable disease following first-line treatment with CVP (cyclophosphamide, vincristine, prednisolone/prednisone) chemotherapy or who achieve a partial or complete response following first-line treatment with CVP (cyclophosphamide, vincristine, prednisolone/prednisone) chemotherapy, or diagnosis of relapsed or refractory, low grade or follicular CD20-positive, B-cell non-Hodgkin's lymphoma. Rheumatoid Arthritis (RA) (init): Diagnosis of moderately to severe active RA, Concurrently on or contraindication, or intolerance to methotrexate. Trial and failure, contraindication, or intolerance (TF/C/I) to a TNF antagonist (eg, adalimumab, etanercept, infliximab). Wegener's Granulomatosis (WG) and Microscopic Polyangiitis (MPA): Diagnosis of WG or MPA, Patient is concurrently on glucocorticoids (eg, prednisone) OR contraindication or intolerance to glucocorticoids (eg, prednisone). Immune or idiopathic thrombocytopenic purpura (ITP): Diagnosis of ITP, TF/C/I to one of the following: corticosteroids, immunoglobulins, or splenectomy. Documented platelet count of less than $50 \times 10^9 /L$. Chronic Lymphocytic Leukemia (CLL): Diagnosis of CLL, used in combination with fludarabine and cyclophosphamide. Pemphigus Vulgaris (PV): Diagnosis of moderate to severe PV, used in combination with a tapering course of glucocorticoids.</p> |
| Age Restrictions | N/A |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|--------------------------------|---|
| Prescriber Restrictions | ITP, CLL, NHL: Prescribed by or in consultation with a hematologist or oncologist. RA: Prescribed by or in consultation with a rheumatologist. WG, MPA: Prescribed by or in consultation with a nephrologist, pulmonologist, or rheumatologist. PV: Prescribed by or in consultation with a dermatologist |
| Coverage Duration | All uses except RA, WG, MPA: 12 mos. RA: 3 months. WG, MPA: 3 months only. |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

RITUXAN HYCELA (S)

Products Affected

- Rituxan Hycela

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | <p>Follicular Lymphoma: Diagnosis of follicular lymphoma. One of the following: 1) Disease is relapsed or refractory OR 2) Patient exhibited complete or partial response to prior treatment with rituximab in combination with chemotherapy OR 3) Disease is non-progressing or stable following prior treatment with first-line cyclophosphamide, vincristine, and prednisone (CVP) chemotherapy OR 4) Both of the following: a) Disease is previously untreated AND b) Medication is used in combination with first-line chemotherapy. Patient receives a full induction dose of intravenous rituximab prior to initiation of therapy.</p> <p>Diffuse Large B-Cell Lymphoma: 1) Diagnosis of diffuse large B-cell lymphoma AND 2) Disease is previously untreated AND 3) Medication is being used in combination with cyclophosphamide, doxorubicin, vincristine, prednisone (CHOP) or other anthracycline-based chemotherapy AND 4) Patient receives a full induction dose of intravenous rituximab prior to initiation of therapy.</p> <p>Chronic Lymphocytic Leukemia: 1) Diagnosis of chronic lymphocytic leukemia AND 2) Medication is being used in combination with fludarabine and cyclophosphamide (FC) therapy AND 3) Patient receives a full induction dose of intravenous rituximab prior to initiation of therapy.</p> |
| Age Restrictions | N/A |
| Prescriber Restrictions | All uses: Prescribed by or in consultation with a hematologist or oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ROZLYTREK (S)

Products Affected

- Rozlytrek

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Non-small cell lung cancer (NSCLC): Diagnosis of metastatic non-small cell lung cancer (NSCLC). Patient has ROS1 rearrangement positive tumor(s). Solid Tumors: Patient has solid tumors with a neurotrophic tyrosine receptor kinase (NTRK) gene fusion (e.g., ETV6-NTRK3, TPM3-NTRK1, TPR-NTRK1, etc.). Disease is without a known acquired resistance mutation (e.g., TRKA G595R, TRKA G667C or TRKC G623R substitutions). Disease is one of the following: metastatic or unresectable (including cases where surgical resection is likely to result in severe morbidity). One of the following: disease has progressed following previous treatment (e.g., surgery, radiation therapy, or systemic therapy) or disease has no satisfactory alternative treatments. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

RUBRACA (S)

Products Affected

- Rubraca

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Ovarian cancer: Diagnosis of epithelial ovarian cancer, fallopian tube cancer, or primary peritoneal cancer. One of the following: 1) Both of the following: a) Presence of deleterious BRCA mutation as detected by a U.S. Food and Drug Administration (FDA)-approved diagnostic test (e.g., FoundationFocus CDxBRCA Assay) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) and b) Trial and failure, contraindication, or intolerance to two or more chemotherapies (e.g., cisplatin, carboplatin), OR 2) Both of the following: a) Disease is recurrent and b) Used for maintenance treatment in patients who are in a complete or partial response to platinum-based chemotherapy (e.g., cisplatin, carboplatin). Prostate cancer: Diagnosis of metastatic castration-resistant prostate cancer. Presence of deleterious BRCA mutation as detected by an FDA-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Patient has received previous treatment with both of the following: 1) Androgen receptor-directed therapy [e.g., Erleada (apalutamide), Xtandi (enzalutamide), Zytiga (abiraterone)], AND 2) A taxane-based chemotherapy [e.g., docetaxel, Jevtana (cabazitaxel)]. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Ovarian cancer: Prescribed by or in consultation with an oncologist. Prostate cancer: Prescribed by or in consultation with an oncologist or urologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

RUCONEST (S)

Products Affected

- Ruconest

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Treatment of hereditary angioedema (HAE) attacks: Diagnosis of HAE. For the treatment of acute HAE attacks. Not used in combination with other approved treatments for acute HAE attacks. |
| Age Restrictions | N/A |
| Prescriber Restrictions | HAE: Prescribed by or in consultation with an immunologist, allergist, or rheumatologist |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

RUZURGI (S)

Products Affected

- Ruzurgi

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Lambert-Eaton Myasthenic Syndrome (LEMS) (initial): Diagnosis of LEMS. |
| Age Restrictions | N/A |
| Prescriber Restrictions | LEMS (initial): Prescribed by or in consultation with a neurologist. |
| Coverage Duration | LEMS (initial): 3 months. LEMS (reauth): 12 months. |
| Other Criteria | LEMS (reauth): Documentation of positive clinical response to therapy (e.g., improvement in dynamometry, Timed 25-Foot Walk Test, Timed Up and Go Test). |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

RYDAPT (S)

Products Affected

- Rydapt

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Acute Myeloid Leukemia (AML): Newly diagnosed acute myeloid leukemia (AML), FMS-like tyrosine kinase 3 (FLT3) mutation-positive as detected by a U.S. Food and Drug Administration (FDA)-approved test (e.g., LeukoStrat CDx FLT3 Mutation Assay) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA), used in combination with standard cytarabine and daunorubicin induction and cytarabine consolidation. Aggressive Systemic Mastocytosis (ASM), Systemic Mastocytosis with Associated Hematological Neoplasm (SM-AHN), Mast Cell Leukemia (MCL): Diagnosis of one of the following: aggressive systemic mastocytosis (ASM), systemic mastocytosis with associated hematological neoplasm (SM-AHN), or mast cell leukemia (MCL). |
| Age Restrictions | N/A |
| Prescriber Restrictions | All uses: Prescribed by or in consultation with a hematologist or oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

SABRIL (S)

Products Affected

- Sabril TABS
- Vigabatrin
- Vigadrone

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Complex Partial Seizures (CPS): For use as adjunctive therapy. Failure, contraindication, or intolerance to two formulary anticonvulsants [eg, Lamictal (lamotrigine), Depakene (valproic acid), Dilantin (phenytoin)]. Infantile Spasms (IS): Diagnosis of infantile spasms. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020
SANDOSTATIN (S)

Products Affected

- Octreotide Acetate

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Acromegaly (initial): Diagnosis of acromegaly. One of the following: A) Inadequate response to surgical resection and/or pituitary irradiation OR B) Patient is not a candidate for surgical resection or pituitary irradiation. Trial and failure, contraindication or intolerance to a dopamine agonist (e.g., bromocriptine or cabergoline) at maximally tolerated doses. Carcinoid tumor (initial): Diagnosis of metastatic carcinoid tumor requiring symptomatic treatment of severe diarrhea or flushing episodes. Vasoactive intestinal peptide tumor (initial): Diagnosis of vasoactive intestinal peptide tumor requiring treatment of profuse watery diarrhea. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | All uses (initial, reauth): 12 months |
| Other Criteria | Acromegaly (reauth): Documentation of positive clinical response to therapy (e.g., reduction or normalization of IGF-1/GH level for same age and sex, reduction in tumor size). Carcinoid tumor (reauth): Patient has improvement in number of diarrhea or flushing episodes. Vasoactive intestinal peptide tumor (reauth): Patient has improvement in number of diarrhea episodes. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

SANDOSTATIN LAR (S)

Products Affected

- Sandostatin Lar Depot

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Acromegaly (initial): Diagnosis of acromegaly. One of the following: A) Inadequate response to surgical resection and/or pituitary irradiation OR B) Patient is not a candidate for surgical resection or pituitary irradiation. Trial and failure or intolerance to a dopamine agonist (e.g., bromocriptine or cabergoline) at maximally tolerated doses. Patient had a trial of short-acting octreotide and responded to and tolerated therapy. Carcinoid tumor (initial): Diagnosis of metastatic carcinoid tumor requiring symptomatic treatment of severe diarrhea or flushing episodes. Patient had a trial of short-acting octreotide and responded to and tolerated therapy. Vasoactive intestinal peptide tumor (initial): Diagnosis of vasoactive intestinal peptide tumor requiring treatment of profuse watery diarrhea. Patient had a trial of short-acting octreotide and responded to and tolerated therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | All uses (initial, reauth): 12 months |
| Other Criteria | Acromegaly (reauth): Documentation of positive clinical response to therapy (e.g., reduction or normalization of IGF-1/GH level for same age and sex, reduction in tumor size). Carcinoid tumor (reauth): Patient has improvement in number of diarrhea or flushing episodes. Vasoactive intestinal peptide tumor (reauth): Patient has improvement in number of diarrhea episodes. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

SCIG (S)

Products Affected

- Cutaquig
- Cuvitru
- Hyqvia
- Xembify

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | All uses (initial, reauth): Contraindications to immune globulin therapy (i.e., IgA deficiency with antibodies to IgA and a history of hypersensitivity or product specific contraindication). |
| Required Medical Information | Initial: Immune globulin will be administered at the minimum effective dose and appropriate frequency for the prescribed diagnosis. Medication is being used subcutaneously. Diagnosis of chronic inflammatory demyelinating polyneuropathy (CIDP) OR one of the following FDA-approved or literature supported diagnoses: 1) Common variable immunodeficiency (CVID), OR 2) Congenital agammaglobulinemia (X-linked or autosomal recessive), OR 3) Severe combined immunodeficiencies (SCID), OR 4) Wiskott-Aldrich syndrome, OR 5) Other primary immunodeficiency with an immunologic evaluation including IgG levels below the normal laboratory value for the patient's age at the time of diagnosis and the patient lacks an adequate response to protein and polysaccharide antigens (i.e., tetanus toxoid or diphtheria toxoid and pneumovax or HiB vaccine). |
| Age Restrictions | N/A |
| Prescriber Restrictions | All uses (initial, reauth): Prescribed by or in consultation with a physician who has specialized expertise in managing patients on SCIG therapy (e.g., immunologist, hematologist, neurologist). |
| Coverage Duration | Initial, reauth: 12 months |
| Other Criteria | Subject to Part B vs. Part D review. All uses (reauth): Patient has experienced an objective improvement on immune globulin therapy and the immune globulin will be administered at the minimum effective dose (by decreasing the dose, increasing the frequency, or implementing both strategies) for maintenance therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

SECUADO (S)

Products Affected

- Secuado

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Diagnosis of schizophrenia. Both of the following: 1) Trial and failure of Saphris (asenapine) and 2) Trial and failure, contraindication, or intolerance to one of the following generic formulary atypical antipsychotic agents: aripiprazole, olanzapine, paliperidone, quetiapine, risperidone, or ziprasidone. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

SEROSTIM (S)

Products Affected

- Serostim INJ 4MG, 5MG, 6MG

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | HIV wasting (Initial): Diagnosis of HIV-associated wasting syndrome or cachexia, and one of the following: unintentional weight loss greater than 10% over the last 12 months, or unintentional weight loss greater than 7.5% over the last 6 months, or loss of 5% body cell mass (BCM) within 6 months, or body mass index (BMI) less than 20 kg/m ² , or patient is male and has BCM less than 35% of total body weight (TBW) and BMI less than 27 kg/m ² , or patient is female and has BCM less than 23% of TBW and BMI less than 27 kg/m ² . Nutritional evaluation since onset of wasting first occurred. Anti-retroviral tx has been optimized to decrease the viral load. Patient has not had weight loss as a result of other underlying treatable conditions (eg, depression, mycobacterium avium complex, chronic infectious diarrhea, or malignancy with the exception of Kaposi's sarcoma limited to skin or mucous membranes). Patient has tried and had an inadequate response or intolerance to dronabinol or megestrol acetate. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Initial/Reauth: Prescribed by or in consultation with an infectious disease specialist. |
| Coverage Duration | Initial: 3 months. Reauth: 6 months |
| Other Criteria | HIV wasting (reauth): Evidence of positive response to therapy. One of the following targets or goals has not been achieved: weight, BCM, BMI. Patient is currently receiving treatment with antiretrovirals. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

SIGNIFOR (S)

Products Affected

- Signifor

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Cushing's disease (initial): Diagnosis of Cushing's disease. One of the following: a) Pituitary surgery has not been curative for the patient or b) Patient is not a candidate for pituitary surgery. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Cushing's disease (initial): Prescribed by or in consultation with an endocrinologist. |
| Coverage Duration | Cushing's disease (initial, reauth): 12 months |
| Other Criteria | Cushing's disease (reauth): Documentation of positive clinical response to therapy (e.g., a clinically meaningful reduction in 24-hour urinary free cortisol levels, improvement in signs or symptoms of the disease). |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

SIGNIFOR LAR (S)

Products Affected

- Signifor Lar

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Acromegaly (initial): Diagnosis of acromegaly. One of the following: a) Inadequate response to surgery or b) Patient is not a candidate for surgery. Cushing's disease (initial): Diagnosis of Cushing's disease. One of the following: a) Pituitary surgery has not been curative for the patient or b) Patient is not a candidate for pituitary surgery. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Cushing's disease (initial): Prescribed by or in consultation with an endocrinologist. |
| Coverage Duration | Acromegaly: Initial: 6 months, Reauth: 12 months. Cushing's disease (init, reauth): 12 months |
| Other Criteria | Acromegaly (reauth): Documentation of positive clinical response to therapy (e.g., patient's growth hormone (GH) level or insulin-like growth factor 1 (IGF-1) level for age and gender has normalized/improved). Cushing's disease (reauth): Documentation of positive clinical response to therapy (e.g., a clinically meaningful reduction in 24-hour urinary free cortisol levels, improvement in signs or symptoms of the disease). |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

SIKLOS (S)

Products Affected

- Siklos

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Sickle Cell Anemia: Diagnosis of sickle cell anemia. Patient has moderate to severe painful crises. One of the following: 1) Patient is less than 18 years of age or 2) Trial and failure, or intolerance to Droxia. |
| Age Restrictions | Patient is 2 years of age or older |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

SILIQ (S)

Products Affected

- Siliq

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Plaque psoriasis (Initial): Diagnosis of moderate to severe plaque psoriasis. One of the following: a) Trial and failure, contraindication, or intolerance to Cosentyx (secukinumab) AND one of the following: Enbrel (etanercept) or Humira (adalimumab), OR b) for continuation of prior Siliq therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Plaque psoriasis (initial): Prescribed by or in consultation with a dermatologist. |
| Coverage Duration | Plaque psoriasis (Initial, reauth): 12 months |
| Other Criteria | Plaque psoriasis (Reauth): Documentation of positive clinical response to Siliq therapy. |

SIMPONI (S)

Products Affected

- Simponi

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Rheumatoid Arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. One of the following: Receiving concurrent therapy with methotrexate (Rheumatrex/Trexall) OR trial and failure, contraindication, or intolerance (TF/C/I) to methotrexate (Rheumatrex/Trexall). One of the following: TF/C/I to both Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR for continuation of prior therapy. Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. One of the following: TF/C/I to both Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR for continuation of prior therapy. Ankylosing Spondylitis (AS) (Initial): Diagnosis of active AS. One of the following: TF/C/I to both Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR for continuation of prior therapy. Ulcerative Colitis (UC) (Initial): Diagnosis of moderately to severely active UC. Patient is corticosteroid dependent (ie, an inability to successfully taper corticosteroids without a return of the symptoms of UC), OR TF/C/I to one of the following: oral aminosalicylate, oral corticosteroid, azathioprine, 6-mercaptopurine. One of the following: TF/C/I to Humira (adalimumab), OR for continuation of prior Simponi therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | RA, AS (initial): Prescribed by or in consultation with a rheumatologist. PsA (initial): Prescribed by or in consultation with a rheumatologist or dermatologist. UC (initial): Prescribed by or in consultation with a gastroenterologist. |
| Coverage Duration | UC (Initial): 12 weeks. UC (Reauth): 12 months. RA, AS, PsA (Initial, reauth): 12 months |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|-----------------------|---|
| Other Criteria | All indications (Reauth): Documentation of positive clinical response to Simponi therapy. |
|-----------------------|---|

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020
SIMPONI ARIA (S)

Products Affected

- Simponi Aria

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Rheumatoid arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. One of the following: Receiving concurrent therapy with methotrexate (Rheumatrex/Trexall), OR trial and failure, contraindication, or intolerance (TF/C/I) to methotrexate (Rheumatrex/Trexall). One of the following: trial and failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR for continuation of prior therapy. Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. One of the following: TF/C/I to both Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR for continuation of prior therapy. Ankylosing Spondylitis (AS) (Initial): Diagnosis of active AS. One of the following: TF/C/I to both Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR for continuation of prior therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | RA, AS (initial): Prescribed by or in consultation with a rheumatologist. PsA (initial): Prescribed by or in consultation with a rheumatologist or dermatologist. |
| Coverage Duration | RA, AS, PsA (Initial, reauth): 12 months |
| Other Criteria | All Indications (Reauth): Documentation of positive clinical response to Simponi therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

SKYRIZI (S)

Products Affected

- Skyrizi

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Plaque psoriasis (Initial): Diagnosis of moderate to severe plaque psoriasis. One of the following: a) Trial and failure, contraindication, or intolerance to Cosentyx (secukinumab) AND one of the following: Enbrel (etanercept) or Humira (adalimumab), OR b) for continuation of prior Skyrizi therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Plaque psoriasis (initial): Prescribed by or in consultation with a dermatologist. |
| Coverage Duration | Plaque psoriasis (Initial, reauth): 12 months |
| Other Criteria | Plaque psoriasis (Reauth): Documentation of positive clinical response to Skyrizi therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

SOLIRIS (S)

Products Affected

- Soliris

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Paroxysmal Nocturnal Hemoglobinuria (PNH), Atypical Hemolytic Uremic Syndrome (aHUS) (initial): Diagnosis of PNH or aHUS. Generalized Myasthenia Gravis (gMG) (initial): Diagnosis of gMG. Patient is anti-acetylcholine (AChR) antibody positive. One of the following: 1) Trial and failure, contraindication, or intolerance (TF/C/I) to two immunosuppressive therapies (e.g., glucocorticoids, azathioprine, cyclosporine, mycophenolate mofetil, methotrexate, tacrolimus), or 2) TF/C/I to one immunosuppressive therapy (e.g., glucocorticoids, azathioprine, cyclosporine, mycophenolate mofetil, methotrexate, tacrolimus), and TF/C/I to chronic plasmapheresis/plasma exchange (PE) or intravenous immunoglobulin (IVIG). |
| Age Restrictions | N/A |
| Prescriber Restrictions | gMG (initial): Prescribed by or in consultation with a neurologist. |
| Coverage Duration | All uses (initial, reauth): 12 months |
| Other Criteria | PNH (reauth): Documentation of positive clinical response (e.g., hemoglobin stabilization, decrease in the number of red blood cell transfusions) to Soliris therapy. aHUS (reauth): Documentation of positive clinical response (e.g., increase in mean platelet counts, hematologic normalization) to Soliris therapy. gMG (reauth): Documentation of positive clinical response to Soliris therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

SOMATULINE DEPOT (S)

Products Affected

- Somatuline Depot

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Acromegaly: Diagnosis of acromegaly. One of the following: A) Inadequate response to one of the following: surgery or radiotherapy, OR B) Not a candidate for one of the following: surgery or radiotherapy. Gastroenteropancreatic neuroendocrine tumors (GEP-NETs) (120mg/0.5mL strength only): Diagnosis of GEP-NETs. Disease is one of the following: (a) unresectable, locally advanced or (b) metastatic. Carcinoid syndrome (120mg/0.5mL strength only): Diagnosis of carcinoid syndrome. Used to reduce the frequency of short-acting somatostatin analog rescue therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | All uses: 12 months |
| Other Criteria | All Indications: Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

SOMAVERT (S)

Products Affected

- Somavert

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Acromegaly (initial): Diagnosis of acromegaly AND Failure to surgery and/or radiation therapy and/or other medical therapies (such as dopamine agonists [e.g., bromocriptine, cabergoline]) unless patient is not a candidate for these treatment options AND trial and failure or intolerance to generic octreotide (a somatostatin analogue) |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an endocrinologist |
| Coverage Duration | Initial and reauth: 12 months |
| Other Criteria | Acromegaly (reauth): Patient has experienced a positive clinical response to therapy (biochemical control, decrease or normalization of IGF-1 levels). |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

SOVALDI (S)

Products Affected

- Sovaldi TABS 200MG

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Criteria will be applied consistent with current AASLD/IDSA guideline. Diagnosis of chronic hepatitis C. All GT1 (except Sovaldi plus Daklinza therapy in liver transplant (tx) patients) and GT4: 1) trial and failure, intolerance or contraindication (TF/I/C) to both of the following: a) sofosbuvir/velpatasvir and b) Mavyret OR 2) For continuation of prior Sovaldi therapy. For GT2 (except liver tx patients, or pediatric patients younger than 12 years of age and weighing between 35-44 kg) or GT3 patients (except pediatric patients younger than 12 years of age and weighing between 35-44 kg), using Sovaldi plus ribavirin: TF/I/C to a) sofosbuvir/velpatasvir AND Mavyret OR b) for continuation of prior Sovaldi therapy. For pediatric patients 12 years of age and older or weighing at least 45 kg, using Sovaldi plus ribavirin with either GT 2 or GT3: TF/I/C to Mavyret, OR for continuation of prior Sovaldi therapy. (continued in Other Criteria) |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with one of the following: Hepatologist, Gastroenterologist, Infectious disease specialist, HIV specialist certified through the American Academy of HIV Medicine. |
| Coverage Duration | 12 to 48 weeks. Criteria will be applied consistent with current AASLD/IDSA guideline. |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|-----------------------|--|
| Other Criteria | All Sovaldi plus Daklinza therapy: One of the following: 1) Patient has not failed a prior HCV NS5A-containing regimen (eg, Daklinza) therapy, OR 2) patient has failed prior therapy with an NS5A-containing regimen AND patient does not have NS5A inhibitor resistance-associated variants detected using commercially available assays. For GT2 and GT3 (except liver tx patients) patients, using Sovaldi plus Daklinza: TF/I/C to sofosbuvir/velpatasvir AND Mavyret, OR for continuation of prior Sovaldi therapy. For GT2 or 3 liver tx recipients without cirrhosis, ONE of the following: 1) Patient has had a TF/I/C to Mavyret OR 2) For continuation of prior Sovaldi therapy. For GT2 or 3 liver tx recipients with decompensated cirrhosis, ONE of the following: 1) Patient has had a TF/I/C to sofosbuvir/velpatasvir OR 2) For continuation of prior Sovaldi therapy. For GT1 liver tx patients using Sovaldi plus Daklinza, TF/I/C to a) Mavyret OR b) continuation of prior Sovaldi therapy. |
|-----------------------|--|

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

SPINRAZA (S)

Products Affected

- Spinraza

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Spinal muscular atrophy (SMA) (initial): Diagnosis of spinal muscular atrophy (SMA) type I, II, or III. Both of the following: a) The mutation or deletion of genes in chromosome 5q resulting in one of the following: 1) Homozygous gene deletion or mutation (e.g., homozygous deletion of exon 7 at locus 5q13) or 2) Compound heterozygous mutation (e.g., deletion of SMN1 exon 7 [allele 1] and mutation of SMN1 [allele 2]) AND b) Patient has at least 2 copies of SMN2. Patient is not dependent on either of the following: 1) Invasive ventilation or tracheostomy or 2) Use of non-invasive ventilation beyond use for naps and nighttime sleep. At least one of the following exams (based on patient age and motor ability) has been conducted to establish baseline motor ability: Hammersmith Infant Neurological Exam (HINE) (infant to early childhood), Hammersmith Functional Motor Scale Expanded (HFMSSE), Upper Limb Module (ULM) Test (Non ambulatory), or Children's Hospital of Philadelphia Infant Test of Neuromuscular Disorders (CHOP INTEND). Spinraza is to be administered intrathecally by, or under the direction of, healthcare professionals experienced in performing lumbar punctures. |
| Age Restrictions | N/A |
| Prescriber Restrictions | SMA (initial, reauth): Prescribed by or in consultation with a neurologist with expertise in the diagnosis of SMA |
| Coverage Duration | Initial: 3 months. Reauth: 6 months. |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|-----------------------|--|
| Other Criteria | <p>SMA (reauth): Documentation of positive clinical response to Spinraza therapy from pretreatment baseline status as demonstrated by the most recent results (less than 1 month prior to request) from one of the following exams: A) Both of the following HINE milestones: 1) One of the following: a) Improvement or maintenance of a previous improvement of at least a 2 point (or maximal score) increase in ability to kick or b) Improvement or maintenance of a previous improvement of at least a 1 point increase in any other HINE milestone (e.g., head control, rolling, sitting, crawling, etc.), excluding voluntary grasp AND 2) One of the following: a) Improvement or maintenance of a previous improvement in more HINE motor milestones than worsening from pretreatment baseline (net positive improvement) or b) Patient has achieved and maintained any new motor milestones from pretreatment baseline when they would otherwise be unexpected to do so (e.g., sit unassisted, stand, walk) OR B) One of the following HFMSE milestones: 1) Improvement or maintenance of a previous improvement of at least a 3 point increase in score from pretreatment baseline OR 2) Patient has achieved and maintained any new motor milestone from pretreatment baseline when they would otherwise be unexpected to do so OR C) One of the following ULM test milestones: 1) Improvement or maintenance of a previous improvement of at least a 2 point increase in score from pretreatment baseline OR 2) Patient has achieved and maintained any new motor milestone from pretreatment baseline when they would otherwise be unexpected to do so OR D) One of the following CHOP INTEND milestones: 1) Improvement or maintenance of a previous improvement of at least a 4 point increase in score from pretreatment baseline OR 2) Patient has achieved and maintained any new motor milestone from pretreatment baseline when they would otherwise be unexpected to do so. Patient continues to not be dependent on either of the following: 1) Invasive ventilation or tracheostomy or 2) use of non-invasive ventilation beyond use for naps and nighttime sleep. Spinraza is to be administered intrathecally by, or under the direction of, healthcare professionals experienced in performing lumbar punctures.</p> |
|-----------------------|--|

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

SPORANOX (S)

Products Affected

- Itraconazole CAPS

- Itraconazole SOLN

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | One of the following: 1) patient has a systemic fungal infection (e.g., aspergillosis, histoplasmosis, blastomycosis), OR 2) all of the following: a) patient has a diagnosis of onychomycosis confirmed by one of the following (CAPSULE ONLY): i) positive potassium hydroxide (KOH) preparation, OR ii) fungal culture, OR iii) nail biopsy, AND b) patient has had a trial and failure, contraindication, or intolerance to oral terbinafine, OR 3) patient has a diagnosis of candidiasis (esophageal or oropharyngeal) that is refractory to treatment with fluconazole (ORAL SOLUTION ONLY). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | Systemic fungal infxn:6mo.Candidiasis:1mo.Fingernail onycho:5wks.Toenail onycho, other:3mo. |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

SPRAVATO (S)

Products Affected

- Spravato 56mg Dose
- Spravato 84mg Dose

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Treatment-resistant depression (TRD): Diagnosis of major depressive disorder (treatment-resistant). Patient has not experienced a clinical meaningful improvement after treatment with at least two antidepressants from different classes for an adequate duration (at least 4 weeks each) in the current depressive episode. Used in combination with an oral antidepressant. |
| Age Restrictions | N/A |
| Prescriber Restrictions | TRD: Prescribed by or in consultation with a psychiatrist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

SPRYCEL (S)

Products Affected

- Sprycel

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Philadelphia chromosome positive (Ph+)/BCR ABL chronic myelogenous leukemia (CML): Diagnosis of Ph+/BCR ABL CML. Ph+/BCR ABL acute lymphoblastic leukemia (ALL): Diagnosis of Ph+/BCR ABL ALL. |
| Age Restrictions | N/A |
| Prescriber Restrictions | All Uses: Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | All Uses: 12 months |
| Other Criteria | All Uses: Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

STELARA (IV) (S)

Products Affected

- Stelara INJ 130MG/26ML

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Diagnosis of moderately to severely active Crohn's disease. One of the following: a) trial and failure, contraindication, or intolerance to Humira (adalimumab), or (b) trial and failure, contraindication, or intolerance to treatment with at least one immunomodulator or corticosteroid [e.g., Purinethol (6-mercaptopurine), Imuran (azathioprine), Sandimmune (cyclosporine A), Prograf (tacrolimus), MTX (methotrexate)]. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a gastroenterologist. |
| Coverage Duration | One time |
| Other Criteria | Stelara is to be administered as an intravenous induction dose. Stelara induction dosing is in accordance with the United States Food and Drug Administration approved labeled dosing for Crohn's Disease: 260 mg for patients weighing 55 kg or less, 390 mg for patients weighing more than 55 kg to 85 kg, or 520 mg for patients weighing more than 85 kg. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

STELARA (S)

Products Affected

- Stelara INJ 45MG/0.5ML, 90MG/ML

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | <p>Plaque psoriasis (Initial - 45mg/0.5mL): Diagnosis of moderate to severe plaque psoriasis. Plaque psoriasis (Initial - 90mg/1mL): Diagnosis of moderate to severe plaque psoriasis. Patient's weight is greater than 100 kg (220 lbs). Psoriatic arthritis (PsA) (Initial - 45mg/0.5mL): Diagnosis of active PsA. PsA (Initial - 90mg/1mL): Diagnosis of active PsA. Patient's weight is greater than 100 kg (220 lbs). Diagnosis of co-existent moderate to severe psoriasis. Plaque psoriasis (Initial): One of the following: a) Trial and failure, contraindication, or intolerance (TF/C/I) to Cosentyx (secukinumab) AND one of the following: Enbrel (etanercept) OR Humira (adalimumab) OR b) for continuation of prior Stelara therapy. PsA (Initial): One of the following: a) TF/C/I to Cosentyx (secukinumab) AND one of the following: Enbrel (etanercept) or Humira (adalimumab), OR b) for continuation of prior Stelara therapy. Crohn's disease (CD) (Initial): Diagnosis of moderately to severely active Crohn's disease. One of the following: a) TF/C/I to Humira (adalimumab) OR b) TF/C/I to treatment with at least one immunomodulator or corticosteroid [e.g., Purinethol (6-mercaptopurine), Imuran (azathioprine), Sandimmune (cyclosporine A), Prograf (tacrolimus), MTX (methotrexate)], OR c) for continuation of prior Stelara therapy. Ulcerative colitis (UC) (Initial): Diagnosis of moderately to severely active UC. One of the following: a) TF/C/I to Humira (adalimumab) OR b) TF/C/I to treatment with at least one immunomodulator or corticosteroid (e.g., Purinethol [6-mercaptopurine], Imuran [azathioprine], aminosalicylates [e.g., mesalamine {Asacol, Pentasa, Rowasa}, olsalazine {Dipentum}, sulfasalazine {Azulfidine, Sulfazine}]), OR c) for continuation of prior Stelara therapy.</p> |
| Age Restrictions | N/A |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|--------------------------------|---|
| Prescriber Restrictions | Plaque psoriasis (initial): Prescribed by or in consultation with a dermatologist. PsA (initial): Prescribed by or in consultation with a dermatologist or rheumatologist. CD and UC (initial): Prescribed by or in consultation with a gastroenterologist. |
| Coverage Duration | All uses (Initial, reauth): 12 months |
| Other Criteria | Reauth (all indications): Documentation of positive clinical response to Stelara therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

STIVARGA (S)

Products Affected

- Stivarga

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Metastatic colorectal cancer (mCRC): All of the following: 1) diagnosis of mCRC, AND 2) trial and failure, contraindication or intolerance to fluoropyrimidine-, oxaliplatin- and irinotecan-based chemotherapy, AND 3) trial and failure, contraindication or intolerance to an anti-VEGF therapy (e.g., Avastin [bevacizumab]), AND 4) one of the following: a) RAS mutation, OR b) both of the following: RAS wild-type (RAS mutation negative tumor) and trial and failure, contraindication or intolerance to an anti-EGFR therapy [e.g., Vectibix (panitumumab), Erbitux (cetuximab)]. Gastrointestinal stromal tumor (GIST): All of the following: 1) diagnosis of locally advanced, unresectable or metastatic GIST, AND 2) trial and failure, contraindication or intolerance to both of the following: a) Gleevec (imatinib mesylate), AND b) Sutent (sunitinib malate). Hepatocellular Carcinoma (HCC): Diagnosis of HCC. Trial and failure or intolerance to Nexavar (sorafenib tosylate). |
| Age Restrictions | N/A |
| Prescriber Restrictions | mCRC, GIST: Prescribed by or in consultation with an oncologist. HCC: Prescribed by or in consultation with an oncologist, hepatologist, or gastroenterologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

STRENSIQ (S)

Products Affected

- Strensiq

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Hypophosphatasia: Diagnosis of perinatal/infantile or juvenile-onset hypophosphatasia AND for patients requesting the 80 mg/0.8 mL vial only: Patient's weight is greater than or equal to 40 kg. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Hypophosphatasia: Prescribed by or in consultation with a specialist experienced in the treatment of inborn errors of metabolism or endocrinologist |
| Coverage Duration | Hypophosphatasia: 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

SUPPRELIN LA (S)

Products Affected

- Supprelin La

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Central Precocious Puberty (CPP) (initial): Diagnosis of CPP (idiopathic or neurogenic). Early onset of secondary sexual characteristics in females less than age 8 or males less than age 9. Advanced bone age of at least one year compared with chronologic age. One of the following: a) patient has undergone gonadotropin-releasing hormone agonist (GnRHa) testing AND Peak luteinizing hormone (LH) level above pre-pubertal range, or b) patient has a random LH level in the pubertal range. |
| Age Restrictions | N/A |
| Prescriber Restrictions | CPP (initial, reauth): Prescribed by or in consultation with a pediatric endocrinologist. |
| Coverage Duration | CPP (init, reauth): 12 months |
| Other Criteria | CPP (reauthorization): LH levels have been suppressed to pre-pubertal levels. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

SUTENT (S)

Products Affected

- Sutent

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Renal cell carcinoma: Diagnosis of advanced or metastatic renal cell carcinoma. Gastrointestinal stromal tumor (GIST): Diagnosis of GIST after disease progression on, or contraindication or intolerance to Gleevec (imatinib). Pancreatic neuroendocrine tumors: Diagnosis of progressive, well-differentiated pancreatic neuroendocrine tumor that is unresectable locally advanced or metastatic disease. Adjuvant treatment of renal cell carcinoma: Diagnosis of renal cell carcinoma (RCC). Used as adjuvant therapy. Patient is at high risk of recurrent RCC following nephrectomy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | All uses: Prescribed by or in consultation with an oncologist |
| Coverage Duration | All uses: 12 months |
| Other Criteria | All Indications: Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

SYLATRON (S)

Products Affected

- Sylatron

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Melanoma: Diagnosis of melanoma with microscopic or gross nodal involvement AND The prescribed medication will be used as adjuvant therapy within 84 days of definitive surgical resection, including complete lymphadenectomy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or dermatologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

SYLVANT (S)

Products Affected

- Sylvant

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Multicentric Castleman's disease (MCD) (Initial): Diagnosis of MCD. Patient is human immunodeficiency virus (HIV) negative and human herpesvirus-8 (HHV-8) negative. |
| Age Restrictions | N/A |
| Prescriber Restrictions | MCD (initial): Prescribed by or in consultation with hematologist/oncologist or rheumatologist. |
| Coverage Duration | MCD (initial, reauth): 6 months |
| Other Criteria | MCD (reauth): Documentation of positive clinical response to Sylvant therapy. Patient is HIV negative and HHV-8 negative. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

SYMDEKO (S)

Products Affected

- Symdeko

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Initial: Diagnosis of cystic fibrosis. One of the following: 1) Patient is homozygous for the F508del mutation as detected by a U.S. Food and Drug Administration (FDA)-cleared cystic fibrosis mutation test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) OR 2) Patient has one of the following mutations on at least one allele in the CF transmembrane conductance regulator (CFTR) gene as detected by a U.S. Food and Drug Administration (FDA)-cleared cystic fibrosis mutation test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA): E56K, P67L, R74W, D110E, D110H, R117C, E193K, L206W, R347H, R352Q, A455E, D579G, 711+3A-G, E831X, S945L, S977F, F1052V, K1060T, A1067T, R1070W, F1074L, D1152H, D1270N, 2789+5G-A, 3272-26A-G, 3849+10kbC-T. |
| Age Restrictions | Initial: Patient is 6 years of age or older |
| Prescriber Restrictions | Initial: Prescribed by or in consultation with a pulmonologist or specialist affiliated with a CF care center |
| Coverage Duration | 12 months |
| Other Criteria | Reauth: Documentation of a positive clinical response to Symdeko (tezacaftor/ivacaftor) therapy (e.g., improvement in lung function or decreased number of pulmonary exacerbations). |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

SYMLIN (S)

Products Affected

- Symlinpen 120
- Symlinpen 60

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Initial: One of the following diagnoses: A) Type 1 diabetes OR B) Type 2 diabetes. Patient has failed to achieve desired glucose control despite optimal insulin therapy. Patient is taking concurrent mealtime insulin therapy (e.g., Humulin, Humalog, Novolin, Novolog). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | Reauth: Patient has experienced an objective response to therapy demonstrated by an improvement in HbA1c from baseline. Patient is receiving concurrent mealtime insulin therapy (e.g., Humulin, Humalog, Novolin, Novolog). |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

SYNAGIS (S)

Products Affected

- Synagis INJ 100MG/ML, 50MG/0.5ML

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | <p>Patient will use palivizumab for immunoprophylaxis of respiratory syncytial virus (RSV) during the peak months of infection in the patient's geographic region AND Patient meets one of the following criteria: 1) Infants born at 28 weeks, six days gestation or earlier and who are younger than 12 months of age at the start of the RSV season OR 2) Diagnosis of chronic lung disease of prematurity, born before 32 weeks, 0 days gestation, received greater than 21% oxygen for at least the first 28 days after birth, and one of the following: a) 12 months of age or younger at the start of the RSV season OR b) greater than 12 months of age to 24 months of age at the start of the RSV season and received medical support (i.e., chronic corticosteroid therapy, diuretic therapy, or supplemental oxygen) within 6 months before the start of the second RSV season. OR 3) Patient is 12 months of age or younger at the start of the RSV season and has one of the following: a) acyanotic heart failure that will require a cardiac surgical procedure and the patient is receiving medication to control congestive heart failure, OR b) moderate to severe pulmonary hypertension OR c) cyanotic heart defect. OR 4) patient is younger than 24 months of age and will or has undergone a cardiac transplantation during the RSV season. OR 5) Patient is 12 months of age or younger at the start of the RSV season with a congenital abnormality or neuromuscular disorder and has an impaired ability to clear secretions from the upper airway due to an ineffective cough. OR 6) Patient is younger than 24 months of age with a lymphocyte count below the normal range for patient's age and has received or will receive a solid organ transplant, hematopoietic stem cell transplant recipient, or chemotherapy during the RSV season.</p> |
| Age Restrictions | N/A |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|--------------------------------|--|
| Prescriber Restrictions | Prescribed by or in consultation with a pediatric specialist (i.e., pulmonologist, neonatologist, neurologist, cardiologist, pediatric intensivist, or infectious disease specialist). |
| Coverage Duration | 5 months (5 doses) during RSV season. |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

SYNDROS (S)

Products Affected

- Syndros

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Nausea and Vomiting Associated with Cancer Chemotherapy (CINV): Patient is receiving cancer chemotherapy. Trial and failure, contraindication, or intolerance (TF/C/I) to a 5HT-3 receptor antagonist (eg, Anzemet [dolasetron], Kytril [granisetron], or Zofran [ondansetron]). TF/C/I to one of the following: Compazine (prochlorperazine), Decadron (dexamethasone), Haldol (haloperidol), Zyprexa (olanzapine). AIDS anorexia: Diagnosis of anorexia with weight loss in patients with AIDS. Patient is on antiretroviral therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | CINV: 6 months. AIDS anorexia: 3 months. |
| Other Criteria | Subject to Part B vs. Part D review. CINV: Approve for continuation of therapy for treatment covered under Part B when patient is receiving chemotherapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

SYNRIBO (S)

Products Affected

- Synribo

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Chronic myelogenous leukemia (CML): Diagnosis of CML in the chronic or accelerated phase AND Patient has tried and has had resistance, relapse, inadequate response, intolerance or is contraindicated to TWO tyrosine kinase inhibitors (i.e., Gleevec [imatinib], Sprycel, Tassigna, and Bosulif, Iclusig). |
| Age Restrictions | CML: 18 years of age or older |
| Prescriber Restrictions | CML: Prescribed by or in consultation with a hematologist/oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

SYPRINE (S)

Products Affected

- Clovique
- Trientine Hydrochloride

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Initial: Diagnosis of Wilson's disease (i.e., hepatolenticular degeneration). Trial and failure, contraindication, or intolerance to a penicillamine product (e.g., Depen, Cuprimine) |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | Reauth: Documentation of a positive clinical response to therapy |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

TABRECTA (S)

Products Affected

- Tabrecta

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Diagnosis of non-small cell lung cancer (NSCLC). Disease is one of the following: recurrent, advanced, metastatic. Presence of mesenchymal-epithelial transition (MET) exon 14 skipping positive tumors as detected with an FDA-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

TAFAMIDIS (S)

Products Affected

- Vyndamax
- Vyndaqel

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Transthyretin-mediated amyloidosis with cardiomyopathy (ATTR-CM) (initial): Diagnosis of transthyretin-mediated amyloidosis with cardiomyopathy (ATTR-CM). One of the following: 1) Patient has a transthyretin (TTR) mutation (e.g., V122I), 2) Cardiac or noncardiac tissue biopsy demonstrating histologic confirmation of TTR amyloid deposits, OR 3) All of the following: i) echocardiogram or cardiac magnetic resonance imaging suggestive of amyloidosis, ii) scintigraphy scan suggestive of cardiac TTR amyloidosis, and iii) absence of light-chain amyloidosis. One of the following: 1) History of heart failure (HF), with at least one prior hospitalization for HF, OR 2) Presence of clinical signs and symptoms of HF (e.g., dyspnea, edema). Patient has New York Heart Association (NYHA) Functional Class I, II, or III heart failure. |
| Age Restrictions | N/A |
| Prescriber Restrictions | ATTR-CM (initial, reauth): Prescribed by or in consultation with a cardiologist |
| Coverage Duration | ATTR-CM (initial, reauth): 12 months |
| Other Criteria | ATTR-CM (reauth): Documentation of positive clinical response to therapy. Patient continues to have New York Heart Association (NYHA) Functional Class I, II, or III heart failure. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

TAFINLAR (S)

Products Affected

- Tafinlar

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | <p>Melanoma: Diagnosis of unresectable or metastatic melanoma AND cancer is BRAF V600E mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) OR both of the following: cancer is BRAF V600E or V600K mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) and medication is used in combination with Mekinist (trametinib). Adjuvant Treatment for Melanoma: Diagnosis of melanoma. Cancer is BRAF V600E or V600K mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Involvement of lymph nodes following complete resection. Used as adjunctive therapy. Medication is used in combination with Mekinist (trametinib). Non-small Cell Lung Cancer (NSCLC): Diagnosis of metastatic non-small cell lung cancer AND cancer is BRAF V600E mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) AND medication is used in combination with Mekinist (trametinib). Anaplastic Thyroid Cancer (ATC): Diagnosis of locally advanced or metastatic anaplastic thyroid cancer. Cancer is BRAF V600E mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Cancer may not be treated with standard locoregional treatment options. Medication is used in combination with Mekinist (trametinib) .</p> |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist. |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|--------------------------|--|
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

TAGRISO (S)

Products Affected

- Tagrisso

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Non-small cell lung cancer (NSCLC): Diagnosis of metastatic NSCLC. One of the following: 1) Patient has known active epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R mutations as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA), OR 2) Both of the following: a) Patient has known active EGFR T790M mutation as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) and b) Patient has experienced disease progression on or after one of the following EGFR Tyrosine Kinase Inhibitors (TKIs): Gilotrif (afatinib), Iressa (gefitinib), Tarceva (erlotinib). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

TAKHZYRO (S)

Products Affected

- Takhzyro

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Prophylaxis of hereditary angioedema (HAE) attacks: Diagnosis of HAE. For prophylaxis against HAE attacks. |
| Age Restrictions | N/A |
| Prescriber Restrictions | HAE (prophylaxis): Prescribed by or in consultation with an immunologist, allergist, or rheumatologist |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

TALTZ (S)

Products Affected

- Taltz

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Plaque psoriasis (Initial): Diagnosis of moderate to severe plaque psoriasis. One of the following: a) Trial and failure, contraindication, or intolerance to Cosentyx (secukinumab) AND one of the following: Enbrel (etanercept) or Humira (adalimumab), OR b) for continuation of prior Taltz therapy. Psoriatic Arthritis (PsA) (initial): Diagnosis of active PsA. One of the following: a) Trial and failure, contraindication, or intolerance to Cosentyx (secukinumab) AND one of the following: Enbrel (etanercept) or Humira (adalimumab), OR b) for continuation of prior Taltz therapy. Ankylosing spondylitis (AS) (initial): Diagnosis of active AS. One of the following: a) Trial and failure, contraindication, or intolerance to Cosentyx (secukinumab) AND one of the following: Enbrel (etanercept) or Humira (adalimumab), OR b) for continuation of prior Taltz therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Plaque Psoriasis (initial): Prescribed by or in consultation with a dermatologist. PsA (initial): Prescribed by or in consultation with a dermatologist or rheumatologist. AS (initial): Prescribed by or in consultation with a rheumatologist. |
| Coverage Duration | Initial, reauth: 12 months |
| Other Criteria | All indications (Reauth): Documentation of positive clinical response to Taltz therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

TALZENNA (S)

Products Affected

- Talzenna

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Breast cancer: Diagnosis of breast cancer. Disease is one of the following: a) locally advanced or b) metastatic. Presence of a deleterious or suspected deleterious germline BRCA-mutation (gBRCAm) as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Disease is human epidermal growth factor receptor 2 (HER2)-negative. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

TARCEVA (S)

Products Affected

- Erlotinib Hydrochloride

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Non-small cell lung cancer (NSCLC): Diagnosis of locally advanced or metastatic (Stage III or IV) NSCLC AND Patient has known active epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutation as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Pancreatic Cancer: Diagnosis of locally advanced, unresectable, or metastatic pancreatic cancer AND erlotinib will be used in combination with gemcitabine. |
| Age Restrictions | N/A |
| Prescriber Restrictions | All uses: Prescribed by or in consultation with an oncologist |
| Coverage Duration | All uses: 12 months |
| Other Criteria | All Indications: Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

TARGRETIN (S)

Products Affected

- Bexarotene
- Targretin GEL

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Cutaneous T-Cell Lymphoma (CTCL): Diagnosis of CTCL. Trial and failure, contraindication, or intolerance to at least one prior therapy (including skin-directed therapies [eg, corticosteroids {ie, clobetasol, diflorasone, halobetasol, augmented betamethasone dipropionate}] or systemic therapies [eg, interferons]). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or dermatologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

TASIGNA (S)

Products Affected

- Tasigna

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Chronic myelogenous leukemia (CML): Diagnosis of Ph+/BCR ABL CML |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

TAVALISSE (S)

Products Affected

- Tavalisse

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Chronic Idiopathic Thrombocytopenic Purpura (ITP) (initial): Diagnosis of chronic immune ITP or relapsed/refractory ITP. Baseline platelet count is less than 30,000/mcL. Trial and failure, contraindication, or intolerance to at least one of the following: corticosteroids, immunoglobulins, splenectomy, thrombopoietin receptor agonists (e.g., Nplate, Promacta), or Rituxan (rituximab). Patient's degree of thrombocytopenia and clinical condition increase the risk of bleeding. |
| Age Restrictions | N/A |
| Prescriber Restrictions | ITP (initial): Prescribed by or in consultation with a hematologist/oncologist. |
| Coverage Duration | ITP (initial, reauth): 12 months |
| Other Criteria | ITP (reauth): Documentation of positive clinical response to Tavalisse therapy as evidenced by an increase in platelet count to a level sufficient to avoid clinically important bleeding. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

TAZVERIK (S)

Products Affected

- Tazverik

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Epithelioid sarcoma: Diagnosis of epithelioid sarcoma. Disease is one of the following: metastatic or locally advanced. Patient is not eligible for complete resection. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

TECENTRIQ (S)

Products Affected

- Tecentriq

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | N/A |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 Months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

TECFIDERA (S)

Products Affected

- Tecfidera Starter Pack

- Tecfidera

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Multiple Sclerosis (MS): Diagnosis of a relapsing form of MS (eg, relapsing-remitting MS, secondary-progressive MS with relapses). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

TEGSEDI (S)

Products Affected

- Tegsedi

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Hereditary transthyretin-mediated amyloidosis (hATTR amyloidosis) (initial): Diagnosis of hATTR amyloidosis with polyneuropathy. Patient has a transthyretin (TTR) mutation (e.g., V30M). One of the following: 1) Patient has a baseline polyneuropathy disability (PND) score less than or equal to IIIb, 2) Patient has baseline familial amyloidotic polyneuropathy (FAP) stage of 1 or 2, OR 3) Patient has a baseline neuropathy impairment score (NIS) between 10 and 130. Presence of clinical signs and symptoms of the disease (e.g., peripheral/autonomic neuropathy). |
| Age Restrictions | N/A |
| Prescriber Restrictions | hATTR amyloidosis (initial): Prescribed by or in consultation with a neurologist |
| Coverage Duration | hATTR amyloidosis (initial, reauth): 12 months |
| Other Criteria | hATTR amyloidosis (reauth): Patient has demonstrated a benefit from therapy (e.g., improved neurologic impairment, slowing of disease progression, quality of life assessment). One of the following: 1) Patient continues to have a PND score less than or equal to IIIb, 2) Patient continues to have a FAP stage of 1 or 2, OR 3) Patient continues to have a NIS between 10 and 130. |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

TERIPARATIDE (S)

Products Affected

- Teriparatide

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Postmenopausal osteoporosis or osteopenia or men with primary or hypogonadal osteoporosis or osteopenia: Diagnosis of one of the following: a) postmenopausal osteoporosis or osteopenia or b) primary or hypogonadal osteoporosis or osteopenia. One of the following: Set I) Both of the following: A) Bone mineral density (BMD) T-score of -2.5 or lower in the lumbar spine, femoral neck, total hip, or radius (one-third radius site) AND B) One of the following: 1) history of low-trauma fracture of the hip, spine, proximal humerus, pelvis, or distal forearm, or 2) trial and failure, contraindication, or intolerance (TF/C/I) to one osteoporosis treatment (e.g., alendronate, risedronate, zoledronic acid, Prolia [denosumab]), or Set II) Both of the following: A) BMD T-score between -1.0 and -2.5 in the lumbar spine, femoral neck, total hip, or radius (one-third radius site) AND B) One of the following: 1) history of low-trauma fracture of the hip, spine, proximal humerus, pelvis, or distal forearm, or 2) both of the following: i) TF/C/I to one osteoporosis treatment (e.g., alendronate, risedronate, zoledronic acid, Prolia [denosumab]) and ii) One of the following FRAX 10-year probabilities: a) Major osteoporotic fracture at 20% or more in the U.S., or the country-specific threshold in other countries or regions, or b) Hip fracture at 3% or more in the U.S., or the country-specific threshold in other countries or regions. Treatment duration of parathyroid hormones [e.g., teriparatide, Tymlos (abaloparatide)] has not exceeded a total of 24 months during the patient's lifetime. Glucocorticoid-Induced Osteoporosis: See Other Criteria section. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|--------------------------|--|
| Coverage Duration | All uses: 24 months (max 24 months of therapy per lifetime) |
| Other Criteria | Glucocorticoid-Induced Osteoporosis: Diagnosis of glucocorticoid-induced osteoporosis. History of prednisone or its equivalent at a dose greater than or equal to 5mg/day for greater than or equal to 3 months. One of the following: 1) BMD T-score less than or equal to -2.5 based on BMD measurements from lumbar spine, femoral neck, total hip, or radius (one-third radius site), or 2) One of the following FRAX 10-year probabilities: a) Major osteoporotic fracture at 20% or more in the U.S., or the country-specific threshold in other countries or regions, or b) Hip fracture at 3% or more in the U.S., or the country-specific threshold in other countries or regions, or 3) History of one of the following fractures resulting from minimal trauma: vertebral compression fx, fx of the hip, fx of the distal radius, fx of the pelvis, or fx of the proximal humerus. TF/C/I to one bisphosphonate (e.g., alendronate). Treatment duration of parathyroid hormones [e.g., teriparatide, Tymlos (abaloparatide)] has not exceeded a total of 24 months during the patient's lifetime. |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

TESTOSTERONE (S)

Products Affected

- Androderm PT24 2MG/24HR, 4MG/24HR
- Striant
- Testosterone GEL 1.62%, 20.25MG/1.25GM, 25MG/2.5GM, 40.5MG/2.5GM, 50MG/5GM
- Testosterone SOLN
- Testosterone Cypionate INJ 100MG/ML, 200MG/ML
- Testosterone Pump
- Testosterone Topical Solution

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Hypogonadism (HG) (Initial): Diagnosis (dx) of HG AND male patient at birth AND one of the following: 1) Two pre-treatment serum total testosterone (T) levels less than 300 ng/dL (10.4 nmol/L) or less than the reference range for the lab OR 2) Both of the following: a) Has a condition that may cause altered sex-hormone binding globulin (SHBG) (eg, thyroid disorder, HIV disease, liver disorder, diabetes, obesity), and b) one pre-treatment calculated free or bioavailable T level less than 5 ng/dL (0.17 nmol/L) or less than reference range for the lab, OR 3) History of bilateral orchiectomy, panhypopituitarism, or a genetic disorder known to cause HG (eg, congenital anorchia, Klinefelter's syndrome). Gender Dysphoria (GD) (off-label): Dx of GD. Patient is a female-to-male transsexual. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | HG(init): (New to T tx:6 mo. New to plan and cont T tx:12 mo), (reauth): 12 mo. GD: 12 mo. |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|-----------------------|--|
| Other Criteria | HG (Reauth): 1) Follow-up total serum T level within or below the normal limits of the reporting lab, or 2) Follow-up total serum T level outside of upper limits of normal for the reporting lab and the dose is adjusted, OR 3) Has a condition that may cause altered SHBG (eg, thyroid disorder, HIV disease, liver disorder, diabetes, obesity), and one of the following: Follow-up calculated free or bioavailable T level within or below the normal limits of the reporting lab, or follow-up calculated free or bioavailable T level outside of upper limits of normal for the reporting lab and the dose is adjusted. |
|-----------------------|--|

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

TESTOSTERONE ENANTHATE (S)

Products Affected

- Testosterone Enanthate INJ

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Hypogonadism (HG) (Initial): Diagnosis (dx) of HG AND male patient at birth AND one of the following: 1) Two pre-treatment serum total testosterone (T) levels less than 300 ng/dL (10.4 nmol/L) or less than the reference range for the lab OR 2) Both of the following: a) Has a condition that may cause altered sex-hormone binding globulin (SHBG) (eg, thyroid disorder, HIV disease, liver disorder, diabetes, obesity), and b) one pre-treatment calculated free or bioavailable T level less than 5 ng/dL (0.17 nmol/L) or less than reference range for the lab, OR 3) History of bilateral orchiectomy, panhypopituitarism, or a genetic disorder known to cause HG (eg, congenital anorchia, Klinefelter's syndrome). Delayed puberty (DP): Dx of DP AND male patient at birth. Breast cancer (BC): Dx of inoperable BC AND used for palliative treatment AND female patient at birth. Gender Dysphoria (GD) (off-label): Dx of GD. Patient is a female-to-male transsexual. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | HG(init): (New to T tx:6 mo. Cont T tx:12 mo), (reauth): 12 mo. BC, GD: 12 mo. DP: 6 mo. |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|-----------------------|--|
| Other Criteria | HG (Reauth): 1) Follow-up total serum T level within or below the normal limits of the reporting lab, or 2) Follow-up total serum T level outside of upper limits of normal for the reporting lab and the dose is adjusted, OR 3) Has a condition that may cause altered SHBG (eg, thyroid disorder, HIV disease, liver disorder, diabetes, obesity), and one of the following: Follow-up calculated free or bioavailable T level within or below the normal limits of the reporting lab, or follow-up calculated free or bioavailable T level outside of upper limits of normal for the reporting lab and the dose is adjusted. |
|-----------------------|--|

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

THALOMID (S)

Products Affected

- Thalomid

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Multiple myeloma (MM): Diagnosis of MM. Used in combination with dexamethasone, unless the patient has an intolerance to steroids. Erythema nodosum leprosum (ENL): Diagnosis of moderate to severe ENL with cutaneous manifestations. Thalomid is not used as monotherapy if moderate to severe neuritis is present. |
| Age Restrictions | N/A |
| Prescriber Restrictions | MM: Prescribed by or in consultation with an oncologist/hematologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

TIBSOVO (S)

Products Affected

- Tibsovo

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Relapsed or refractory Acute Myeloid Leukemia (AML): Diagnosis of AML. Disease is relapsed or refractory. Patient has an isocitrate dehydrogenase-1 (IDH1) mutation as detected by a U.S. Food and Drug Administration (FDA)-approved test (e.g., Abbott RealTime IDH1 assay) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Newly-Diagnosed AML: Diagnosis of newly-diagnosed AML. Patient has an isocitrate dehydrogenase-1 (IDH1) mutation as detected by an FDA-approved test (e.g., Abbott RealTime IDH1 assay) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). One of the following: 1) patient is greater than or equal to 75 years old OR 2) patient has comorbidities that preclude use of intensive induction chemotherapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist/oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

TIGLUTIK (S)

Products Affected

- Tiglutik

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Amyotrophic Lateral Sclerosis (ALS): Diagnosis of ALS. Trial and failure or intolerance to generic riluzole tablets. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

TOLSURA (S)

Products Affected

- Tolsura

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Diagnosis of one of the following: Blastomycosis, Histoplasmosis, or Aspergillosis. Trial and failure or intolerance to generic itraconazole capsules. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 6 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

TOPICAL RETINOID (S)

Products Affected

- Avita
- Tretinoin CREA
- Tretinoin GEL
- Tretinoin Microsphere
- Tretinoin Microsphere Pump

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Acne vulgaris: Diagnosis of acne vulgaris (i.e., acne). |
| Age Restrictions | PA applies to members 26 years of age or older |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

TRACLEER (S)

Products Affected

- Bosentan

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH AND PAH is symptomatic AND One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH. |
| Age Restrictions | N/A |
| Prescriber Restrictions | PAH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist. |
| Coverage Duration | PAH (Initial): 6 months. PAH (Reauth): 12 months |
| Other Criteria | PAH (Reauth): Documentation of positive clinical response to therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

TRELSTAR (S)

Products Affected

- Trelstar
- Trelstar Mixject

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Prostate Cancer: Diagnosis of advanced or metastatic prostate cancer. Trial and failure, contraindication, or intolerance to any brand Lupron formulation. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

TREMFYA (S)

Products Affected

- Tremfya

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Plaque psoriasis (Initial): Diagnosis of moderate to severe plaque psoriasis. One of the following: set A) 1) Trial and failure, contraindication, or intolerance (TF/C/I) to Enbrel (etanercept) or Humira (adalimumab) AND 2) TF/C/I to Cosentyx (secukinumab), OR set B) for continuation of prior Tremfya therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Plaque psoriasis (initial): Prescribed by or in consultation with a dermatologist. |
| Coverage Duration | Plaque psoriasis (Initial, reauth): 12 months |
| Other Criteria | Plaque psoriasis (Reauth): Documentation of positive clinical response to Tremfya therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

TRIKAFTA (S)

Products Affected

- Trikafta

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Cystic Fibrosis (CF) (initial): Diagnosis of CF. Patient has at least one F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene as detected by an FDA-cleared cystic fibrosis mutation test or a test performed at a Clinical Laboratory Improvement Amendments (CLIA)-approved facility. |
| Age Restrictions | CF (initial): 12 years of age or older. |
| Prescriber Restrictions | CF (initial): Prescribed by or in consultation with a pulmonologist or specialist affiliated with a CF care center. |
| Coverage Duration | CF (initial, reauth): 12 months |
| Other Criteria | CF (reauth): Documentation of a positive clinical response to Trikafta therapy (e.g., improvement in lung function [percent predicted forced expiratory volume in one second {PPFEV1}] or decreased number of pulmonary exacerbations). |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

TRIPTODUR (S)

Products Affected

- Triptodur

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Central Precocious Puberty (CPP) (initial): Diagnosis of CPP (idiopathic or neurogenic). Early onset of secondary sexual characteristics in females less than age 8 or males less than age 9. Advanced bone age of at least one year compared with chronologic age. One of the following: a) patient has undergone gonadotropin-releasing hormone agonist (GnRHa) testing AND Peak luteinizing hormone (LH) level above pre-pubertal range, or b) patient has a random LH level in the pubertal range. Trial and failure or intolerance to Lupron Depot-Ped. |
| Age Restrictions | N/A |
| Prescriber Restrictions | CPP (initial): Prescribed by or in consultation with a pediatric endocrinologist. |
| Coverage Duration | CPP (Initial, reauth): 12 months |
| Other Criteria | CPP (reauthorization): LH levels have been suppressed to pre-pubertal levels. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

TUKYSA (S)

Products Affected

- Tukysa

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Breast cancer: Diagnosis of breast cancer. Disease is one of the following: a) advanced unresectable or b) metastatic. Disease is human epidermal growth factor receptor 2 (HER2)-positive. Used in combination with trastuzumab and capecitabine. Patient has received one or more prior anti-HER2 based regimens (e.g., trastuzumab, pertuzumab, ado-trastuzumab emtansine). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

TURALIO (S)

Products Affected

- Turalio

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Tenosynovial Giant Cell Tumor (TGCT): Diagnosis of TGCT. Patient is symptomatic. Patient is not a candidate for surgery due to worsening functional limitation or severe morbidity with surgical removal. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist/oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

TYKERB (S)

Products Affected

- Tykerb

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Breast Cancer: Diagnosis of human epidermal growth factor receptor 2 (HER2)-positive metastatic or recurrent breast cancer. Used in combination with one of the following: Trastuzumab, Xeloda (capecitabine), or aromatase inhibitors [eg, Aromasin (exemestane), Femara (letrozole), Arimidex (anastrozole)]. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

TYMLOS (S)

Products Affected

- Tymlos

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Diagnosis of postmenopausal osteoporosis or osteopenia. One of the following: Set I) Both of the following: A) Bone mineral density (BMD) T-score of -2.5 or lower in the lumbar spine, femoral neck, total hip, or radius (one-third radius site) AND B) One of the following: 1) history of low-trauma fracture of the hip, spine, proximal humerus, pelvis, or distal forearm, or 2) trial and failure, contraindication, or intolerance (TF/C/I) to one osteoporosis treatment (e.g., alendronate, risedronate, zoledronic acid, Prolia [denosumab]), or Set II) Both of the following: A) BMD T-score between -1.0 and -2.5 in the lumbar spine, femoral neck, total hip, or radius (one-third radius site) AND B) One of the following: 1) history of low-trauma fracture of the hip, spine, proximal humerus, pelvis, or distal forearm, or 2) both of the following: i) TF/C/I to one osteoporosis treatment (e.g., alendronate, risedronate, zoledronic acid, Prolia [denosumab]) and ii) one of the following FRAX (Fracture Risk Assessment Tool) 10-year probabilities: a) major osteoporotic fracture at 20% or more in the U.S., or the country-specific threshold in other countries or regions, or b) hip fracture at 3% or more in the U.S., or the country-specific threshold in other countries or regions. Treatment duration of parathyroid hormones (e.g., Forteo [teriparatide], Tymlos [abaloparatide]) has not exceeded a total of 24 months during the patient's lifetime. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 24 months (max 24 months of therapy per lifetime) |
| Other Criteria | N/A |

TYSABRI (S)

Products Affected

- Tysabri

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | <p>Multiple Sclerosis (MS): Diagnosis of a relapsing form of MS (eg, relapsing-remitting MS, secondary-progressive MS with relapses). One of the following: 1) Trial and failure, contraindication, or intolerance (TF/C/I) to one of the following disease-modifying therapies for MS: Aubagio (teriflunomide), Avonex (interferon beta-1a), Betaseron (interferon beta-1b), Copaxone/Glatopa (glatiramer acetate), Extavia (interferon beta-1b), Gilenya (fingolimod), Lemtrada (alemtuzumab), Mavenclad (cladribine), Mayzent (siponimod), Ocrevus (ocrelizumab), Plegridy (peginterferon beta-1a), Rebif (interferon beta-1a), Tecfidera (dimethyl fumarate), or Zinbryta (daclizumab), 2) Patient is not a candidate for any of the drugs listed as prerequisites due to the severity of their MS, or 3) for continuation of prior therapy. Not used in combination with another disease-modifying therapy for MS. Crohn's Disease (CD) (initial): Diagnosis of moderate to severe CD with evidence of inflammation (eg, elevated C-reactive protein [CRP], elevated erythrocyte sedimentation rate, presence of fecal leukocytes). TF/C/I to one of the following conventional therapies: corticosteroids, 6-mercaptopurine (6MP [Purinethol], azathioprine (Imuran), methotrexate, aminosalicylates (eg, sulfasalazine, mesalamine, olsalazine). TF/C/I to a TNF-inhibitor (eg, Cimzia [certolizumab pegol], Humira [adalimumab], Remicade [infliximab]). CD (initial and reauth): Not used in combination with an immunosuppressant (eg, 6-MP, azathioprine, cyclosporine, or methotrexate). Not used in combination with a TNF-inhibitor (eg, Enbrel [etanercept], Humira [adalimumab], or Remicade [infliximab]).</p> |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|--------------------------|--|
| Coverage Duration | MS: 12mo. CD (Init): 3 mo. CD (Reauth): 6 mo if not on steroids. Otherwise, 3 mo. |
| Other Criteria | CD (reauth): Documentation of positive clinical response (eg, improved disease activity index) to Tysabri therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

UBRELVY (S)

Products Affected

- Ubrelyvy

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Initial: Diagnosis of migraine with or without aura. Will be used for the acute treatment of migraine. Will not be used for preventive treatment of migraine. Patient has less than 15 headache days per month. Trial and failure or intolerance to one triptan (e.g., eletriptan, rizatriptan, sumatriptan) or a contraindication to all triptans. If patient has 4 or more headache days per month, patient must meet one of the following: a) currently being treated with Elavil (amitriptyline) or Effexor (venlafaxine) unless there is a contraindication or intolerance to these medications, OR b) currently being treated with Depakote/Depakote ER (divalproex sodium) or Topamax (topiramate) unless there is a contraindication or intolerance to these medications, OR c) currently being treated with a beta blocker (i.e., atenolol, propranolol, nadolol, timolol, or metoprolol) unless there is a contraindication or intolerance to these medications. |
| Age Restrictions | Initial: 18 years of age or older. |
| Prescriber Restrictions | Initial, Reauth: Prescribed by or in consultation with a neurologist or pain specialist. |
| Coverage Duration | Initial: 3 months. Reauth: 12 months. |
| Other Criteria | Reauth: Patient has experienced a positive response to therapy (e.g., reduction in pain, photophobia, phonophobia, nausea). Will not be used for preventive treatment of migraine. |

UDENYCA (S)

Products Affected

- Udenyca

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | <p>Febrile neutropenia (FN) prophylaxis: Patient will be receiving prophylaxis for FN due to one of the following: 1) Patient is receiving National Cancer Institute’s Breast Intergroup, INT C9741 dose dense chemotherapy protocol for primary breast cancer, 2) patient is receiving a dose-dense chemotherapy regimen for which the incidence of FN is unknown, 3) patient is receiving chemotherapy regimen(s) associated with greater than 20% incidence of FN, 4) both of the following: a) patient is receiving chemotherapy regimen(s) associated with 10-20% incidence of FN, AND b) patient has one or more risk factors associated with chemotherapy-induced infection, FN, or neutropenia, OR 5) Both of the following: a) patient is receiving myelosuppressive anticancer drugs associated with neutropenia, AND b) patient has a history of FN or dose-limiting event during a previous course of chemotherapy (secondary prophylaxis). Treatment of FN (off-label): Patient has received or is receiving myelosuppressive anticancer drugs associated with neutropenia. Diagnosis of FN. Patient is at high risk for infection-associated complications. Acute radiation syndrome (ARS) (off-label): Patient was/will be acutely exposed to myelosuppressive doses of radiation (hematopoietic subsyndrome of ARS).</p> |
| Age Restrictions | N/A |
| Prescriber Restrictions | All uses: Prescribed by or in consultation with a hematologist/oncologist |
| Coverage Duration | ARS: 1 mo. FN (prophylaxis, treatment): 3 mo or duration of tx. |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ULTOMIRIS (S)

Products Affected

- Ultomiris

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Paroxysmal nocturnal hemoglobinuria (PNH) (initial): Diagnosis of PNH. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | PNH (initial, reauth): 12 months |
| Other Criteria | PNH (reauth): Documentation of positive clinical response (e.g., hemoglobin stabilization, decrease in the number of red blood cell transfusions) to Ultomiris therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

UPTRAVI (S)

Products Affected

- Uptravi

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Pulmonary arterial hypertension (PAH) (initial): Diagnosis of PAH AND Patient is symptomatic AND One of the following: a) Diagnosis of PAH was confirmed by right heart catheterization OR b) patient is currently on any therapy for the diagnosis of PAH. One of the following: a) trial and failure, contraindication, or intolerance to a PDE5 inhibitor [i.e., Adcirca (tadalafil), Revatio (sildenafil)] or Adempas (riociguat), and trial and failure, contraindication, or intolerance to an endothelin receptor antagonist [e.g. Letairis (ambrisentan), Opsumit (macitentan), or Tracleer (bosentan)] OR b) For continuation of prior Uptravi therapy. Not taken in combination with a prostanoid/prostacyclin analogue [e.g., Flolan (epoprostenol), Ventavis (iloprost), Tyvaso/Remodulin/Orenitram (treprostinil)]. |
| Age Restrictions | N/A |
| Prescriber Restrictions | PAH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist. |
| Coverage Duration | Initial: 6 months. Reauth: 12 months |
| Other Criteria | PAH (Reauth): Documentation of positive clinical response to Uptravi therapy. Not taken in combination with a prostanoid/prostacyclin analogue [e.g., Flolan (epoprostenol), Ventavis (iloprost), Tyvaso/Remodulin/Orenitram (treprostinil)]. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

VALCHLOR (S)

Products Affected

- Valchlor

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Mycosis fungoides-type cutaneous T-cell lymphoma (MF-CTCL) (initial): All of the following: 1) diagnosis of Stage IA MF-CTCL, OR diagnosis of Stage IB MF-CTCL, AND 2) patient has received at least one prior skin-directed therapy [e.g., topical corticosteroids, bexarotene topical gel (Targretin topical gel), etc.]. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or dermatologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

VARIZIG (S)

Products Affected

- Varizig INJ 125UNIT/1.2ML

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | Presence of contraindications to immune globulin therapy (i.e., IgA deficiency with antibodies to IgA and a history of hypersensitivity or product specific contraindication). |
| Required Medical Information | Immune globulin is being used intramuscularly. The immune globulin is being used for passive immunization or post exposure-prophylaxis of varicella. Patient is considered a high risk individual (i.e., immune compromised, pregnant woman, newborn of mother with varicella, premature infant, and infant less than 1 year old). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 3 months (approve one dose only) |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

VELCADE (S)

Products Affected

- Velcade

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Multiple myeloma (MM): Diagnosis of MM. Mantle cell lymphoma (MCL): Diagnosis of MCL. |
| Age Restrictions | N/A |
| Prescriber Restrictions | MM, MCL: Prescribed by or in consultation with an oncologist/hematologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

VENCLEXTA (S)

Products Affected

- Venclexta

- Venclexta Starting Pack

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Chronic lymphocytic leukemia (CLL)/Small lymphocytic lymphoma (SLL): Diagnosis of CLL or SLL. Acute Myeloid Leukemia (AML): Diagnosis of newly diagnosed AML. Used in combination with azacitidine, or decitabine, or low-dose cytarabine. One of the following: 1) age 75 years or older OR 2) comorbidities that preclude use of intensive induction chemotherapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist or oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

VENTAVIS (S)

Products Affected

- Ventavis

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Pulmonary arterial hypertension (PAH) (Initial): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH. |
| Age Restrictions | N/A |
| Prescriber Restrictions | PAH (initial): Prescribed by or in consultation with a pulmonologist or cardiologist. |
| Coverage Duration | PAH (Initial): 6 months. (Reauth): 12 months |
| Other Criteria | Subject to Part B vs D review. PAH (Reauth): Documentation of positive clinical response to therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

VERZENIO (S)

Products Affected

- Verzenio

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Breast Cancer: Diagnosis of advanced or metastatic breast cancer. Disease is hormone receptor (HR)-positive and human epidermal growth factor receptor 2 (HER2)-negative. One of the following: a) used in combination with an aromatase inhibitor (e.g., Arimidex [anastrozole], Aromasin [exemestane], Femara [letrozole]) and patient is a postmenopausal woman, OR b) used in combination with Faslodex (fulvestrant) and disease has progressed following endocrine therapy, OR c) used as monotherapy and disease has progressed following endocrine therapy and patient has already received at least one prior chemotherapy regimen. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

VIMIZIM (S)

Products Affected

- Vimizim

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Mucopolysaccharidosis (initial): Diagnosis of Mucopolysaccharidosis type IVA (MPS IVA, Morquio A syndrome) confirmed by both of the following: a) documented clinical signs and symptoms of the disease (e.g., kyphoscoliosis, genu valgum, pectus carinatum, gait disturbance, growth deficiency, etc.) and b) documented reduced fibroblast or leukocyte GALNS enzyme activity or molecular genetic testing of GALNS. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | Initial, reauth: 12 months |
| Other Criteria | Reauth: Documentation of positive clinical response to Vimizim therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

VIMOVO (S)

Products Affected

- Naproxen/esomeprazole Magnesium

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | (initial): All of the following: 1) diagnosis of one of the following: a) osteoarthritis, OR b) rheumatoid arthritis, OR c) ankylosing spondylitis, OR d) juvenile idiopathic arthritis, AND 2) one of the following: a) history of peptic ulcer disease, OR b) history of gastrointestinal (GI) bleeding, obstruction, or perforation, OR c) erosive esophagitis, OR d) used in combination with aspirin, AND 3) trial and failure, or intolerance to esomeprazole and naproxen sodium administered together as separate components. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | All uses (initial, reauth): 3 months. |
| Other Criteria | All uses (reauth): documentation of positive clinical response to Vimovo therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

VITRAKVI (S)

Products Affected

- Vitrakvi

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Presence of solid tumors (e.g., salivary gland, soft tissue sarcoma, infantile fibrosarcoma, thyroid cancer, lung, melanoma, colon, etc.). Disease is positive for neurotrophic receptor tyrosine kinase (NTRK) gene fusion (e.g. ETV6-NTRK3, TPM3-NTRK1, LMNA-NTRK1, etc.). Disease is without a known acquired resistance mutation [e.g., TRKA G595R substitution, TRKA G667C substitution, or other recurrent kinase domain (solvent front and xDFG) mutations]. Disease is one of the following: metastatic or unresectable (including cases where surgical resection is likely to result in severe morbidity). One of the following: Disease has progressed on previous treatment (e.g., surgery, radiotherapy, or systemic therapy) OR Disease has no satisfactory alternative treatments. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

VIZIMPRO (S)

Products Affected

- Vizimpro

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Non-small cell lung cancer (NSCLC): Diagnosis of NSCLC. Disease is metastatic. Disease is positive for one of the following epidermal growth factor receptor (EGFR) mutations: exon 19 deletion or exon 21 L858R substitution. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

VOSEVI (S)

Products Affected

- Vosevi

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Criteria will be applied consistent with current AASLD/IDSA guideline. All patients: Diagnosis of chronic hepatitis C, patient is without decompensated liver disease (defined as Child-Pugh Class B or C), and patient is not receiving Vosevi in combination with another HCV direct acting antiviral agent [e.g., Harvoni, Zepatier]. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with one of the following: Hepatologist, Gastroenterologist, Infectious disease specialist, HIV specialist certified through the American Academy of HIV Medicine. |
| Coverage Duration | 12 weeks. Criteria will be applied consistent with current AASLD/IDSA guideline. |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

VOTRIENT (S)

Products Affected

- Votrient

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Renal cell carcinoma (RCC): Diagnosis of advanced/metastatic RCC. Soft tissue sarcoma: Diagnosis of advanced soft tissue sarcoma and patient received at least one prior chemotherapy (e.g., ifosfamide, doxorubicin, cisplatin, dacarbazine, docetaxel, oxaliplatin, etc.). |
| Age Restrictions | N/A |
| Prescriber Restrictions | All uses: Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

VPRIV (S)

Products Affected

- Vpriv

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Gaucher disease: Diagnosis of type 1 Gaucher disease. Patient has evidence of symptomatic disease (e.g., moderate to severe anemia, thrombocytopenia, bone disease, hepatomegaly, or splenomegaly). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | Gaucher disease: 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

VUMERITY (S)

Products Affected

- Vumerity

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Multiple Sclerosis (MS): Diagnosis of a relapsing form of MS (e.g., relapsing-remitting MS, secondary-progressive MS with relapses). One of the following: a) Failure after a trial of at least 4 weeks, contraindication, or intolerance to two of the following disease-modifying therapies for MS: Aubagio (teriflunomide), Gilenya (fingolimod), or Tecfidera (dimethyl fumarate), OR b) for continuation of prior therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

VYXEOS (S)

Products Affected

- Vyxeos

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Newly diagnosed therapy related acute myeloid leukemia (t-AML): Diagnosis of t-AML. Acute myeloid leukemia myelodysplasia-related changes (AML-MRC): Diagnosis of AML-MRC. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist/oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

WAKIX (S)

Products Affected

- Wakix

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Narcolepsy (initial): Diagnosis (Dx) of narcolepsy as confirmed by sleep study (unless the prescriber provides justification confirming that a sleep study would not be feasible). Trial and failure, contraindication or intolerance to both generic modafinil and generic armodafinil. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | Initial, reauth: 12 months |
| Other Criteria | Narcolepsy (reauth): Documentation of positive clinical response to therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

XALKORI (S)

Products Affected

- Xalkori

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Non-small cell lung cancer (NSCLC): Diagnosis of advanced or metastatic NSCLC AND One of the following: A) Patient has an anaplastic lymphoma kinase (ALK)-positive tumor as detected with a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) or B) Patient has MET amplification- or ROS1 rearrangements-positive tumor as detected with an FDA-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). |
| Age Restrictions | N/A |
| Prescriber Restrictions | NSCLC: Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

XCOPRI (S)

Products Affected

- Xcopri

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Diagnosis of partial onset seizures. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

XELJANZ (S)

Products Affected

- Xeljanz
- Xeljanz Xr

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Rheumatoid arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. Psoriatic arthritis (PsA) (Initial): Diagnosis of active PsA. RA/PsA (initial): One of the following: Failure, contraindication, or intolerance to both Enbrel (etanercept) and Humira (adalimumab), or attestation demonstrating a trial may be inappropriate, OR patient has a documented needle-phobia to the degree that the patient has previously refused any injectable therapy or medical procedure (refer to DSM-IV-TR 300.29/F40.2 for specific phobia diagnostic criteria), OR for continuation of prior tofacitinib therapy. Ulcerative colitis (UC) (Initial): Diagnosis of moderately to severely active UC. Trial and failure, contraindication or intolerance to one of the following conventional therapies: 6-mercaptopurine (Purinethol), aminosalicylate [e.g., mesalamine (Asacol, Pentasa, Rowasa), olsalazine (Dipentum), sulfasalazine (Azulfidine, Sulfazine)], azathioprine (Imuran), or corticosteroids (e.g., prednisone, methylprednisolone). Trial and failure, contraindication, or intolerance to Humira (adalimumab), OR patient has a documented needle-phobia to the degree that the patient has previously refused any injectable therapy or medical procedure (refer to DSM-V-TR F40.2 for specific phobia diagnostic criteria), OR for continuation of prior tofacitinib therapy. All indications: Patient is not receiving tofacitinib in combination with a potent immunosuppressant (eg, azathioprine, cyclosporine). |
| Age Restrictions | N/A |
| Prescriber Restrictions | RA (initial): Prescribed by or in consultation with a rheumatologist. PsA (initial): Prescribed by or in consultation with a dermatologist or rheumatologist. UC (initial): Prescribed by or in consultation with a gastroenterologist. |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|--------------------------|---|
| Coverage Duration | RA/PsA (initial, reauth): 12 months. UC (init): 4 mo. UC (reauth): 12 mo. |
| Other Criteria | All Indications (Reauth): Documentation of positive clinical response to tofacitinib therapy. Patient is not receiving tofacitinib in combination with a potent immunosuppressant (eg, azathioprine, cyclosporine). |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

XENAZINE (S)

Products Affected

- Tetrabenazine

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Chorea associated with Huntington's Disease (HD) (Initial): Diagnosis of chorea in patients with Huntington's disease. Tardive dyskinesia (Initial): Diagnosis of tardive dyskinesia. One of the following: 1) Patient has persistent symptoms of tardive dyskinesia despite a trial of dose reduction, tapering, or discontinuation of the offending medication, OR 2) Patient is not a candidate for a trial of dose reduction, tapering, or discontinuation of the offending medication. Tourette's syndrome (Initial): Patient has tics associated with Tourette's syndrome. Failure, contraindication, or intolerance to Haldol (haloperidol). |
| Age Restrictions | N/A |
| Prescriber Restrictions | HD (Initial): Prescribed by or in consultation with a neurologist. Tardive dyskinesia, Tourette's syndrome (Initial): Prescribed by or in consultation with neurologist or psychiatrist. |
| Coverage Duration | All uses: (initial) 3 months. (Reauth) 12 months. |
| Other Criteria | All indications (Reauth): Documentation of clinical response and benefit from therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

XEOMIN (S)

Products Affected

- Xeomin

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Cervical Dystonia (CD) (init): Diagnosis of CD (also known as spasmodic torticollis). Blepharospasm (initial): Diagnosis of blepharospasm. History of previous use of Botox (onabotulinumtoxinA) for the treatment of blepharospasm. Upper limb spasticity (ULS) (init): Diagnosis of upper limb spasticity. Chronic Sialorrhea (CS) (init): Diagnosis of chronic sialorrhea. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | All indications (init, reauth): 3 months (for 1 dose) |
| Other Criteria | CD, blepharospasm, ULS (reauth): Confirmed improvement in symptoms with initial Xeomin treatment. At least 3 months have elapsed or will have elapsed since the last treatment with Xeomin. CS (reauth): Confirmed improvement in symptoms with initial Xeomin treatment. At least 4 months have elapsed or will have elapsed since the last treatment with Xeomin. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

XERMELO (S)

Products Affected

- Xermelo

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Carcinoid syndrome diarrhea (Initial): Diagnosis of carcinoid syndrome diarrhea AND diarrhea is inadequately controlled by a stable dose of somatostatin analog (SSA) therapy (e.g., octreotide [Sandostatin, Sandostatin LAR], lanreotide [Somatuline Depot]) for at least 3 months AND used in combination with SSA therapy. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Initial: Prescribed by or in consultation with an oncologist, endocrinologist, or gastroenterologist |
| Coverage Duration | Initial: 6 months. Reauth: 12 months |
| Other Criteria | Carcinoid syndrome diarrhea (Reauthorization): Documentation of a positive clinical response to Xermelo therapy AND Xermelo will continue to be used in combination with SSA therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

XGEVA (S)

Products Affected

- Xgeva

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Multiple Myeloma (MM)/Bone metastasis from solid tumors (BMST): One of the following: 1) Diagnosis of multiple myeloma OR 2) diagnosis of solid tumors (eg, breast cancer, kidney cancer, lung cancer, prostate cancer, thyroid cancer), AND documented evidence of one or more metastatic bone lesions. Giant cell tumor of bone (GCTB): Both of the following: 1) diagnosis of giant cell tumor of bone AND 2) One of the following: a) tumor is unresectable, OR b) surgical resection is likely to result in severe morbidity. Hypercalcemia of malignancy (HCM): Both of the following: 1) diagnosis of hypercalcemia of malignancy, AND 2) Trial and failure, contraindication, or intolerance to one intravenous bisphosphonate (eg, pamidronate, Zometa (zoledronic acid). |
| Age Restrictions | N/A |
| Prescriber Restrictions | GCTB, HCM: Prescribed by or in consultation with an oncologist |
| Coverage Duration | MM/BMST, GCTB: 12 mo. HCM: 2 mo. |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

XIAFLEX (S)

Products Affected

- Xiaflex

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Dupuytren's contracture (DC) (initial, reauth): Diagnosis of Dupuytren's contracture with a palpable cord AND Prescriber is enrolled in the Xiaflex REMS program for Dupuytren's contracture AND Patient has a positive "table top test" (defined as the inability to simultaneously place the affected finger and palm flat against a table top) AND Patient has a documented contracture of at least 20 degrees flexion for a metacarpophalangeal joint or a proximal interphalangeal joint AND Patient has a flexion deformity that results in functional limitations. Peyronie's disease (PD) (initial, reauth): Diagnosis of Peyronie's disease AND Prescriber is enrolled in the Xiaflex REMS program for Peyronie's disease AND Patient has a palpable plaque and curvature deformity of at least 30 degrees at the start of therapy AND The plaques do not involve the penile urethra AND Patient has a curvature deformity that results in pain (e.g., pain upon erection or intercourse). |
| Age Restrictions | Initial (DC, PD): 18 years of age or older |
| Prescriber Restrictions | N/A |
| Coverage Duration | DC, PD (Initial and reauth): 12 months |
| Other Criteria | Peyronie's disease (reauth): patient has a new plaque that results in a curvature deformity. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

XIFAXAN (S)

Products Affected

- Xifaxan

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Travelers' diarrhea (TD) (200 mg strength only): Diagnosis of travelers' diarrhea, AND one of the following: a) Trial and failure, contraindication, or intolerance to one of the following: Cipro (ciprofloxacin), Levaquin (levofloxacin), ofloxacin, Zithromax (azithromycin) OR b) resistance to all of the following: Cipro (ciprofloxacin), Levaquin (levofloxacin), ofloxacin, Zithromax (azithromycin). Prophylaxis of hepatic encephalopathy (HE) recurrence (550mg strength only): Used for the prophylaxis of hepatic encephalopathy recurrence, AND trial and failure, contraindication or intolerance to lactulose. Treatment of HE: Used for the treatment of HE. Trial and failure, contraindication, or intolerance to lactulose. Irritable bowel syndrome with diarrhea (IBS-D) (550mg strength only) (initial): Diagnosis of IBS-D, AND trial and failure, contraindication or intolerance to an antidiarrheal agent [eg, loperamide]. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | TD: 14 days. HE (prophylaxis, treatment): 12 months. IBS-D (initial, reauth): 2 weeks. |
| Other Criteria | IBS-D (reauth): Patient experiences IBS-D symptom recurrence. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

XOLAIR (S)

Products Affected

- Xolair

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Asthma (init): Diagnosis of moderate to severe persistent allergic asthma. Positive skin test or in vitro reactivity to a perennial aeroallergen. Pretreatment serum immunoglobulin (Ig)E level between 30 to 700 IU/mL for patients 12 years of age and older OR 30 to 1300 IU/mL for patients 6 years to less than 12 years of age. Patient is currently being treated with one of the following unless there is a contraindication or intolerance to these medications: a) Both of the following: i) High-dose inhaled corticosteroid (ICS) [e.g., greater than 500 mcg fluticasone propionate equivalent/day] and ii) additional asthma controller medication [e.g., leukotriene receptor antagonist, long-acting beta-2 agonist (LABA), theophylline], OR b) One maximally-dosed combination ICS/LABA product [e.g., Advair (fluticasone propionate/salmeterol), Dulera (mometasone/formoterol), Symbicort (budesonide/formoterol)]. Chronic Idiopathic Urticaria (CIU) (init): Diagnosis of CIU. Persistent symptoms (itching and hives) for at least 4 consecutive weeks despite titrating to an optimal dose with a second generation H1 antihistamine, unless there is a contraindication or intolerance to H1 antihistamines. Patient has tried and had an inadequate response or intolerance or contraindication to at least one of the following additional therapies: H2 antagonist, leukotriene receptor antagonist, H1 antihistamine, hydroxyzine, doxepin. Used concurrently with an H1 antihistamine, unless there is a contraindication or intolerance to H1 antihistamines. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Asthma (init/reauth): Prescribed by or in consultation with an allergist/immunologist, or pulmonologist. CIU (init): Prescribed by or in consultation with an allergist/immunologist, or dermatologist |
| Coverage Duration | Asthma (init): 6 months, Asthma (reauth): 12 months. CIU (init): 3 months, (reauth) 6 months |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|-----------------------|---|
| Other Criteria | <p>Asthma (reauth): Documentation of positive clinical response to therapy (e.g., Reduction in number of asthma exacerbations, improvement in forced expiratory volume in 1 second (FEV1), or decreased use of rescue medications). Patient is currently being treated with one of the following unless there is a contraindication or intolerance to these medications: a) Both of the following: i) inhaled corticosteroid (ICS) and ii) additional asthma controller medication [e.g., leukotriene receptor antagonist, long-acting beta-2 agonist (LABA), theophylline], OR b) A combination ICS/LABA product [e.g., Advair (fluticasone propionate/salmeterol), Dulera (mometasone/formoterol), Symbicort (budesonide/formoterol)].</p> <p>CIU (reauth): Patient's disease status has been re-evaluated since the last authorization to confirm the patient's condition warrants continued treatment. Patient has experienced one or both of the following: Reduction in itching severity from baseline or Reduction in the number of hives from baseline.</p> |
|-----------------------|---|

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

XOSPATA (S)

Products Affected

- Xospata

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Diagnosis of acute myeloid leukemia (AML). Disease is relapsed or refractory. Patient has a FMS-like tyrosine kinase (FLT3) mutation as determined by a U.S. Food and Drug Administration (FDA)-approved test (e.g., LeukoStrat CDx FLT3 Mutation Assay) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist or oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

XPOVIO (S)

Products Affected

- Xpovio 100 Mg Once Weekly
- Xpovio 40 Mg Once Weekly
- Xpovio 40 Mg Twice Weekly
- Xpovio 60 Mg Once Weekly
- Xpovio 60 Mg Twice Weekly
- Xpovio 80 Mg Once Weekly
- Xpovio 80 Mg Twice Weekly

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Diagnosis of relapsed or refractory multiple myeloma (RRMM). Patient has received at least four prior therapies. Disease is refractory to all of the following: 1) Two proteasome inhibitors (e.g., bortezomib, carfilzomib), 2) Two immunomodulatory agents (e.g., lenalidomide, thalidomide), and 3) An anti-CD38 monoclonal antibody (e.g. daratumumab). Used in combination with dexamethasone. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist/hematologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

XTANDI (S)

Products Affected

- Xtandi

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Castration-resistant or castration-recurrent prostate cancer (CRPC): Diagnosis of castration-resistant (chemical or surgical) or recurrent prostate cancer. Metastatic castration-sensitive prostate cancer (M-CSPC): Diagnosis of metastatic castration-sensitive prostate cancer. |
| Age Restrictions | N/A |
| Prescriber Restrictions | CRPC, M-CSPC: Prescribed by or in consultation with an oncologist or urologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

XURIDEN (S)

Products Affected

- Xuriden

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Hereditary orotic aciduria (Initial): Diagnosis of hereditary orotic aciduria. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Initial: Prescribed by or in consultation with a medical geneticist or other specialist that treats inborn errors of metabolism |
| Coverage Duration | Initial, reauth: 12 months |
| Other Criteria | Hereditary orotic aciduria (reauth): Documentation of positive clinical response to Xuriden therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

XYREM (S)

Products Affected

- Xyrem

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Narcolepsy with cataplexy (Narcolepsy Type 1)(initial): Diagnosis of narcolepsy as confirmed by sleep study (unless the prescriber provides justification confirming that a sleep study would not be feasible), AND symptoms of cataplexy are present, AND symptoms of excessive daytime sleepiness (eg, irrepressible need to sleep or daytime lapses into sleep) are present. Narcolepsy without cataplexy (Narcolepsy Type 2)(initial): Diagnosis of narcolepsy as confirmed by sleep study (unless the prescriber provides justification confirming that a sleep study would not be feasible), AND symptoms of cataplexy are absent, AND symptoms of excessive daytime sleepiness (eg, irrepressible need to sleep or daytime lapses into sleep) are present, AND trial and failure, contraindication, or intolerance to one of the following: 1) amphetamine-based stimulant (eg, amphetamine, dextroamphetamine), OR 2) methylphenidate-based stimulant. |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | All uses (initial, reauth): 12 months |
| Other Criteria | Narcolepsy Type 1 (reauth): Documentation demonstrating a reduction in the frequency of cataplexy attacks associated with Xyrem therapy, OR documentation demonstrating a reduction in symptoms of excessive daytime sleepiness associated with Xyrem therapy. Narcolepsy Type 2 (reauth): Documentation demonstrating a reduction in symptoms of excessive daytime sleepiness associated with Xyrem therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

YERVOY (S)

Products Affected

- Yervoy

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | N/A |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

YONSA (S)

Products Affected

- Yonsa

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Prostate Cancer: Diagnosis of metastatic castration-resistant (chemical or surgical) or recurrent prostate cancer. Used in combination with methylprednisolone. Trial and failure or intolerance to Xtandi (enzalutamide). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or urologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ZALTRAP (S)

Products Affected

- Zaltrap

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Colon and/or rectal cancer: Diagnosis of metastatic colon and/or rectal cancer. Ziv-aflibercept is being used in combination with 5-fluorouracil, leucovorin, and irinotecan (FOLFIRI) regimen. Patient has disease that is resistant to or has progressed following an oxaliplatin-containing regimen [e.g., 5-fluorouracil, leucovorin, and oxaliplatin (FOLFOX)]. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ZAVESCA (S)

Products Affected

- Miglustat

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Gaucher disease: Diagnosis of mild to moderate type 1 Gaucher disease. Patient is unable to receive enzyme replacement therapy due to one of the following conditions: allergy or hypersensitivity to enzyme replacement therapy, poor venous access, unavailability of enzyme replacement therapy (e.g., Cerezyme, VPRIV). |
| Age Restrictions | N/A |
| Prescriber Restrictions | N/A |
| Coverage Duration | Gaucher disease: 12 months |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

ZEJULA (S)

Products Affected

- Zejula

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer: Diagnosis of advanced epithelial ovarian cancer, advanced fallopian tube cancer, or advanced primary peritoneal cancer. Used for maintenance treatment in patients who are in a complete or partial response to first-line platinum-based chemotherapy (e.g., cisplatin, carboplatin). Recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer: Diagnosis of recurrent epithelial ovarian cancer, recurrent fallopian tube cancer, or recurrent primary peritoneal cancer. Used for maintenance treatment in patients who are in a complete or partial response to platinum-based chemotherapy (e.g., cisplatin, carboplatin). Treatment of advanced ovarian cancer after three or more chemotherapies: Diagnosis of advanced ovarian cancer, advanced fallopian tube cancer, or advanced primary peritoneal cancer. Patient has been treated with three or more prior chemotherapy regimens. Patient's cancer is associated with homologous recombination deficiency (HRD) positive status defined by one of the following: (a) a deleterious or suspected deleterious BRCA mutation or (b) both of the following: (1) genomic instability and (2) cancer has progressed more than 6 months after response to the last platinum-based chemotherapy (e.g., cisplatin, carboplatin). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ZELBORAF (S)

Products Affected

- Zelboraf

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Melanoma: Diagnosis of unresectable melanoma or metastatic melanoma. Cancer is BRAFV600 mutant type (MT) as detected by a U.S. Food and Drug Administration (FDA)-approved test (eg, cobas 4600 BRAFV600 Mutation Test) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). Erdheim-Chester Disease: Diagnosis of Erdheim-Chester disease AND Disease is BRAFV600 mutant type (MT). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Melanoma: Prescribed by or in consultation with an oncologist. Erdheim-Chester Disease: Prescribed by or in consultation with a hematologist/oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | All indications: Approve for continuation of therapy. |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

ZIEXTENZO (S)

Products Affected

- Ziextenzo

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | <p>Febrile neutropenia (FN) prophylaxis: Patient will be receiving prophylaxis for FN due to one of the following: 1) Patient is receiving National Cancer Institute’s Breast Intergroup, INT C9741 dose dense chemotherapy protocol for primary breast cancer, 2) patient is receiving a dose-dense chemotherapy regimen for which the incidence of FN is unknown, 3) patient is receiving chemotherapy regimen(s) associated with greater than 20% incidence of FN, 4) both of the following: a) patient is receiving chemotherapy regimen(s) associated with 10-20% incidence of FN, AND b) patient has one or more risk factors associated with chemotherapy-induced infection, FN, or neutropenia, OR 5) Both of the following: a) patient is receiving myelosuppressive anticancer drugs associated with neutropenia, AND b) patient has a history of FN or dose-limiting event during a previous course of chemotherapy (secondary prophylaxis). Acute radiation syndrome (ARS): Patient was/will be acutely exposed to myelosuppressive doses of radiation (hematopoietic subsyndrome of ARS). Treatment of FN: Patient has received or is receiving myelosuppressive anticancer drugs associated with neutropenia. Diagnosis of FN. Patient is at high risk for infection-associated complications.</p> |
| Age Restrictions | N/A |
| Prescriber Restrictions | All uses: Prescribed by or in consultation with a hematologist/oncologist |
| Coverage Duration | ARS: 1 mo. FN (prophylaxis, treatment): 3 mo or duration of tx. |
| Other Criteria | All Indications: Trial and failure or intolerance to both of the following: Neulasta/Neulasta Onpro AND Udenyca. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ZOLINZA (S)

Products Affected

- Zolinda

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Cutaneous T-cell lymphoma (CTCL): Diagnosis of CTCL. Progressive, persistent or recurrent disease on or contraindication or intolerance to two systemic therapies (e.g., bexarotene, romidepsin, etc.). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist/oncologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ZORBTIVE (S)

Products Affected

- Zorbtive

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Short Bowel Syndrome (SBS): Diagnosis of SBS. Patient is currently receiving specialized nutritional support (eg, intravenous parenteral nutrition, fluid, and micronutrient supplements). Patient has not previously received 4 weeks of treatment with Zorbtive. |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with a gastroenterologist. |
| Coverage Duration | SBS: 4 weeks. |
| Other Criteria | N/A |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

ZYDELIG (S)

Products Affected

- Zydelig

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Chronic lymphocytic leukemia (CLL): Diagnosis of CLL. Used in combination with Rituxan (rituximab). The patient has relapsed on at least one prior therapy (eg, purine analogues [fludarabine, pentostatin, cladribine], alkylating agents [chlorambucil, cyclophosphamide], or monoclonal antibodies [rituximab]). Patient is a candidate for Rituxan (rituximab) monotherapy due to presence of other comorbidities (eg, coronary artery disease, peripheral vascular disease, diabetes mellitus, pulmonary disease [COPD]). Follicular Lymphoma (FL): Diagnosis of FL. The patient has relapsed on at least two prior systemic therapies (eg, rituximab, alkylating agents [cyclophosphamide, chlorambucil], anthracyclines [doxorubicin, daunorubicine], purine analogs [fludarabine]). Small lymphocytic lymphoma (SLL): Diagnosis of SLL. The patient has relapsed on at least two prior systemic therapies (eg, rituximab, alkylating agents [cyclophosphamide, chlorambucil], anthracyclines [doxorubicin, daunorubicine], purine analogs [fludarabine]). |
| Age Restrictions | N/A |
| Prescriber Restrictions | All uses: Prescribed by or in consultation with an oncologist/hematologist. |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ZYKADIA (S)

Products Affected

- Zykadia

| PA Criteria | Criteria Details |
|------------------------------|--|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Non-small cell lung cancer (NSCLC): Diagnosis of NSCLC that is metastatic or recurrent. Tumor is anaplastic lymphoma kinase (ALK)-positive as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA). |
| Age Restrictions | N/A |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ZYTIGA (NON-PREFERRED) (S)

Products Affected

- Zytiga TABS 500MG

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Metastatic Castration-Resistant Prostate Cancer (mCRPC): Diagnosis of metastatic castration-resistant (chemical or surgical) or recurrent prostate cancer. Used in combination with prednisone. Trial and failure, or intolerance to Xtandi (enzalutamide). Metastatic Castration-Sensitive Prostate Cancer (mCSPC): Diagnosis of metastatic high-risk castration-sensitive prostate cancer. Used in combination with prednisone. |
| Age Restrictions | N/A |
| Prescriber Restrictions | mCRPC, mCSPC: Prescribed by or in consultation with an oncologist or urologist |
| Coverage Duration | mCRPC, mCSPC: 12 months |
| Other Criteria | Approve for continuation of prior therapy. |

Prior Authorization Criteria
Scott and White
Effective: September 1, 2020

ZYTIGA (PREFERRED) (S)

Products Affected

- Abiraterone Acetate

| PA Criteria | Criteria Details |
|------------------------------|---|
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Exclusion Criteria | N/A |
| Required Medical Information | Metastatic Castration-Resistant Prostate Cancer (mCRPC): Diagnosis of metastatic castration-resistant (chemical or surgical) or recurrent prostate cancer. Used in combination with prednisone. Metastatic Castration-Sensitive Prostate Cancer (mCSPC): Diagnosis of metastatic high-risk castration-sensitive prostate cancer. Used in combination with prednisone. |
| Age Restrictions | N/A |
| Prescriber Restrictions | mCRPC, mCSPC: Prescribed by or in consultation with an oncologist or urologist |
| Coverage Duration | mCRPC, mCSPC: 12 months |
| Other Criteria | Approve for continuation of prior therapy |

PART B VERSUS PART D

Products Affected

- Abelcet
- Acetylcysteine SOLN
- Acyclovir Sodium INJ 1000MG, 500MG, 50MG/ML
- Adriamycin INJ 10MG, 2MG/ML, 50MG
- Adrucil INJ 2.5GM/50ML, 500MG/10ML, 5GM/100ML
- Akynzeo CAPS
- Albuterol Sulfate NEBU
- Ambisome
- Amino Acid INJ 50MG/ML; 50MG/ML
- Aminosyn INJ 148MEQ/L; 1280MG/100ML; 980MG/100ML; 1280MG/100ML; 300MG/100ML; 720MG/100ML; 940MG/100ML; 720MG/100ML; 400MG/100ML; 440MG/100ML; 860MG/100ML; 420MG/100ML; 520MG/100ML; 160MG/100ML; 44MG/100ML; 800MG/100ML
- Aminosyn 7%/electrolytes INJ 124MEQ/L; 900MG/100ML; 690MG/100ML; 96MEQ/L; 900MG/100ML; 210MG/100ML; 510MG/100ML; 660MG/100ML; 510MG/100ML; 10MEQ/L; 280MG/100ML; 310MG/100ML; 30MMOLE/L; 65MEQ/L; 610MG/100ML; 300MG/100ML; 65MEQ/L; 370MG/100ML; 120MG/100ML; 44MG/100ML; 560MG/100ML
- Aminosyn 8.5%/electrolytes INJ 142MEQ/L; 1100MG/100ML; 850MG/100ML; 98MEQ/L; 1100MG/100ML; 260MG/100ML; 620MG/100ML; 810MG/100ML; 624MG/100ML; 10MEQ/L; 340MG/100ML; 380MG/100ML; 30MEQ/L; 65MEQ/L; 750MG/100ML; 370MG/100ML; 65MEQ/L; 460MG/100ML; 150MG/100ML; 44MG/100ML; 680MG/100ML
- Aminosyn II INJ 50.3MEQ/L; 695MG/100ML; 713MG/100ML; 490MG/100ML; 517MG/100ML; 350MG/100ML; 210MG/100ML; 462MG/100ML; 700MG/100ML; 735MG/100ML; 120MG/100ML; 209MG/100ML; 505MG/100ML; 371MG/100ML; 31.3MEQ/L; 280MG/100ML; 140MG/100ML; 189MG/100ML; 350MG/100ML; 71.8MEQ/L; 993MG/100ML; 1018MG/100ML; 700MG/100ML; 738MG/100ML; 500MG/100ML; 300MG/100ML; 660MG/100ML; 1000MG/100ML; 1050MG/100ML; 172MG/100ML; 298MG/100ML; 722MG/100ML; 530MG/100ML; 38MEQ/L; 400MG/100ML; 200MG/100ML; 270MG/100ML; 500MG/100ML
- Aminosyn II 8.5%/electrolytes
- Aminosyn M INJ 65MEQ/L; 448MG/100ML; 343MG/100ML; 40MEQ/L; 448MG/100ML; 105MG/100ML; 252MG/100ML; 329MG/100ML; 252MG/100ML; 3MEQ/L; 140MG/100ML; 154MG/100ML; 3.5MMOLE/L; 13MEQ/L; 300MG/100ML; 147MG/100ML; 40MEQ/L; 182MG/100ML; 56MG/100ML; 31MG/100ML; 280MG/100ML

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

- Aminosyn-hbc INJ 7.1MEQ/100ML; 660MG/100ML; 507MG/100ML; 660MG/100ML; 154MG/100ML; 789MG/100ML; 1576MG/100ML; 265MG/100ML; 206MG/100ML; 1.12GM/100ML; 228MG/100ML; 448MG/100ML; 221MG/100ML; 272MG/100ML; 88MG/100ML; 33MG/100ML; 789MG/100ML
- Aminosyn-pf INJ 46MEQ/L; 698MG/100ML; 1227MG/100ML; 527MG/100ML; 820MG/100ML; 385MG/100ML; 312MG/100ML; 760MG/100ML; 1200MG/100ML; 677MG/100ML; 180MG/100ML; 427MG/100ML; 812MG/100ML; 495MG/100ML; 3.4MEQ/L; 70MG/100ML; 512MG/100ML; 180MG/100ML; 44MG/100ML; 673MG/100ML
- Aminosyn-pf 7%
- Aminosyn-rf INJ 113MEQ/L; 600MG/100ML; 429MG/100ML; 462MG/100ML; 726MG/100ML; 535MG/100ML; 726MG/100ML; 726MG/100ML; 330MG/100ML; 165MG/100ML; 528MG/100ML
- Amphotec
- Amphotericin B INJ
- Anzemet TABS
- Aprepitant
- Astagraf XL
- Azasan
- Azathioprine INJ
- Azathioprine TABS
- Baclofen INJ
- Bethkis
- Bleomycin
- Bleomycin Sulfate INJ
- Budesonide SUSP
- Cladribine
- Clinimix 2.75%/dextrose 5%
- Clinimix 4.25%/dextrose 10%
- Clinimix 4.25%/dextrose 20%
- Clinimix 4.25%/dextrose 25%
- Clinimix 5%/dextrose 15%
- Clinimix 5%/dextrose 20%
- Clinimix 5%/dextrose 25%
- Clinimix E 2.75%/dextrose 10% INJ 570MG/100ML; 317MG/100ML; 33MG/100ML; 10GM/100ML; 283MG/100ML; 132MG/100ML; 165MG/100ML; 201MG/100ML; 159MG/100ML; 51MG/100ML; 110MG/100ML; 454MG/100ML; 154MG/100ML; 261MG/100ML; 187MG/100ML; 138MG/100ML; 217MG/100ML; 112MG/100ML; 116MG/100ML; 50MG/100ML; 11MG/100ML; 160MG/100ML
- Clinimix E 2.75%/dextrose 5%
- Clinimix E 4.25%/dextrose 10%
- Clinimix E 4.25%/dextrose 25%
- Clinimix E 4.25%/dextrose 5%
- Clinimix E 5%/dextrose 15%
- Clinimix E 5%/dextrose 20%
- Clinimix E 5%/dextrose 25%
- Clinimix N14g30e
- Clinimix N9g15e
- Clinimix N9g20e
- Clinisol Sf 15%
- Clinolipid
- Clonidine Hydrochloride INJ
- Cromolyn Sodium NEBU
- Cyclophosphamide CAPS
- Cyclosporine CAPS
- Cyclosporine Modified
- Cytarabine INJ 100MG/ML, 20MG/ML
- Cytarabine Aqueous
- Deferoxamine Mesylate
- Dobutamine Hcl INJ 250MG/20ML, 500MG/40ML
- Dobutamine Hcl/d5w INJ 5%; 1MG/ML
- Dobutamine Hydrochloride/dextrose
- Dobutamine/dextrose 5% INJ 5%; 2MG/ML, 5%; 4MG/ML

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

- Dopamine Hcl INJ 160MG/ML, 80MG/ML
- Dopamine Hydrochloride INJ
- Dopamine Hydrochloride/dextrose
- Dopamine/d5w INJ 5%; 3.2MG/ML
- Doxorubicin Hcl INJ 10MG, 2MG/ML, 50MG
- Emend SUSR
- Engerix-b
- Envarsus Xr
- Everolimus TABS 0.25MG, 0.5MG, 0.75MG
- Fentanyl Citrate INJ 1000MCG/20ML, 100MCG/2ML, 2500MCG/50ML, 250MCG/5ML, 500MCG/10ML, 50MCG/ML
- Floxuridine INJ
- Fluorouracil INJ 1GM/20ML, 2.5GM/50ML, 500MG/10ML, 5GM/100ML
- Freamine Hbc 6.9%
- Freamine III INJ 89MEQ/L; 710MG/100ML; 950MG/100ML; 3MEQ/L; 24MG/100ML; 1400MG/100ML; 280MG/100ML; 690MG/100ML; 910MG/100ML; 730MG/100ML; 530MG/100ML; 560MG/100ML; 10MMOLE/L; 120MG/100ML; 1120MG/100ML; 590MG/100ML; 10MEQ/L; 400MG/100ML; 150MG/100ML; 660MG/100ML
- Gablofen
- Ganciclovir INJ 500MG, 500MG/10ML
- Gengraf
- Granisetron Hcl TABS
- Hecoria CAPS 0.5MG, 1MG
- Hepagam B
- Hepatamine
- Heplisav-b
- Hyperhep B S/d
- Hyperrab
- Hyperrab S/d INJ 1500UNIT/10ML, 300UNIT/2ML
- Imogam Rabies-ht
- Imovax Rabies (h.d.c.v.)
- Intralipid INJ 20GM/100ML
- Ipratropium Bromide INHALATION SOLN 0.02%
- Ipratropium Bromide/albuterol Sulfate
- Kedrab
- Leucovorin Calcium INJ 100MG/10ML
- Levalbuterol NEBU
- Levalbuterol Hcl NEBU
- Levalbuterol Hydrochloride
- Lioresal Intrathecal
- Milrinone In Dextrose INJ 5%; 20MG/100ML, 5%; 40MG/200ML
- Milrinone Lactate INJ 10MG/10ML, 20MG/20ML, 50MG/50ML
- Morphine Sulfate INJ 10MG/ML, 150MG/30ML, 1MG/ML, 4MG/ML, 5MG/ML, 8MG/ML
- Mycophenolate Mofetil
- Mycophenolic Acid Dr
- Nabi-hb
- Nebupent
- Nephramine
- Nutrilipid
- Ondansetron Hcl SOLN
- Ondansetron Hcl TABS 24MG
- Ondansetron Hydrochloride TABS
- Ondansetron Odt
- Pentamidine Isethionate INHALATION SOLR
- Perforomist
- Plenamine
- Premasol
- Procalamine
- Prograf PACK
- Prosol
- Rabavert
- Rapamune SOLN
- Recombivax Hb
- Sandimmune SOLN
- Sirolimus SOLN
- Sirolimus TABS

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

- Synthamin 17
- Tacrolimus CAPS
- Tobramycin NEBU
- Travasol INJ 52MEQ/L;
1760MG/100ML; 880MG/100ML;
34MEQ/L; 1760MG/100ML;
372MG/100ML; 406MG/100ML;
526MG/100ML; 492MG/100ML;
492MG/100ML; 526MG/100ML;
356MG/100ML; 500MG/100ML;
356MG/100ML; 390MG/100ML;
34MG/100ML; 152MG/100ML
- Trimethobenzamide Hydrochloride
- Trophamine INJ 97MEQ/L;
0.54GM/100ML; 1.2GM/100ML;
0.32GM/100ML; 0; 0; 0.5GM/100ML;
0.36GM/100ML; 0.48GM/100ML;
0.82GM/100ML; 1.4GM/100ML;
1.2GM/100ML; 0.34GM/100ML;
0.48GM/100ML; 0.68GM/100ML;
0.38GM/100ML; 5MEQ/L;
0.025GM/100ML; 0.42GM/100ML;
0.2GM/100ML; 0.24GM/100ML;
0.78GM/100ML
- Vinblastine Sulfate INJ 1MG/ML
- Vincasar Pfs
- Vincristine Sulfate INJ
- Xopenex
- Xopenex Concentrate
- Yupelri
- Zortress

Details

This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.

INDEX

| | |
|---|-------|
| 7 | |
| 7t Lido Gel | 199 |
| A | |
| Abelcet | 453 |
| Abiraterone Acetate..... | 452 |
| Abstral | 243 |
| Acetylcysteine | 453 |
| Actemra | 2, 3 |
| Actemra Actpen..... | 3 |
| Actemra IV (s)..... | 2 |
| Actemra Sc (s)..... | 3 |
| Acthar | 133 |
| Actimmune | 4 |
| Actimmune (s)..... | 4 |
| Acyclovir Sodium | 453 |
| Adcetris | 5 |
| Adcetris (s) | 5 |
| Adcirca (s)..... | 6 |
| Adderall Xr (s) | 7 |
| Adempas..... | 8 |
| Adempas (s)..... | 8 |
| Adriamycin..... | 453 |
| Adrucil..... | 453 |
| Afinitor | 9 |
| Afinitor (s)..... | 9 |
| Afinitor Disperz..... | 10 |
| Afinitor Disperz (s) | 10 |
| Aimovig..... | 11 |
| Aimovig (s) | 11 |
| Akynzeo | 453 |
| Albuterol Sulfate | 453 |
| Aldurazyme | 12 |
| Aldurazyme (s)..... | 12 |
| Alecensa | 13 |
| Alecensa (s)..... | 13 |
| Aliqopa..... | 14 |
| Aliqopa (s) | 14 |
| Alosetron Hydrochloride | 203 |
| Alpha-1 Proteinase Inhibitor, Non-preferred (s).. | 15 |
| Alpha-1 Proteinase Inhibitor, Prolastin (s) | 16 |
| Alprazolam..... | 35 |
| Alprazolam Er..... | 35 |
| Alprazolam Intensol..... | 35 |
| Alprazolam Odt..... | 35 |
| Alprazolam Xr | 35 |
| Alunbrig | 17 |
| Alunbrig (s)..... | 17 |
| Alyq..... | 6 |
| Ambisome..... | 453 |
| Ambrisentan..... | 195 |
| Amino Acid..... | 453 |
| Aminosyn..... | 453 |
| Aminosyn 7%/electrolytes | 453 |
| Aminosyn 8.5%/electrolytes | 453 |
| Aminosyn II | 453 |
| Aminosyn II 8.5%/electrolytes | 453 |
| Aminosyn M | 453 |
| Aminosyn-hbc..... | 454 |
| Aminosyn-pf | 454 |
| Aminosyn-pf 7% | 454 |
| Aminosyn-rf..... | 454 |
| Amitriptyline Hcl | 147 |
| Amitriptyline Hydrochloride | 147 |
| Amnesteem | 167 |
| Amphetamine (s)..... | 18 |
| Amphetamine/dextroamphetamine | 7, 18 |
| Amphotec | 454 |
| Amphotericin B..... | 454 |
| Ampyra | 19 |
| Ampyra (s) | 19 |
| Anadrol-50 | 20 |
| Anadrol-50 (s)..... | 20 |
| Androderm | 383 |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | | | |
|----------------------------|-----|---|-----|
| Androxy..... | 21 | Berinert (s) | 36 |
| Androxy (s) | 21 | Besponsa | 37 |
| Anzemet | 454 | Besponsa (s) | 37 |
| Apokyn | 22 | Betaseron..... | 230 |
| Apokyn (s)..... | 22 | Bethkis | 454 |
| Aprepitant..... | 454 | Bexarotene | 374 |
| Aralast Np | 15 | Bivigam..... | 170 |
| Aranesp (s) | 23 | Bleomycin..... | 454 |
| Aranesp Albumin Free | 23 | Bleomycin Sulfate..... | 454 |
| Arcalyst | 25 | Blincyto..... | 38 |
| Arcalyst (s) | 25 | Blincyto (s)..... | 38 |
| Armodafinil | 255 | Bortezomib..... | 39 |
| Arzerra..... | 26 | Bortezomib (s) | 39 |
| Arzerra (s) | 26 | Bosentan..... | 392 |
| Asceniv | 170 | Bosulif..... | 40 |
| Ascomp/codeine | 143 | Bosulif (s)..... | 40 |
| Astagraf XL..... | 454 | Botox..... | 41 |
| Aubagio | 27 | Botox (s)..... | 41 |
| Aubagio (s)..... | 27 | Braftovi | 43 |
| Auryxia..... | 28 | Braftovi (s) | 43 |
| Auryxia (s)..... | 28 | Briviact..... | 44 |
| Austedo | 29 | Briviact (s) | 44 |
| Austedo (s) | 29 | Brukinsa | 45 |
| Avita | 391 | Brukinsa (s)..... | 45 |
| Avonex | 230 | Budesonide..... | 454 |
| Avonex Pen | 230 | Butalbital Compound/codeine | 143 |
| Ayvakit..... | 30 | Butalbital/acetaminophen | 143 |
| Ayvakit (s)..... | 30 | Butalbital/acetaminophen/caffeine | 143 |
| Azasan | 454 | Butalbital/acetaminophen/caffeine/codeine | 143 |
| Azathioprine | 454 | Butalbital/aspirin/caffeine..... | 143 |
| | | Butalbital/aspirin/caffeine/codeine | 143 |
| B | | Bynfezia Pen | 46 |
| Baclofen | 454 | Bynfezia (s)..... | 46 |
| Balversa..... | 31 | | |
| Balversa (s)..... | 31 | C | |
| Bavencio..... | 32 | Cablivi..... | 47 |
| Bavencio (s)..... | 32 | Cablivi (s)..... | 47 |
| Beleodaq..... | 33 | Cabometyx | 48 |
| Beleodaq (s)..... | 33 | Cabometyx (s)..... | 48 |
| Benlysta..... | 34 | Calquence..... | 49 |
| Benlysta (s)..... | 34 | Calquence (s) | 49 |
| Benzodiazepines (s)..... | 35 | Capacet..... | 143 |
| Berinert..... | 36 | Caprelsa..... | 50 |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | | | |
|-------------------------------------|-----|------------------------------------|-----|
| Caprelsa (s)..... | 50 | Clinimix E 4.25%/dextrose 5%..... | 454 |
| Carimune Nanofiltered..... | 170 | Clinimix E 5%/dextrose 15%..... | 454 |
| Carisoprodol..... | 51 | Clinimix E 5%/dextrose 20%..... | 454 |
| Carisoprodol (s)..... | 51 | Clinimix E 5%/dextrose 25%..... | 454 |
| Cayston..... | 52 | Clinimix N14g30e..... | 454 |
| Cayston (s)..... | 52 | Clinimix N9g15e..... | 454 |
| Cephadyn..... | 143 | Clinimix N9g20e..... | 454 |
| Cerdelga..... | 53 | Clinisol Sf 15%..... | 454 |
| Cerdelga (s)..... | 53 | Clinolipid..... | 454 |
| Cerezyme..... | 54 | Clonidine Hydrochloride..... | 454 |
| Cerezyme (s)..... | 54 | Clovique..... | 364 |
| Chlordiazepoxide Hcl..... | 35 | Cometriq..... | 62 |
| Chlordiazepoxide Hydrochloride..... | 35 | Cometriq (s)..... | 62 |
| Chlordiazepoxide/amitriptyline..... | 147 | Copiktra..... | 63 |
| Chlorzoxazone..... | 146 | Copiktra (s)..... | 63 |
| Cholbam..... | 55 | Corlanor..... | 64 |
| Cholbam (s)..... | 55 | Corlanor (s)..... | 64 |
| Chorionic Gonadotropin..... | 56 | Cosentyx..... | 65 |
| Chorionic Gonadotropin (s)..... | 56 | Cosentyx (s)..... | 65 |
| Cialis (s)..... | 57 | Cosentyx Sensoready Pen..... | 65 |
| Ciclodan..... | 58 | Cotellic..... | 66 |
| Ciclopirox (s)..... | 58 | Cotellic (s)..... | 66 |
| Ciclopirox Nail Lacquer..... | 58 | Crinone..... | 67 |
| Cimzia..... | 59 | Crinone (s)..... | 67 |
| Cimzia (s)..... | 59 | Cromolyn Sodium..... | 454 |
| Cimzia Starter Kit..... | 59 | Cutaquig..... | 328 |
| Cinryze..... | 61 | Cuvitru..... | 328 |
| Cinryze (s)..... | 61 | Cyclobenzaprine Hydrochloride..... | 146 |
| Cladribine..... | 454 | Cyclophosphamide..... | 454 |
| Claravis..... | 167 | Cyclosporine..... | 454 |
| Clinimix 2.75%/dextrose 5%..... | 454 | Cyclosporine Modified..... | 454 |
| Clinimix 4.25%/dextrose 10%..... | 454 | Cyproheptadine Hcl..... | 141 |
| Clinimix 4.25%/dextrose 20%..... | 454 | Cyproheptadine Hydrochloride..... | 141 |
| Clinimix 4.25%/dextrose 25%..... | 454 | Cyramza..... | 68 |
| Clinimix 4.25%/dextrose 5%..... | 454 | Cyramza (s)..... | 68 |
| Clinimix 5%/dextrose 15%..... | 454 | Cystaran..... | 69 |
| Clinimix 5%/dextrose 20%..... | 454 | Cystaran (s)..... | 69 |
| Clinimix 5%/dextrose 25%..... | 454 | Cytarabine..... | 454 |
| Clinimix E 2.75%/dextrose 10%..... | 454 | Cytarabine Aqueous..... | 454 |
| Clinimix E 2.75%/dextrose 5%..... | 454 | | |
| Clinimix E 4.25%/dextrose 10%..... | 454 | D | |
| Clinimix E 4.25%/dextrose 25%..... | 454 | Dacogen (s)..... | 70 |
| | | Dalfampridine Er..... | 19 |

Prior Authorization Criteria
 Scott and White
 Effective: September 1, 2020

| | |
|---|---------|
| Daliresp | 71 |
| Daliresp (s) | 71 |
| Daraprim..... | 72 |
| Daraprim (s) | 72 |
| Darzalex | 73 |
| Darzalex (s) | 73 |
| Daurismo | 74 |
| Daurismo (s)..... | 74 |
| Decitabine..... | 70 |
| Deferasirox | 75 |
| Deferasirox (s)..... | 75 |
| Deferoxamine Mesylate | 454 |
| Dexchlorpheniramine Maleate | 141 |
| Dexedrine | 77 |
| Dexmethylphenidate (s) | 76 |
| Dexmethylphenidate Hcl..... | 76 |
| Dexmethylphenidate Hcl Er..... | 76 |
| Dexmethylphenidate Hydrochloride | 76 |
| Dextroamphetamine (s)..... | 77 |
| Dextroamphetamine Sulfate..... | 77 |
| Dextroamphetamine Sulfate Er | 77 |
| Diacomit..... | 78 |
| Diacomit (s)..... | 78 |
| Diclofenac Sodium..... | 281 |
| Dobutamine Hcl | 454 |
| Dobutamine Hcl/d5w | 454 |
| Dobutamine Hydrochloride/dextrose | 454 |
| Dobutamine/dextrose 5% | 454 |
| Dopamine Hcl | 455 |
| Dopamine Hydrochloride..... | 455 |
| Dopamine Hydrochloride/dextrose | 455 |
| Dopamine/d5w | 455 |
| Doxepin Hcl | 147 |
| Doxepin Hydrochloride..... | 79, 147 |
| Doxepin Topical (s)..... | 79 |
| Doxorubicin Hcl..... | 455 |
| Dronabinol..... | 214 |
| Duobrii | 80 |
| Duobrii (s) | 80 |
| Dupixent..... | 81 |
| Dupixent (s)..... | 81 |

| | |
|--------------------------------|-----|
| E | |
| Egrifta | 83 |
| Egrifta (s) | 83 |
| Egrifta Sv | 83 |
| Elaprase..... | 84 |
| Elaprase (s)..... | 84 |
| Eligard..... | 85 |
| Eligard (s)..... | 85 |
| Elzonris | 86 |
| Elzonris (s) | 86 |
| Emend | 455 |
| Emflaza | 87 |
| Emflaza (s)..... | 87 |
| Emgality..... | 88 |
| Emgality (s)..... | 88 |
| Empliciti..... | 89 |
| Empliciti (s) | 89 |
| Enbrel | 90 |
| Enbrel (s)..... | 90 |
| Enbrel Mini | 90 |
| Enbrel Sureclick..... | 90 |
| Endari..... | 91 |
| Endari (s)..... | 91 |
| Engerix-b..... | 455 |
| Entyvio..... | 92 |
| Entyvio (s)..... | 92 |
| Envarsus Xr..... | 455 |
| Epclusa..... | 93 |
| Epclusa Non-preferred (s)..... | 93 |
| Epclusa Preferred (s)..... | 94 |
| Epidiolex..... | 95 |
| Epidiolex (s)..... | 95 |
| Epoprostenol (s)..... | 96 |
| Epoprostenol Sodium..... | 96 |
| Erbitux..... | 97 |
| Erbitux (s) | 97 |
| Erivedge | 98 |
| Erivedge (s)..... | 98 |
| Erleada | 99 |
| Erleada (s)..... | 99 |
| Erlotinib Hydrochloride | 373 |
| Esbriet | 100 |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|----------------------|--------|
| Esbriet (s) | 100 |
| Esgic | 143 |
| Estazolam | 35 |
| Evenity | 101 |
| Evenity (s) | 101 |
| Everolimus | 9, 455 |
| Exjade | 75 |
| Exondys 51 | 102 |
| Exondys 51 (s) | 102 |
| Extavia | 230 |
| Eylea | 103 |
| Eylea (s) | 103 |

F

| | |
|--|----------|
| Fabrazyme | 104 |
| Fabrazyme (s) | 104 |
| Farydak | 105 |
| Farydak (s) | 105 |
| Fasenra | 106 |
| Fasenra (s) | 106 |
| Fasenra Pen | 106 |
| Fentanyl (s) | 108 |
| Fentanyl Citrate | 243, 455 |
| Fentanyl Citrate Oral Transmucosal | 108 |
| Ferriprox | 109 |
| Ferriprox (s) | 109 |
| Ferriprox Twice-a-day | 109 |
| Firazyr | 110 |
| Firazyr (s) | 110 |
| Firdapse | 111 |
| Firdapse (s) | 111 |
| Firmagon | 112 |
| Firmagon (s) | 112 |
| Flebogamma Dif | 170 |
| Floxuridine | 455 |
| Fluorouracil | 455 |
| Folotyn | 113 |
| Folotyn (s) | 113 |
| Forteo | 114 |
| Forteo (s) | 114 |
| Freamine Hbc 6.9% | 455 |
| Freamine III | 455 |
| Fulphila | 116 |

| | |
|--------------------|-----|
| Fulphila (s) | 116 |
|--------------------|-----|

G

| | |
|---|-----|
| Gablofen | 455 |
| Galafold | 117 |
| Galafold (s) | 117 |
| Gamastan | 118 |
| Gamastan S/d | 118 |
| Gamastan S/d (s) | 118 |
| Gamifant | 119 |
| Gamifant (s) | 119 |
| Gammagard Liquid | 170 |
| Gammagard S/d Iga Less Than 1mcg/ml | 170 |
| Gammaked | 170 |
| Gammaplex | 170 |
| Gamunex-c | 170 |
| Ganciclovir | 455 |
| Gattex | 120 |
| Gattex (s) | 120 |
| Gazyva | 121 |
| Gazyva (s) | 121 |
| Gengraf | 455 |
| Genotropin | 130 |
| Genotropin Miniquick | 130 |
| Gilenya | 122 |
| Gilenya (s) | 122 |
| Gilotrif | 123 |
| Gilotrif (s) | 123 |
| Glassia | 15 |
| Glatiramer Acetate | 124 |
| Glatiramer Acetate (s) | 124 |
| Glatopa | 124 |
| Gleevec (s) | 125 |
| Glydo | 199 |
| Gocovri | 126 |
| Gocovri (s) | 126 |
| Granisetron Hcl | 455 |
| Growth Hormone, Non-preferred (s) | 127 |
| Growth Hormone, Preferred (s) | 130 |

H

| | |
|---------------------------|-----|
| H.p. Acthar Gel (s) | 133 |
| Haegarda | 135 |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | | | |
|--|----------|--|-----|
| Haegarda (s) | 135 | Icatibant Acetate | 110 |
| Halaven..... | 136 | Iclusig..... | 152 |
| Halaven (s) | 136 | Iclusig (s) | 152 |
| Harvoni..... | 137 | Idhifa..... | 153 |
| Harvoni (s)..... | 137 | Idhifa (s)..... | 153 |
| Hecoria | 455 | Ilaris | 154 |
| Hepagam B..... | 455 | Ilaris (s)..... | 154 |
| Hepatamine..... | 455 | Ilumya | 155 |
| Heplisav-b | 455 | Ilumya (s)..... | 155 |
| Herceptin | 138 | Imatinib Mesylate | 125 |
| Herceptin (s)..... | 138 | Imbruvica | 156 |
| Hetlioz | 139 | Imbruvica (s)..... | 156 |
| Hetlioz (s)..... | 139 | Imfinzi..... | 157 |
| Hizentra | 140 | Imfinzi (s)..... | 157 |
| Hizentra (s)..... | 140 | Imogam Rabies-ht..... | 455 |
| Hrm - Antihistamines | 141 | Imovax Rabies (h.d.c.v.)..... | 455 |
| Hrm - Antipsychotics | 142 | Inbrija..... | 158 |
| Hrm - Butalbital | 143 | Inbrija (s)..... | 158 |
| Hrm - Endocrine..... | 144 | Increlex | 159 |
| Hrm - Phenobarbital, Pentobarbital..... | 145 | Increlex (s) | 159 |
| Hrm - Skeletal Muscle Relaxants..... | 146 | Inflectra | 160 |
| Hrm - Tca | 147 | Inflectra (s)..... | 160 |
| Humatrope..... | 127 | Ingrezza..... | 162 |
| Humatrope Combo Pack | 127 | Ingrezza (s)..... | 162 |
| Humira..... | 148 | Inlyta | 163 |
| Humira (s) | 148 | Inlyta (s) | 163 |
| Humira Pediatric Crohns Disease Starter Pack . | 148 | Inrebic | 164 |
| Humira Pen..... | 148 | Inrebic (s) | 164 |
| Humira Pen-cd/uc/hs Starter | 148 | Intralipid..... | 455 |
| Humira Pen-ps/uv Starter | 148 | Intron A..... | 165 |
| Hydroxyprogesterone (s)..... | 150 | Intron A (s)..... | 165 |
| Hydroxyprogesterone Caproate..... | 150, 213 | Intron A W/diluent..... | 165 |
| Hydroxyzine Hcl | 141 | Ipratropium Bromide | 455 |
| Hydroxyzine Hydrochloride..... | 141 | Ipratropium Bromide/albuterol Sulfate..... | 455 |
| Hydroxyzine Pamoate | 141 | Iressa | 166 |
| Hyperhep B S/d..... | 455 | Iressa (s)..... | 166 |
| Hyperrab..... | 455 | Isotretinoin | 167 |
| Hyperrab S/d | 455 | Isotretinoin (s)..... | 167 |
| Hyqvia | 328 | Istodax..... | 168 |
| I | | Istodax (overfill) | 168 |
| Ibrance..... | 151 | Istodax (s)..... | 168 |
| Ibrance (s)..... | 151 | Isturisa..... | 169 |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|-------------------|-----|
| Isturisa (s)..... | 169 |
| Itraconazole..... | 346 |
| Ivig (s)..... | 170 |

J

| | |
|----------------------|-----|
| Jadenu..... | 75 |
| Jadenu Sprinkle..... | 75 |
| Jakafi..... | 173 |
| Jakafi (s)..... | 173 |
| Jevtana..... | 174 |
| Jevtana (s)..... | 174 |
| Juxtapid..... | 175 |
| Juxtapid (s)..... | 175 |

K

| | |
|------------------------------|-----|
| Kadcyla..... | 176 |
| Kadcyla (s)..... | 176 |
| Kalbitor..... | 177 |
| Kalbitor (s)..... | 177 |
| Kalydeco..... | 178 |
| Kalydeco (s)..... | 178 |
| Kanuma..... | 179 |
| Kanuma (s)..... | 179 |
| Kedrab..... | 455 |
| Keveyis..... | 180 |
| Keveyis (s)..... | 180 |
| Kevzara..... | 181 |
| Kevzara (s)..... | 181 |
| Keytruda..... | 182 |
| Keytruda (s)..... | 182 |
| Kineret..... | 183 |
| Kineret (s)..... | 183 |
| Kisqali..... | 184 |
| Kisqali (s)..... | 184 |
| Kisqali Femara 200 Dose..... | 185 |
| Kisqali Femara 400 Dose..... | 185 |
| Kisqali Femara 600 Dose..... | 185 |
| Kisqali-femara Pack (s)..... | 185 |
| Klofensaid II..... | 281 |
| Korlym..... | 186 |
| Korlym (s)..... | 186 |
| Koselugo..... | 187 |
| Koselugo (s)..... | 187 |

| | |
|--------------------|-----|
| Krystexxa..... | 188 |
| Krystexxa (s)..... | 188 |
| Kuvan..... | 189 |
| Kuvan (s)..... | 189 |
| Kynamro..... | 190 |
| Kynamro (s)..... | 190 |
| Kyprolis..... | 191 |
| Kyprolis (s)..... | 191 |

L

| | |
|--------------------------------------|----------|
| Lartruvo..... | 192 |
| Lartruvo (s)..... | 192 |
| Lazanda..... | 243 |
| Ledipasvir/sofosbuvir..... | 137 |
| Lemtrada..... | 193 |
| Lemtrada (s)..... | 193 |
| Lenvima (s)..... | 194 |
| Lenvima 10 Mg Daily Dose..... | 194 |
| Lenvima 12mg Daily Dose..... | 194 |
| Lenvima 14 Mg Daily Dose..... | 194 |
| Lenvima 18 Mg Daily Dose..... | 194 |
| Lenvima 20 Mg Daily Dose..... | 194 |
| Lenvima 24 Mg Daily Dose..... | 194 |
| Lenvima 4 Mg Daily Dose..... | 194 |
| Lenvima 8 Mg Daily Dose..... | 194 |
| Letairis..... | 195 |
| Letairis (s)..... | 195 |
| Leucovorin Calcium..... | 455 |
| Leukine..... | 196 |
| Leukine (s)..... | 196 |
| Leuprolide Acetate..... | 207 |
| Levalbuterol..... | 455 |
| Levalbuterol Hcl..... | 455 |
| Levalbuterol Hydrochloride..... | 455 |
| Libtayo..... | 198 |
| Libtayo (s)..... | 198 |
| Lidocaine..... | 199, 200 |
| Lidocaine And Tetracaine Cream..... | 199 |
| Lidocaine Hcl..... | 199 |
| Lidocaine Hcl Jelly..... | 199 |
| Lidocaine Topical (s)..... | 199 |
| Lidocaine/prilocaine..... | 199 |
| Lidocaine-prilocaine-cream Base..... | 199 |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | | | |
|---------------------------------|----------|--|----------|
| Lidoderm (s)..... | 200 | Mekinist (s)..... | 218 |
| Lioresal Intrathecal..... | 455 | Mektovi..... | 220 |
| Lonsurf..... | 201 | Mektovi (s)..... | 220 |
| Lonsurf (s)..... | 201 | Metadate Er..... | 221 |
| Lorazepam..... | 35 | Metadate Er-ritalin Sr (s)..... | 221 |
| Lorazepam Intensol..... | 35 | Methitest..... | 226 |
| Lorbrenea..... | 202 | Methocarbamol..... | 146 |
| Lorbrenea (s)..... | 202 | Methotrexate Injection (s)..... | 222 |
| Lotronex (s)..... | 203 | Methylin Chew (s)..... | 223 |
| Lumizyme..... | 204 | Methylphenidate (s)..... | 224 |
| Lumizyme (s)..... | 204 | Methylphenidate Er (s)..... | 225 |
| Lumoxiti..... | 205 | Methylphenidate Hcl Sr..... | 221 |
| Lumoxiti (s)..... | 205 | Methylphenidate Hydrochloride..... | 223, 224 |
| Lupaneta Pack..... | 206 | Methylphenidate Hydrochloride CD..... | 225 |
| Lupaneta Pack (s)..... | 206 | Methylphenidate Hydrochloride Er..... | 221, 225 |
| Lupron (s)..... | 207 | Methylphenidate Hydrochloride Er (la)..... | 225 |
| Lupron Depot (1-month)..... | 208 | Methyltestosterone..... | 226 |
| Lupron Depot (3-month)..... | 208 | Methyltestosterone (s)..... | 226 |
| Lupron Depot (4-month)..... | 208 | Miglustat..... | 443 |
| Lupron Depot (6-month)..... | 208 | Milrinone In Dextrose..... | 455 |
| Lupron Depot (s)..... | 208 | Milrinone Lactate..... | 455 |
| Lupron Depot Ped (s)..... | 209 | Mirvaso..... | 228 |
| Lupron Depot-ped (1-month)..... | 209 | Mirvaso (s)..... | 228 |
| Lupron Depot-ped (3-month)..... | 209 | Mitoxantrone Hcl..... | 245 |
| Lynparza..... | 210, 211 | Modafinil..... | 290 |
| Lynparza (s)..... | 210 | Morphine Sulfate..... | 455 |
| Lynparza Tablet (s)..... | 211 | Mozobil..... | 229 |
| M | | Mozobil (s)..... | 229 |
| Makena..... | 213 | Ms Interferons (non-preferred) (s)..... | 230 |
| Makena (s)..... | 213 | Ms Interferons (preferred) (s)..... | 231 |
| Margesic..... | 143 | Mulpleta..... | 232 |
| Marinol (s)..... | 214 | Mulpleta (s)..... | 232 |
| Marten-tab..... | 143 | Myalept..... | 233 |
| Mavenclad..... | 215 | Myalept (s)..... | 233 |
| Mavenclad (s)..... | 215 | Mycophenolate Mofetil..... | 455 |
| Mavyret..... | 216 | Mycophenolic Acid Dr..... | 455 |
| Mavyret (s)..... | 216 | Mylotarg..... | 234 |
| Mayzent..... | 217 | Mylotarg (s)..... | 234 |
| Mayzent (s)..... | 217 | Myorisan..... | 167 |
| Mayzent Starter Pack..... | 217 | Myozyme..... | 204 |
| Megestrol Acetate..... | 144 | N | |
| Mekinist..... | 218 | Nabi-hb..... | 455 |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | | | |
|---------------------------------------|-----|--------------------------------|----------|
| Naglazyme..... | 235 | Nutropin Aq Nuspin 5..... | 130 |
| Naglazyme (s) | 235 | Nutropin Aq Pen | 130 |
| Naproxen/esomeprazole Magnesium | 414 | Nuvigil (s)..... | 255 |
| Natpara | 236 | O | |
| Natpara (s)..... | 236 | Ocaliva | 256 |
| Nebupent | 455 | Ocaliva (s)..... | 256 |
| Nephramine | 455 | Ocrevus | 257 |
| Nerlynx..... | 237 | Ocrevus (s)..... | 257 |
| Nerlynx (s)..... | 237 | Octagam | 170 |
| Neulasta..... | 238 | Octreotide Acetate | 326 |
| Neulasta (s)..... | 238 | Odomzo..... | 258 |
| Neulasta Onpro Kit..... | 238 | Odomzo (s)..... | 258 |
| Nexavar | 239 | Ofev..... | 259 |
| Nexavar (s) | 239 | Ofev (s) | 259 |
| Nexletol | 240 | Olumiant | 261 |
| Nexletol (s)..... | 240 | Olumiant (s) | 261 |
| Ninlaro..... | 242 | Omnitrope | 127 |
| Ninlaro (s) | 242 | Ondansetron Hcl | 455 |
| Non-preferred Tirf (s)..... | 243 | Ondansetron Hydrochloride..... | 455 |
| Norditropin Flexpro..... | 127 | Ondansetron Odt..... | 455 |
| Norditropin Nordiflex Pen | 127 | Onmel..... | 262 |
| Northera..... | 244 | Onmel (s) | 262 |
| Northera (s) | 244 | Onpattro | 263 |
| Novantrone (s)..... | 245 | Onpattro (s)..... | 263 |
| Novarel | 56 | Opdivo..... | 264 |
| Nplate | 247 | Opdivo (s) | 264 |
| Nplate (s)..... | 247 | Opsumit..... | 265 |
| Nubeqa | 248 | Opsumit (s)..... | 265 |
| Nubeqa (s) | 248 | Orencia | 266, 267 |
| Nucala..... | 249 | Orencia Clickject | 267 |
| Nucala (s) | 249 | Orencia IV (s) | 266 |
| Nuedexta..... | 251 | Orencia Sc (s)..... | 267 |
| Nuedexta (s) | 251 | Orenitram | 268 |
| Nulojix..... | 252 | Orenitram (s)..... | 268 |
| Nulojix (s) | 252 | Oriahnn | 269 |
| Nuplazid | 253 | Oriahnn (s) | 269 |
| Nuplazid (s)..... | 253 | Orilissa | 270 |
| Nurtec | 254 | Orilissa (s)..... | 270 |
| Nurtec (s)..... | 254 | Orkambi | 271, 272 |
| Nutrilipid..... | 455 | Orkambi (s)..... | 271 |
| Nutropin Aq Nuspin 10..... | 130 | Orkambi Granules (s)..... | 272 |
| Nutropin Aq Nuspin 20..... | 130 | Orphenadrine Citrate Er..... | 146 |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | | | |
|---------------------------------|-----|---|-----|
| Osphena..... | 273 | Pliaglis..... | 199 |
| Osphena (s)..... | 273 | Pomalyst..... | 284 |
| Otezla | 274 | Pomalyst (s) | 284 |
| Otezla (s) | 274 | Portrazza | 285 |
| Oxandrin (s)..... | 275 | Portrazza (s) | 285 |
| Oxandrolone | 275 | Poteligeo | 286 |
| Oxazepam..... | 35 | Poteligeo (s) | 286 |
| Oxbryta..... | 276 | Pregnyl W/diluent Benzyl Alcohol/nacl | 56 |
| Oxbryta (s)..... | 276 | Premasol..... | 455 |
| Oxervate | 277 | Premium Lidocaine | 199 |
| Oxervate (s)..... | 277 | Privigen | 170 |
| P | | | |
| Panzyga | 170 | Procalamine..... | 455 |
| Part B Versus Part D | 453 | Procysbi..... | 287 |
| Pegasys..... | 278 | Procysbi (s) | 287 |
| Pegasys (s)..... | 278 | Prograf..... | 455 |
| Pegasys Proclick..... | 278 | Prolastin-c | 16 |
| Pegintron | 279 | Promacta | 288 |
| Peg-intron (s)..... | 279 | Promacta (s) | 288 |
| Peg-intron Redipen..... | 279 | Promethazine Hcl..... | 141 |
| Pemazyre | 280 | Promethazine Hcl Plain..... | 141 |
| Pemazyre (s)..... | 280 | Promethazine Hydrochloride | 141 |
| Pennsaid | 281 | Promethazine Vc Plain..... | 141 |
| Pennsaid (s) | 281 | Promethazine/phenylephrine..... | 141 |
| Pentamidine Isethionate | 455 | Promethegan | 141 |
| Pentobarbital Sodium | 145 | Prosol | 455 |
| Perforomist..... | 455 | Provigil (s) | 290 |
| Perjeta..... | 282 | Pulmozyme | 292 |
| Perjeta (s)..... | 282 | Pulmozyme (s) | 292 |
| Perphenazine/amitriptyline..... | 147 | Pyrimethamine | 72 |
| Phenadoz | 141 | Q | |
| Phenergan | 141 | Qinlock..... | 293 |
| Phenobarbital..... | 145 | Qinlock (s) | 293 |
| Phenobarbital Sodium | 145 | Qualaquin (s)..... | 294 |
| Phrenilin Forte..... | 143 | Quinine Sulfate | 294 |
| Piqray (s) | 283 | R | |
| Piqray 200mg Daily Dose | 283 | Rabavert | 455 |
| Piqray 250mg Daily Dose | 283 | Radicava..... | 295 |
| Piqray 300mg Daily Dose | 283 | Radicava (s) | 295 |
| Plegridy | 230 | Rapamune | 455 |
| Plegridy Starter Pack..... | 230 | Rasuvo..... | 222 |
| Plenammine | 455 | Ravicti..... | 296 |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | | | |
|-------------------------------------|-----|------------------------------------|----------|
| Ravicti (s) | 296 | Rozlytrek (s)..... | 320 |
| Rebif | 231 | Rubraca | 321 |
| Rebif Rebidose | 231 | Rubraca (s) | 321 |
| Rebif Rebidose Titration Pack | 231 | Ruconest..... | 322 |
| Rebif Titration Pack | 231 | Ruconest (s) | 322 |
| Recombivax Hb..... | 455 | Ruzurgi..... | 323 |
| Regranex..... | 297 | Ruzurgi (s) | 323 |
| Regranex (s) | 297 | Rydapt..... | 324 |
| Relexxii | 225 | Rydapt (s)..... | 324 |
| Remicade..... | 298 | S | |
| Remicade (s)..... | 298 | Sabril | 325 |
| Remodulin | 300 | Sabril (s)..... | 325 |
| Remodulin (s)..... | 300 | Saizen | 127 |
| Renflexis..... | 301 | Saizen Click.easy | 127 |
| Renflexis (s) | 301 | Saizenprep Reconstitutionkit | 127 |
| Repatha..... | 303 | Sandimmune | 455 |
| Repatha (s)..... | 303 | Sandostatin (s)..... | 326 |
| Repatha Pushtrex System | 303 | Sandostatin Lar (s) | 327 |
| Repatha Sureclick..... | 303 | Sandostatin Lar Depot..... | 327 |
| Retacrit | 306 | Scig (s) | 328 |
| Retacrit (s)..... | 306 | Secuado | 329 |
| Retevmo | 308 | Secuado (s)..... | 329 |
| Retevmo (s) | 308 | Serostim | 330 |
| Revatio (s) | 309 | Serostim (s)..... | 330 |
| Revatio Injection (s)..... | 310 | Signifor | 331 |
| Revatio Suspension (s)..... | 311 | Signifor (s) | 331 |
| Revcovi..... | 312 | Signifor Lar..... | 332 |
| Revcovi (s) | 312 | Signifor Lar (s)..... | 332 |
| Revlimid | 313 | Siklos..... | 333 |
| Revlimid (s)..... | 313 | Siklos (s) | 333 |
| Reyvow | 314 | Sildenafil..... | 310 |
| Reyvow (s) | 314 | Sildenafil Citrate | 309, 311 |
| Rilutek (s)..... | 315 | Siliq..... | 334 |
| Riluzole | 315 | Siliq (s)..... | 334 |
| Rinvoq..... | 316 | Simponi | 335 |
| Rinvoq (s)..... | 316 | Simponi (s)..... | 335 |
| Rituxan | 317 | Simponi Aria..... | 337 |
| Rituxan (s)..... | 317 | Simponi Aria (s)..... | 337 |
| Rituxan Hycela..... | 319 | Sirolimus | 455 |
| Rituxan Hycela (s)..... | 319 | Skyrizi..... | 338 |
| Romidepsin | 168 | Skyrizi (s)..... | 338 |
| Rozlytrek | 320 | Sofosbuvir/velpatasvir | 94 |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | | | |
|---------------------------|----------|-----------------------------|-------|
| Soliris | 339 | Synribo (s)..... | 363 |
| Soliris (s) | 339 | Synthamin 17 | 456 |
| Somatuline Depot..... | 340 | Syprine (s)..... | 364 |
| Somatuline Depot (s)..... | 340 | T | |
| Somavert..... | 341 | Tabrecta..... | 365 |
| Somavert (s) | 341 | Tabrecta (s) | 365 |
| Sovaldi..... | 342 | Tacrolimus | 456 |
| Sovaldi (s) | 342 | Tadalafil | 6, 57 |
| Spinraza..... | 344 | Tafamidis (s)..... | 366 |
| Spinraza (s)..... | 344 | Tafinlar..... | 367 |
| Sporanox (s) | 346 | Tafinlar (s) | 367 |
| Spravato (s) | 347 | Tagrisso..... | 369 |
| Spravato 56mg Dose | 347 | Tagrisso (s)..... | 369 |
| Spravato 84mg Dose | 347 | Takhzyro | 370 |
| Sprycel..... | 348 | Takhzyro (s) | 370 |
| Sprycel (s) | 348 | Taltz | 371 |
| Stelara..... | 349, 350 | Taltz (s)..... | 371 |
| Stelara (iv) (s)..... | 349 | Talzenna..... | 372 |
| Stelara (s)..... | 350 | Talzenna (s)..... | 372 |
| Stivarga..... | 352 | Tarceva (s) | 373 |
| Stivarga (s) | 352 | Targretin..... | 374 |
| Strensiq..... | 353 | Targretin (s) | 374 |
| Strensiq (s) | 353 | Tasigna..... | 375 |
| Striant | 383 | Tasigna (s)..... | 375 |
| Supprelin La | 354 | Tavalisse | 376 |
| Supprelin La (s)..... | 354 | Tavalisse (s) | 376 |
| Sutent..... | 355 | Tazverik | 377 |
| Sutent (s) | 355 | Tazverik (s)..... | 377 |
| Sylatron | 356 | Tecentriq | 378 |
| Sylatron (s) | 356 | Tecentriq (s)..... | 378 |
| Sylvant..... | 357 | Tecfidera | 379 |
| Sylvant (s) | 357 | Tecfidera (s) | 379 |
| Symdeko..... | 358 | Tecfidera Starter Pack..... | 379 |
| Symdeko (s) | 358 | Tegsedi..... | 380 |
| Symlin (s) | 359 | Tegsedi (s)..... | 380 |
| Symlinpen 120 | 359 | Temazepam..... | 35 |
| Symlinpen 60 | 359 | Tencon..... | 143 |
| Synagis | 360 | Teriparatide | 381 |
| Synagis (s)..... | 360 | Teriparatide (s)..... | 381 |
| Syndros..... | 362 | Testosterone | 383 |
| Syndros (s) | 362 | Testosterone (s)..... | 383 |
| Synribo | 363 | Testosterone Cypionate..... | 383 |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | | | |
|---------------------------------------|-----|------------------------------|-----|
| Testosterone Enanthate | 385 | Tysabri | 401 |
| Testosterone Enanthate (s) | 385 | Tysabri (s) | 401 |
| Testosterone Pump | 383 | <i>U</i> | |
| Testosterone Topical Solution..... | 383 | Ubrelvy | 403 |
| Tetrabenazine | 427 | Ubrelvy (s) | 403 |
| Thalomid | 387 | Udenyca | 404 |
| Thalomid (s) | 387 | Udenyca (s) | 404 |
| Thioridazine Hcl..... | 142 | Ultomiris | 405 |
| Tibsovo..... | 388 | Ultomiris (s) | 405 |
| Tibsovo (s) | 388 | Uptravi | 406 |
| Tiglutik | 389 | Uptravi (s) | 406 |
| Tiglutik (s)..... | 389 | <i>V</i> | |
| Tobramycin | 456 | Valchlor..... | 407 |
| Tolsura..... | 390 | Valchlor (s) | 407 |
| Tolsura (s) | 390 | Vanadom | 51 |
| Topical Retinoid (s)..... | 391 | Vanatol Lq | 143 |
| Tracleer (s) | 392 | Varizig..... | 408 |
| Travasol..... | 456 | Varizig (s) | 408 |
| Trelstar | 393 | Velcade | 409 |
| Trelstar (s) | 393 | Velcade (s) | 409 |
| Trelstar Mixject..... | 393 | Veletri | 96 |
| Tremfya | 394 | Venclexta | 410 |
| Tremfya (s)..... | 394 | Venclexta (s) | 410 |
| Treprostinil | 300 | Venclexta Starting Pack..... | 410 |
| Tretinoin | 391 | Ventavis | 411 |
| Tretinoin Microsphere..... | 391 | Ventavis (s) | 411 |
| Tretinoin Microsphere Pump | 391 | Verzenio..... | 412 |
| Trientine Hydrochloride..... | 364 | Verzenio (s)..... | 412 |
| Trikafta | 395 | Vigabatrin | 325 |
| Trikafta (s)..... | 395 | Vigadrone..... | 325 |
| Trimethobenzamide Hydrochloride | 456 | Vimizim | 413 |
| Triptodur..... | 396 | Vimizim (s) | 413 |
| Triptodur (s) | 396 | Vimovo (s) | 414 |
| Trophamine | 456 | Vinblastine Sulfate..... | 456 |
| Tukysa | 397 | Vincasar Pfs | 456 |
| Tukysa (s)..... | 397 | Vincristine Sulfate | 456 |
| Turalio | 398 | Vitrakvi | 415 |
| Turalio (s)..... | 398 | Vitrakvi (s) | 415 |
| Tykerb | 399 | Vizimpro | 416 |
| Tykerb (s) | 399 | Vizimpro (s) | 416 |
| Tymlos..... | 400 | Vosevi | 417 |
| Tymlos (s) | 400 | | |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

| | |
|--------------------|-----|
| Vosevi (s) | 417 |
| Votrient..... | 418 |
| Votrient (s) | 418 |
| Vpriv | 419 |
| Vpriv (s) | 419 |
| Vtol Lq | 143 |
| Vumerity..... | 420 |
| Vumerity (s) | 420 |
| Vyndamax | 366 |
| Vyndaqel | 366 |
| Vyxeos..... | 421 |
| Vyxeos (s) | 421 |

W

| | |
|----------------|-----|
| Wakix | 422 |
| Wakix (s)..... | 422 |

X

| | |
|---------------------------|-----|
| Xalkori..... | 423 |
| Xalkori (s) | 423 |
| Xcopri..... | 424 |
| Xcopri (s)..... | 424 |
| Xeljanz | 425 |
| Xeljanz (s) | 425 |
| Xeljanz Xr..... | 425 |
| Xembify..... | 328 |
| Xenazine (s)..... | 427 |
| Xeomin | 428 |
| Xeomin (s)..... | 428 |
| Xermelo..... | 429 |
| Xermelo (s)..... | 429 |
| Xgeva | 430 |
| Xgeva (s) | 430 |
| Xiaflex..... | 431 |
| Xiaflex (s)..... | 431 |
| Xifaxan | 432 |
| Xifaxan (s)..... | 432 |
| Xolair..... | 433 |
| Xolair (s) | 433 |
| Xopenex | 456 |
| Xopenex Concentrate | 456 |
| Xospata..... | 435 |
| Xospata (s)..... | 435 |

| | |
|---------------------------------|-----|
| Xpovio (s) | 436 |
| Xpovio 100 Mg Once Weekly | 436 |
| Xpovio 40 Mg Once Weekly | 436 |
| Xpovio 40 Mg Twice Weekly..... | 436 |
| Xpovio 60 Mg Once Weekly | 436 |
| Xpovio 60 Mg Twice Weekly..... | 436 |
| Xpovio 80 Mg Once Weekly | 436 |
| Xpovio 80 Mg Twice Weekly..... | 436 |
| Xtandi..... | 437 |
| Xtandi (s) | 437 |
| Xuriden | 438 |
| Xuriden (s) | 438 |
| Xyrem | 439 |
| Xyrem (s) | 439 |

Y

| | |
|------------------|-----|
| Yervoy..... | 440 |
| Yervoy (s) | 440 |
| Yonsa | 441 |
| Yonsa (s)..... | 441 |
| Yupelri | 456 |

Z

| | |
|---------------------|-----|
| Zaltrap..... | 442 |
| Zaltrap (s)..... | 442 |
| Zavesca (s) | 443 |
| Zebutal | 143 |
| Zejula | 444 |
| Zejula (s)..... | 444 |
| Zelboraf..... | 445 |
| Zelboraf (s)..... | 445 |
| Zemaira | 15 |
| Zenatane | 167 |
| Zenzedi..... | 77 |
| Ziextenzo..... | 446 |
| Ziextenzo (s) | 446 |
| Zolinza | 447 |
| Zolinza (s) | 447 |
| Zorbtive..... | 448 |
| Zorbtive (s)..... | 448 |
| Zortress | 456 |
| Zydelig | 449 |
| Zydelig (s)..... | 449 |

Prior Authorization Criteria

Scott and White

Effective: September 1, 2020

Zykadia..... 450
Zykadia (s)..... 450
Zytiga 451

Zytiga (non-preferred) (s)451
Zytiga (preferred) (s).....452